

PEDIATRIC NUTRITION

Product Guide 2022

Contents

RESOURCES 5
MEDICAL INFORMATION
Contents by Product Usage
TERM INFANT FORMULAS
Similac® 360 Total Care®. 28 Similac® 360 Total Care® Sensitive. 46 Similac Pro-Advance®. 30 Similac® Advance®. 32 Pure Bliss™ by Similac®. 34 Similac® Organic. 36 Similac® Organic with A2 Milk. 38 Similac® Advance® Kosher Cholov Yisroel. 40 Similac® Advance® Lamehadrin. 42 Similac Soy Isomil®. 44 Similac Pro-Sensitive®. 50 Similac Pro-Total Comfort®. 52 Similac Total Comfort®. 54 Similac® For Spit-Up® Non-GMO 58

EleCare® (for infants)	
HUMAN MILK FORTIFICATION	
Nutrient Composition of Preterm Human Milk	. 67
Concentrated Liquid	. 70
Similac® Human Milk Fortifier Concentrated Liquid	. 72
Nutrient Comparison of Preterm Human Milk + Similac® Special Care® 30	. 75



PREMATURE INFANT FORTIFIERS & FORMULAS	CHILDREN
Similac® Special Care® 20 82 Similac® Special Care® 24 84 Similac® Special Care® 24 High Protein 86 Similac® Special Care® 30 88 Similac® Liqui-Mix® System 90 Similac® Liqui-Mix® System, SSC 20 + SSC 24 92 Similac® Liqui-Mix® System, SSC 20 + SSC 24 HP 95 Similac® Liqui-Mix® System, SSC 20 + SSC 30 96 Similac® Liqui-Mix® System, SSC 24 + SSC 30 97 Similac® Liqui-Mix® System, SSC 24 HP + SSC 30 98 Similac® NeoSure® 100	EleCare® Jr 12 PediaSure® Grow & Gain 12 PediaSure® Grow & Gain Shake Mix 12 PediaSure® Grow & Gain with Fiber 13 PediaSure Harvest™ 13 PediaSure® Reduced Calorie 13 PediaSure SideKicks® 13 PediaSure® Enteral Formula 1.0 Cal 13 PediaSure® Enteral Formula 1.0 Cal with Fiber 14 PediaSure® 1.5 Cal 14 PediaSure® 1.5 Cal with Fiber 14
DIETARY SUPPLEMENT	PediaSure® Peptide 1.0 Cal </td
Similac® Probiotic Tri-Blend	Vital® Peptide 1.5 Cal
Description 108 Custom Feeding System 110 Similac® With Iron 24 112 Pedialyte® 114 Similac® 5% Glucose Water 115	Go & Grow by Similac® Non-GMO with 2'-FL HMO Toddler Drink
Similac 3% Glucose Water	THERAPEUTIC HYDRATION
Similac® Water (Sterilized) 116 Nipples 117 Similac® CustomFeed™ Breast Milk Storage Bottle 118 Bottles & Caps 120 Volu-Feed® 121	Pedialyte® Classic

edialyte® Sport	174 176
METABOLIC PRODUCTS	
alcilo XD®	
yclinex®-1	
lutarex®-1	
lutarex®-2	
ominex®-1	192
ominex®-2	194
Valex®-1	196
Valex®-2	198
etonex®-1	200
etonex®-2	202
henex®-1	204
henex®-2	206
ro-Phree®	208
ropimex®-1	210
ropimex®-2	212
roViMin®	214
CF^{g}	216
/rex®-1	218

ADDITIONAL INFORMATION

Newborn Metric Conversion Tables	
Pediatric Metric Conversion Tables	
Abbott Infant Products: Approximate Powder	
Container Yield	
Macronutrient Values for Pediatric Formula Powders 230	
Pediatric Product HCPCS and NDC-format Codes 234	

Every effort is made to ensure that all product information is correct at the time of publication. Values listed are subject to change. Please refer to the product label or packaging for the most current ingredient, allergen, and nutrient profile information.

Values listed in the Nutrition Information tables are for liquid products except as noted. Values per liter are calculated from values per 100 Cal. No entry indicates value is not available.

Potential Renal Solute Load (PRSL) PRSL = [Protein (g) \times 5.714] + Na (mOsm) + K (mOsm) + CI (mOsm) + P (mOsm)



Customer Service and Product Support

For customer assistance and general information, call our toll-free number (800) 227-5767, Monday – Friday, 8:30 a.m. – 5:00 p.m. EST. You may also submit questions or comments on-line at www.AbbottNutrition.com/contact-us. The Abbott Consumer Relations Department is staffed with representatives who can address your specific questions.

Nutritional Products

Abbott Consumer Relations 2900 Easton Square PI Columbus, OH 43219 (800) 227-5767

Metabolic Products (Inherited Metabolic Disorders)

For healthcare professionals with nutrition questions: Abbott Metabolics Hotline (800) 986-8755

E-business Customers

E-business Department 625 Cleveland Avenue Columbus, Ohio 43215-1724 1-877-4Abbott

Feeding Expert

By phone: (800) 986-8800 Online: www.FeedingExpert.com

How To Order Abbott Products

Hospitals/Institutions

Hospitals and other institutions can order Abbott products by calling toll-free, (800) 551-5838, Monday – Friday, 8:00 a.m. – 5:30 p.m. EST.

On-line

Information about Abbott products can be found at:

- www.AbbottNutrition.com for healthcare professionals
- www.Similac.com

e-Abbott

Connecting our customers to the products they need. Anytime, anyplace, and anywhere.

www.e-abbott.com offers customer support and product support:

- Orders and invoice management
- Chargeback / sales trace
- Electronic funds transfer (EFT)

Store Locator

Our Store Locator can help you find major retail stores by ZIP code that carry the specific Abbott products you would like to purchase.

• On www.AbbottNutrition.com, click on the Store Locator link

Pharmacist

Abbott products may be ordered through a pharmacist.

Locate a Representative

If you would like to speak with an Abbott sales representative, please contact us at (800) 551-5838. Let us know your area of specialization and the product you are inquiring about, and our product experts will be happy to assist you.

Home Delivery

Order at www.abbottstore.com for the convenience of having products delivered to your door directly from the manufacturer.

- Free standard ground shipping on orders over \$35
- Save 10%* when you set up repeat orders with Schedule & Save
- · Orders can be shipped to military addresses
- Orders are processed Monday-Friday and are delivered within 3-5 business days
- Priority orders are delivered within 2-3 business days; overnight orders within 1-2 business days

If you don't have Internet access, call (800) 258-7677 to place an order.[†]

Abbott

Information helpful to healthcare professionals and consumers regarding pediatric nutritional products can be found at www.AbbottNutrition.com:

- Specific product information
- Frequently asked questions
- Information specific to very low-birth-weight babies
- · Breastfeeding support information
- Third-party coverage information, including HCPCS product codes and NDC format codes
- Store locater
- Current, science-based nutrition information, including patient education materials and professional enrichment opportunities

^{*}Discount excludes Nepro, Suplena, & EleCare.

AbbottStore

To order Abbott products, see www.AbbottStore.com for:

- Infant nutrition products
- Child nutrition products
- Delivery information
- Schedule & Save program



Similac® Rewards

Designed for new parents who are looking for information on pregnancy, parenting, and infant nutrition, www.Similac.com contains:

- Up to \$400* in benefits with membership
- Ongoing tips and ideas about all things baby
- Email updates including helpful guides, tips, and nutritional information tailored specifically to baby's growth
- Personalized gifts and benefits customized for your specific needs

Merchandise available at select participating OB offices and hospitals only.

* Offers may vary.



Feeding Expert

Abbott has created quick and easy access to experts available to answer feeding-related questions:

- Helpful information on breastfeeding, supplementing, formula feeding, and fussiness
- Tips for introducing solids to toddlers
- Tools such as Feeding Tracker, Tummy Trouble, and Diaper Decoder
- The Similac® Baby Journal App

Live help with feeding questions; bilingual services (English/ Spanish) are available

- By phone
 - 800-986-8800 (Similac and Prenatal nutrition)
 - 800-986-8793 (PediaSure® picky eaters nutrition)



WebNova[™] Neonutrition Optimizer

- A Web-based program to help manage the enteral nutrition of infants in the NICU
- Stores nutrient profiles for commercial and customized feedings
- Compares nutrient profiles of feedings to each other or to a reference



Abbott Health Institute

At Abbott Health Institute (ANHI), we believe nutrition changes lives. Our goal is to improve health worldwide by connecting and empowering healthcare professionals and patients through science-based nutrition education and resources.

We offer:

- Free, accredited continuing education
- Videos that give practical advice on applying nutrition to practice
- · Short podcasts on focused topics
- Printable materials—like infographics and toolkits on nutrition topics that matter

Explore what we have to offer. Visit www.ANHI.org today.



Abbott Pathway Plus Reimbursement Support

Navigating the insurance coverage process for healthcare expenses can be challenging and time-consuming.

That's why Pathway Plus has a dedicated team offering personalized support to verify insurance coverage for Abbott products when they are recommended by an HCP for a medical need. Visit the Pathway Plus website at www.pathway-plus.com for resources and support, or call 1-855-217-0698.







MEDICAL INFORMATION

Contents by Product Usage

Not a complete listing of all possible products/uses.

Allergy to Cow's Milk Protein EleCare® (for infants) EleCare® Jr Similac® Alimentum® Similac® Soy Isomil®	124 56
łuman Milk Alternative/Supplemental F	ormulas
Pure Bliss [™] by Similac [®]	34
Similac® 360 Total Care®	28
Similac® 360 Total Care® Sensitive	46
Similac® Advance®	32
Similac® Advance® Kosher Cholov Yisroel	
Similac® Advance® Lamehadrin	42
Similac® Alimentum®	56
Similac® For Spit-Up Non-GMO Similac® NeoSure®	58
Similac NeoSure	
Similac® Organic with A2 Milk	38
Similac Pro-Advance®	30
Similac Pro-Sensitive®	48
Similac Pro-Total Comfort®	52
Similac Sensitive®	50
Similac Total Comfort®	54
Similac® Soy Isomil®	44
Similac® With Iron 24	112
luman Milk Fortification	
Liquid Protein Fortifier	78
Similac® Human Milk Fortifiers 70	0, 72, 78
Similac® Special Care® 30	88

Celiac Disease

All Abbott pediatric products are gluten-free.

Clear Liquid Pedialyte Products114, 164-178
Colic Symptoms (due to protein sensitivity) Similac® Alimentum®
Colitis EleCare® (for infants) 60 EleCare® Jr 124 Similac® Alimentum® 56 PediaSure® Peptide 1.0 Cal 146 PediaSure® Peptide 1.5 Cal 148 Vital® Peptide 1.5 Cal 150
Dehydration/Oral Hydration Pedialyte Products114, 164-178
Diarrhea 60 EleCare® (for infants) 60 EleCare® Jr. 124 Pedialyte Products 114, 164-178 PediaSure® Peptide 1.0 Cal 146 PediaSure® Peptide 1.5 Cal 148 Vital® Peptide 1.5 Cal 150 Similac® Alimentum® 56
Disaccharidase Deficiencies ProViMin®

Fat Intolerance/Malabsorption EleCare® (for infants)	124 146 148 214
Food Allergy EleCare® (for infants) EleCare® Jr Similac® Alimentum®	124
Fussiness/Gas/Spit-up Similac® 360 Total Care®	46 56 48 52 50
Galactosemia EleCare® (for infants) EleCare® Jr Similac® Soy Isomil®	124
Gastrointestinal Health Similac® Probiotic Tri-Blend	104

Gluten Intolerance/Sensitivity

All Abbott pediatric products are gluten-free.

Halal

Most Abbott products are certified Halal. Please see www.ifanca.org for more information.

Calcilo XD®	182
Cyclinex®-1	184
Cyclinex®-2	186
EleCare® (for infants)	
EleCare® Jr	
Glutarex®-1	
Glutarex®-2	190
Go & Grow by Similac® Non-GMO with 2'-FL HMO	
Toddler Drink	156
Go & Grow by Similac® Toddler Drink	158
Go & Grow by Similac® Non-GMO Sensitive with 2'-FL	
HMO Toddler Drink	160
Hominex®-1	192
Hominex®-2	
I-Valex®-1	
I-Valex®-2	198
Ketonex®-1	
Ketonex®-2	
Pedialyte® Classic	164
Pedialyte AdvancedCare®	
Pedialyte® Electrolyte Water	168
Pedialyte® with Immune Support Powder	170
Pedialyte® Sport	172
Pedialyte® Freezer Pops	
Pedialyte AdvancedCare® Plus	
PediaSure® Grow & Gain	
PediaSure® Grow & Gain Shake Mix	
PediaSure® Grow & Gain with Fiber	
PediaSure Harvest™	
PediaSure® Reduced Calorie	
PediaSure SideKicks®	
PediaSure® Enteral Formula 1.0 Cal	
PediaSure® Enteral Formula 1.0 Cal with Fiber	140

PediaSure® 1.5 Cal	
PediaSure® 1.5 Cal with Fiber	144
Phenex®-1	204
Phenex®-2	206
Pro-Phree®	208
Propimex®-1	210
Propimex®-2	
ProViMin®	214
Pure Bliss® by Similac®	34
Pure Bliss® by Similac® Toddler Drink	154
RCF®	216
Similac® 360 Total Care®	28
Similac® 360 Total Care® Sensitive	46
Similac® 5% Glucose Water	115
Similac® 10% Glucose Water	115
Similac® Advance®	32
Similac® For Spit-Up Non-GMO	58
Similac® Human Milk Fortifiers	. 70, 72, 78
Similac® Advance® Lamehadrin	42
Similac® NeoSure®	100
Similac® Organic	36
Similac® Organic with A2 Milk	38
Similac® Organic Toddler Drink with A2 Milk	152
Similac® PM 60/40	62
Similac Pro-Advance®	30
Similac Pro-Sensitive®	48
Similac Pro-Total Comfort®	52
Similac Sensitive®	50
Similac® Soy Isomil®	44
Similac® Special Care® 20	82
Similac® Special Care® 24	84
Similac® Special Care® 24 High Protein	86
Similac® Special Care® 30	88
Similac Total Comfort®	54
Similac® Water (Sterilized)	
Similac® With Iron 24	
Tyrex®-1	218
Tyrex®-2	
,	

Hypercalcemia Calcilo XD® Similac® PM 60/40	182 62
Hypoallergenic EleCare® (for infants) EleCare® Jr Similac® Alimentum®	124
nborn Errors of Metabolism	
Calcilo XD® Cyclinex®-1 Cyclinex®-2 Glutarex®-1 Glutarex®-2 Hominex®-2 Hominex®-2 I-Valex®-1 I-Valex®-1 Ketonex®-1 Ketonex®-2 Phenex®-1 Phenex®-2 Pro-Phree® Propimex®-1 Propimex®-1 Propimex®-2 ProViMin® RCF® Tyrex®-1 Tyrex®-1 Tyrex®-2	

Kosher

Most Abbott products are certified Kosher by ①-D on packaging. Please see www.oukosher.org for more information.

Calcilo XD®	
Go & Grow by Similac® Non-GMO with 2'-FL HMO	
Toddler Drink	
Go & Grow by Similac® Toddler Drink	
Go & Grow by Similac® Sensitive Non-GMO with 2'-FL	
HMO Toddler Drink	
EleCare® (for infants)	
EleCare® Jr124	
Pedialyte® Classic	
Pedialyte AdvancedCare®	
Pedialyte® Electrolyte Water	
Pedialyte® with Immune Support Powder	
Pedialyte® Sport	
Pedialyte® Freezer Pops	
Pedialyte AdvancedCare® Plus178	
Pedialyte Products	
ProViMin® 214	
Pure Bliss® by Similac® 34	
Pure Bliss® by Similac® Toddler Drink	
RCF®	
Similac® 360 Total Care®	
Similac® 360 Total Care® Sensitive	
Similac® 5% Glucose Water	
Similac® 10% Glucose Water 115	
Similac® Advance®	
Similac® Advance® Kosher Cholov Yisroel	
Similac® Advance® Lamehadrin42	
Similac® For Spit-Up Non-GMO58	
PediaSure® Grow & Gain with Fiber126	
Similac® Human Milk Fortifiers	
Similac® NeoSure® 100	
Similac® Organic	
Similac® Organic with A2 Milk	
Similac® Organic Toddler Drink with A2 Milk152	
Similac® PM 60/40 62	
Similac Pro-Advance®	
Similac Pro-Sensitive®	
Similac Pro-Total Comfort®	

	Similac Sensitive®	44 82 84 86 88
N	on-GMO*	
	Cyclinex®-1	184
	Cyclinex®-2	
	Go & Grow by Similac® Non-GMO with 2'-FL HMO	
	Toddler Drink	156
	Go & Grow by Similac® Sensitive Non-GMO with 2'-F	=1
	Go & Grow by Similac® Sensitive Non-GMO with 2'-F HMO Toddler Drink	160
	Glutarex®-1	188
	Glutarex®-2	
	Hominex®-1	
	Hominex®-2	
	I-Valex®-1	
	I-Valex®-2	
	Ketonex®-1	200
	Ketonex®-2	202
	PediaSure® Grow & Gain	126
	PediaSure® Grow & Gain Shake Mix	128
	PediaSure® Grow & Gain with Fiber	130
	PediaSure Harvest™	132
	Phenex®-1	204
	Phenex®-2	206
	Pro-Phree®	
	Propimex®-1	
	Propimex®-2	212
	Pure Bliss® by Similac®	34
	Pure Bliss® by Similac® Toddler Drink Similac® 360 Total Care®	154
	Similac® 360 Total Care®	28
	Similac® 360 Total Care® Sensitive	46

Similac® Advance® Lamehadrin 42 Similac® For Spit-Up Non-GMO .58 Similac® Organic .36 Similac® Organic with A2 Milk .38 Similac® Organic Toddler Drink with A2 Milk .152 Similac Pro-Advance® .30 Similac Pro-Sensitive® .48	Protein Maldigestion 60 EleCare® (for infants)
Similac Pro-Total Comfort® 52 Tyrex®-1 218	Renal Dysfunction Similac® PM 60/40
Tyrex®-2	
*Ingredients not genetically engineered. uitable for Lactose Sensitivity/Intolerance	Seizures Managed with Ketogenic Diet RCF®216
	Toddlers/Children
EleCare® (for infants)	EleCare® Jr124
Go & Grow by Similac® Sensitive Non-GMO with 2'-FL	Go & Grow by Similac® Non-GMO with
HMO Toddler Drink	2'-FL HMO Toddler Drink156
Pedialyte Products	Go & Grow by Similac® Toddler Drink
Similac® 360 Total Care® Sensitive	Go & Grow by Similac® Sensitive Non-GMO with 2'-FL
Similac® Alimentum®	HMO Toddler Drink160
Similac® For Spit-Up Non-GMO	Pedialyte® Classic
Similac Pro-Sensitive®	Pedialyte AdvancedCare®
Similac Pro-Total Comfort®	Pedialyte® with Immune Support Powder
Similac Sensitive®	Pedialyte® Freezer Pops
Similac® Soy Isomil®	Pedialyte AdvancedCare® Plus
Similac Total Comfort®	Pedialyte Products114, 164-178
	Pure Bliss® by Similac® Toddler Drink154
remature/Low-Birth-Weight Infants	Similac® Organic Toddler Drink with A2 Milk152
Liquid Protein Fortifier78	
Similac® Human Milk Fortifiers	
Similac® NeoSure® 100	

 Similac
 Special Care® 20
 82

 Similac® Special Care® 24
 84

 Similac® Special Care® 24 High Protein
 86

 Similac® Special Care® 30
 88

Similac® Probiotic Tri-Blend Dietary Supplement....... 104

Recommended Storage Guidelines for Abbott Products

 Store unopened Abbott Nutrition pediatric products (including infant formulas) between 32° F and

95° F; the most desirable range of unopened containers is 55°-75° F. Storage at these temperatures will assure the highest quality product, both aesthetically and nutritionally

- Prolonged exposure to temperatures below 32° F or above 95° F could affect the physical consistency of the product and is not recommended
- Abbott pediatric formula products should not be warmed and then refrigerated
- Do not freeze Abbott pediatric formulas; freezing can render liquid products unusable
- Store Abbott pediatric **powder** products in a dry, cool area. Do not store the actual container of powder in the refrigerator
- Use **powdered** formula within 1 month after opening
- Pour prepared formula into individual feeding bottles or cups, cap, and store in the refrigerator, Refer to product label for appropriate storage times
- Do not leave prepared formula unrefrigerated
- · Once oral feeding begins, use formula within 1 hour or discard
- Do not reuse prefilled feeding bottles
- Keep unopened packets of Similac® Human Milk Fortifier Concentrated Liquids at room temperature in the protective carton; avoid extreme temperatures
- Excludes Similac Probiotic Tri-blend. Please see page 109.

Hangtime for Abbott Products

In each health care facility, it is important to involve the appropriate professionals in developing procedures for using safe techniques in the preparation and use of tube feedings and in setting quality assurance protocols for monitoring those techniques.

The Academy of Nutrition and Dietetics (formerly American Dietetic Association) suggests a hangtime of 4 to 8 hours is acceptable for commercially sterile ready-to-feed products when carefully poured from the container into a tubefeeding setup. This includes Ready To Feed infant formulas, Pedialyte®, and PediaSure® products.

Any feedings that are reconstituted with water or modified in any way should be prepared using aseptic technique and should hang for no more than 4 hours. This includes concentrated liquid and powder formulas, fortified human milk, and any feedings to which other ingredients are added.

Adapted from:

Pediatric Nutrition Practice Group. Steele C, Collins E, eds. Infant and Pediatric Feedings: Guidelines for Preparation of Human Milk and Formula in Health Care Facilities. 3rd ed. Chicago, IL: Academy of Nutrition and Dietetics: 2019:17.

Campbell SM: Preventing Microbial Contamination of Enteral Formulas and Delivery Systems. Columbus, Ohio: Ross Products Division, Abbott Laboratories, 2003.

Bankhead R. Boullata J. Brantlev S. et al. A.S.P.E.N. Enteral Nutrition Practice Recommendations, JPEN, 2009;33:122-167.

Expiration Dates

The expiration date of Abbott products is the first day of the month listed on the product.

- Buy and use the formula by the date shown on the product container
- The vitamin content and the physical stability of the product cannot be guaranteed beyond the "Use By" date because both may degrade with time

WARNINGS

- Abbott products are for use under medical supervision.
- Abbott products are intended for oral and tube feeding; not for parenteral (IV) use.
- Never use a microwave oven to warm formula.
 Serious burns can result.
- Powdered infant formulas are not sterile and should not be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor.
- Store 2-fl-oz RTF infant formula bottles in a dark environment; prolonged exposure to light should be avoided.

Mixing Instructions: Powder

- Abbott data on calorically dense feedings is limited. Hypocaloric and hypercaloric formulas should be used under the direction of a health care professional
- 27 Cal/fl oz or more calorically dense formula may not supply enough water for some infants. Hydration status should be monitored and water supplied from other sources if necessary
- For improved tolerance, it is best to increase caloric density slowly, by 2- to 4-Cal/fl oz increments

The following tables show the quantity of water to mix with the number of unpacked, level scoop(s) of powder to arrive at the approximate caloric densities shown. Use only the scoop provided in the container. To maintain freshness, pour prepared formula into individual feeding bottles, cap, store in refrigerator, and use within 24 hours.

Similac® 360 Total Care®, Similac® 360 Total Care® Sensitive, Similac® Advance®, Similac® Advance® Lamehadrin, Similac® Alimentum®, Similac® Organic, Similac® Organic with A2 Milk, Similac® PM 60/40, Similac Pro-Advance®, Similac Pro-Sensitive®, Similac Pro-Total Comfort®, Similac Sensitive®, Similac® Soy Isomil®, Similac Total Comfort®

POWDERED PRODUCTS with 8.6 - 8.8 grams powder/scoop				
Caloric Density (Cal/fl oz)	Water, fl oz	Unpacked, Level Scoop	Approximate Yield (fl oz)	
20 (standard)	2	1	2	
22	31/2	2	4	
24	5	3	51/2	
26	11/2	1	2	
27	41/4	3	5	
28	51/2	4	61/2	
30	5	4	6	

Similac® NeoSure® (9.6 g scoop)

POWDER with 9.6 grams powder/scoop					
Caloric Density (Cal/fl oz)	Water (fl oz)	Unpacked, Level Scoop	Approximate Yield (fl oz)		
20	41/2	2	5		
22 (standard)	2	1	2		
24	51/2	3	6		
26	5	3	6		
27	8	5	9		
28	3	2	31/2		
30	7	5	8		

Household Measurements (per cup)

Measurements are based upon 1 unpacked, level, dry measuring cup. For most accurate results, powder should be weighed on a scale that reads in grams.			
Product	Approximate grams of powder		
Similac Advance	100 g		
Similac Advance Lamehadrin	100 g		
Similac Alimentum	100 g		
Similac Alimentum with 2'-FL HMO	100 g		
Similac Organic	100 g		
Similac Organic with A2 Milk	100 g		
Similac NeoSure	100 g		
Similac PM 60/40	100 g		
Similac Pro-Advance	100 g		
Similac Pro-Sensitive	100 g		
Similac Pro-Total Comfort	100 g		
Similac 360 Total Care	100 g		
Similac 360 Total Care Sensitive	100 g		
Similac Sensitive	100 g		
Similac Soy Isomil	100 g		
Similac Total Comfort	100 g		
EleCare products	Measurement by cup not recommended. Weighing powder or use of scoop is suggested.		

To Reconstitute Large Recipe, All Similac Powders[‡] (except EleCare[®] for Infants and EleCare[®] Jr)

The following table shows the quantity of water to mix with 1 unpacked, level, dry measuring cup (not the enclosed scoop) of powder (~100 g) to arrive at the approximate caloric densities shown.

POWDER Mixing Chart			
Caloric Density (Cal/fl oz)	Water (fl oz)	Unpacked, Level, Dry Measuring Cup (approx. 100 g)	Approximate Yield (fl oz)
20	23	1	26
21	22	1	25
22	21	1	24
23	20	1	23
24	19	1	22
25	18	1	21
26	17	1	20
27	16	1	19
28	151/2	1	18
29	15	1	171/2
30	141/2	1	17

For most accurate results, formula powder should be weighed on a scale that reads in grams.

EleCare® (ages 0-12 months)

EleCare (ages 0-12 mo	EleCare (ages 0-12 months), 9.4 g/scoop				
Caloric Density	Water (fl oz)	Scoops* (g)	Approximate Yield (fl oz) [†]		
20 (standard)	2	1 scoop (9.4 g)	2		
22	31/2	2 scoops (18.8 g)	4		
24	8	5 scoops (47 g)	9		
26	1½	1 scoop (9.4 g)	2		
27	7	5 scoops (47 g)	8		
30	5	4 scoops (37.6 g)	6		

EleCare Jr (ages 1 year and up)

EleCare Jr Unflavored & Vanilla, 9.5 g/scoop				
Caloric Density	Water (fl oz)	Scoops* (g)	Approximate Yield (fl oz) [†]	
30	5	4 scoops (38 g)	6	
35	5	5 scoops (47.5 g)	6	
40	31/2	4 scoops (38 g)	4	
45	41/2	6 scoops (57 g)	6	

EleCare Jr Banana, 9 g/scoop				
Caloric Density	Water (fl oz)	Scoops* (g)	Approximate Yield (fl oz) [†]	
30	6	5 scoops (45 g)	7	
35	5	5 scoops (45 g)	6	
40	5	6 scoops (54 g)	6	
45	31/2	5 scoops (45 g)	41/2	

EleCare Jr Chocolate, 8.9 g/scoop				
Caloric Density	Water (fl oz)	Scoops* (g)	Approximate Yield (fl oz) [†]	
30	6	5 scoops (44.5 g)	7	
35	5	5 scoops (44.5 g)	6	
40	5	6 scoops (53.4 g)	6	
45	31/2	5 scoops (44.5 g)	41/2	

^{*}All scoops are unpacked, level.

Household Measurements (per cup)

Measurements are based upon 1 unpacked, level, dry measuring cup. For most accurate results, powder should be weighed on a scale that reads in grams.		
Product Approximate grams of powder		
EleCare	103	
EleCare Jr Unflavored & Vanilla	104	
EleCare Jr Banana	105	
EleCare Jr Chocolate	106	

EleCare (ages 0-12 months)

The following table shows the quantity of water to mix with 1 unpacked, level, dry measuring cup (not the enclosed scoop) of powder to arrive at the approximate caloric densities shown.

Caloric Density	Water (fl oz)	Unpacked, level, dry measuring cup (≈103 g)*	Approximate Yield (fl oz) [†]
20	22	1	25
21	21	1	24
22	20	1	23
23	19	1	22
24	18	1	21
25	17	1	20
26	16	1	19
27	15 ½	1	18
28	15	1	18
29	14 ½	1	17
30	14	1	17

^{*} Household measures are based upon 1 unpacked, level, dry measuring cup. The value provided is approximate as household measure results can vary significantly based on the measuring device and individual methods. Abbott is not responsible for the accuracy of individual users' household measures. For most accurate results, powder should be weighed on a scale that reads in grams.

[†] Yields are rounded to nearest whole number after calculations.

[†] Yields are rounded to the nearest whole number.

EleCare Jr (ages 1 year and up)

The following table shows the quantity of water to mix with 1 unpacked, level, dry measuring cup (not the enclosed scoop) of powder to arrive at the approximate caloric densities shown.

EleCare Jr Unflavored & Vanilla				
Caloric Density	Water (fl oz)	Unpacked, level, dry measuring cup (≈104 g)*	Approximate Yield (fl oz) [†]	
30	13½	1	16	
35	111/4	1	14	
40	91/2	1	12	
45	81/4	1	11	

EleCare Jr Banana			
Caloric Density	Water (fl oz)	Unpacked, level, dry measuring cup (≈105 g)*	Approximate Yield (fl oz) [†]
30	14	1	17
35	111/2	1	14
40	9¾	1	12
45	81/4	1	11

EleCare Jr Chocolate			
Caloric Density	Water (fl oz)	Unpacked, level, dry measuring cup (≈106 g)*	Approximate Yield (fl oz) [†]
30	14	1	17
35	111/2	1	14
40	9¾	1	12
45	81/2	1	11

^{*} Household measures are based upon 1 unpacked, level, dry measuring cup. The value provided is approximate as household measure results can vary significantly based on the measuring device and individual methods. Abbott is not responsible for the accuracy of individual users' household measures. For most accurate results, powder should be weighed on a scale that reads in grams.

Mixing Instructions: Concentrated Liquid

- Abbott data on calorically dense feedings is limited. Hypocaloric and hypercaloric formulas should be used under the direction of a health care professional
- 27 Cal/fl oz or more calorically dense formula may not supply enough water for some infants. Hydration status should be monitored and water supplied from other sources if necessary
- For improved tolerance, it is best to increase caloric density slowly, by 2- to 4-Cal/fl oz increments

[†] Yields are rounded to the nearest whole number

Similac® Advance®, Similac Sensitive®, Similac® Soy Isomil®

The following table shows the quantity of water to mix with the quantity of concentrated liquid to arrive at the approximate caloric densities shown.

To maintain freshness, pour prepared formula into individual feeding bottles, cap, store in refrigerator, and use within 48 hours.

LIQUID Mixing Chart (small volumes)			
Caloric Density (Cal/fl oz)	Water (fl oz)	Concentrated Liquid (fl oz)	Approximate Yield (fl oz)
20 (standard)	1	1	2
22	21/2	3	51/2
24	2	3	5
26	3	51/2	81/2
27	1	2	3
28	2	41/2	61/2
30	1	3	4

The following table shows the quantity of water to mix with one 13-fl-oz (384-mL) can of Concentrated Liquid to arrive at the approximate caloric densities shown. To maintain freshness, pour prepared formula into individual feeding bottles, cap, store in refrigerator, and use within 48 hours.

LIQUID Mixing Chart (13-fl-oz can)				
Caloric Density (Cal/fl oz)	Water (fl oz)	Concentrated Liquid (fl oz)	Approximate Yield (fl oz)	
20 (standard)	13	13	26	
22	11	13	24	
24	9	13	22	
26	7	13	20	
27	6	13	19	
28	51/2	13	181/2	
30	41/2	13	171/2	

Mixing Instructions: Similac® Human Milk Fortifiers

These products are for premature and low-birth-weight Infants. Proper hygiene, handling, and storage are important when preparing fortified human milk. Follow directions as specified. Improper dilution may be harmful.

Powder/Concentrated Liquids

The Powder or Concentrated Liquid Fortifiers are to be mixed with measured amounts of human milk to provide an additional 2 or 4 Cal/fl oz, as shown.

POWDER		
Additional Cal		Similac
Desired	Human Milk	Human Milk Fortifier
2 Cal/fl oz	50 mL	1 packet (0.9 g)
4 Cal/fl oz	25 mL	1 packet (0.9 g)

CONCENTRATED LIQUI	D		
Additional Cal		Similac	Approximate
Desired	Human Milk	Human Milk Fortifier	Yield
2 Cal/fl oz	50 mL	1 packet (5 mL)	55 mL
4 Cal/fl oz	25 mL	1 packet (5 mL)	30 mL

For high-calorie Human Milk Fortification options, see pages 76-79.

Immediate Use

- Pour desired quantity of human milk into clean feeding container.
- Product separation is normal when storing Concentrated Liquid. Massage packet contents until uniform.
- To open packet, tear top off completely along perforation.
- Add contents to human milk and shake until mixed well.
 If bottle feeding, use within 1 hour or discard.

When Stored

- Divide unused fortified human milk into individual bottles or syringes. Refrigerate and use within 24 hours.
- After refrigeration, warm to room temperature.
- Settling may occur; shake for 10 seconds or until mixed well. Feed promptly.

Note: Similac Human Milk Fortifier Concentrated Liquids are light-sensitive. Keep protective carton lid closed when not in use. Keep unopened packets at room temperature in the protective carton.

Caution: Similac Human Milk Fortifiers are nutritionally incomplete by themselves and cannot be ingested alone.

Preparation and Use: PediaSure® Tube Feeding

PediaSure® Enteral Formula 1.0 Cal with Fiber, PediaSure® 1.5 Cal with Fiber, PediaSure® Peptide 1.0 Cal, PediaSure® Peptide 1.5 Cal

All liquid medical foods, regardless of type of administration system, require careful handling because they can support microbial growth. Follow these instructions for clean technique and proper setup to reduce the potential for microbial contamination.

NOTE: Failure to follow the **Instructions for Use** increases the potential for microbial contamination and reduces hangtime.

- Follow physician's instructions.
- Administer product at room temperature.
- THOROUGHLY wash hands with soap and water before handling container or feeding set.

- Turn container upside down and SHAKE VIGOROUSLY, using a twisting motion for at least 10 seconds.
- DO NOT touch any part of the container or feeding set that comes into contact with the formula.
- Visually inspect foil for signs of leakage. DO NOT use if leakage is apparent.
- When initiating feeding, adjust flow rate and volume according to patient's condition and tolerance.
- Additional fluid requirements should be met by giving water between or after feedings or when flushing the tube.

For Use with Enteral Feeding Pumps

- Remove dust cover from RTH Safety Screw Cap.
- Remove dust cover from feeding set connector.
- Insert feeding set connector into port of RTH Safety Screw Cap and completely pierce foil.
- Turn connector collar clockwise until it is securely fastened.
- Close clamp on feeding set before inverting container.
- Invert container and suspend, using hanging ring on bottom of container.

Precautions

- Follow directions for use provided by manufacturer of feeding sets.
- Unless a shorter hangtime is specified by the set manufacturer, hang product for up to 48 hours after initial connection when clean technique and only one new set are used. Otherwise hang for no more than 24 hours.
- Store unopened at room temperature; avoid extreme temperatures.





Similac[®] 360 Total Care[®]

Infant formula with Iron



Description/Indications

A **20 Cal/fl oz**, nutritionally complete, non-GMO, milk-based, iron fortified infant formula designed to support the immune system, brain development and digestive health. Key ingredients found in breast milk including 5 HMO prebiotics. For use as a supplement or alternative to breastfeeding.

Features

- Designed to support the immune system, brain development, and digestive health.
- 5 HMO Prebiotics. HMOs are immune nourishing prebiotics structurally identical to those in breast milk.¹
- DHA, lutein, vitamin E.
- Similac 360 Total Care is our closest formula to breast milk.
- Similac 360 Total Care may soften stools to be more like those of the breastfed infant.
- No artificial growth hormones.*
- Non-GMO.§
- No palm olein oil.
- Kosher/Halal

Precaution

- Not for infants or children with galactosemia
- 1. Hill DR, Chow JM, & Buck RH. Nutrients. 2021;13(10):3364
- * No significant difference has been shown between milk derived from rbST-treated and non-rbST-treated cows.
- § Ingredients not genetically engineered.

Availability: Hospital/Institutional

Size	Container	List No.
Custom Feedi	ng System	
Ready To Feed	d: (20 Cal/fl oz)	
2 fl oz (59 mL)	plastic bottle; 48/case	68136

Availability: Retail

Size	Container	List No.
Ready To Fe	ed: (20 Cal/fl oz)	
2 fl oz (59 m	L) plastic bottle; 12/case	68133
2 fl oz (59 m	L) plastic bottle; 24/case	68130
8 fl oz (237 i	mL) plastic bottle; 24/case	68138
32 fl oz (946	mL) plastic bottle;6/case	68145
`		

Powder: (with measuring scoop)

20.6 oz (584 g); yields 151 ti oz container; 4/case	68063
30.3 oz (859 g); yields 222 fl oz container; 4/case	68065
30.8 oz (874 g); yields 226 fl oz can; 4/case	68116
36 oz (1.02 kg); yields 264 fl oz container;6/case	68111
40 oz (1.13 kg); yields 293 fl oz container; 6/case	68067

-- (FO4 -)...: alala 4F4 fl -- -- -- -- 4/----

Powder: (Stick Pack)

0.60 oz (17.2 g); yields 16 4-fl-oz bottles; 16/case

Ingredients

Ready To Feed: Water, Nonfat Milk, Lactose, High Oleic Safflower Oil, Soy Oil, Coconut Oil, Whey Protein Concentrate. Less than 0.5% of: Human Milk Oligosaccharides (2'-Fucosyllactose, Lacto-N-Tetraose, 3-Fucosyllactose, 6'-Sialyllactose, 3'-Sialyllactose), C. Cohnii Oil', M. Alpina Oil', Beta-Carotene, Lutein, Ascorbic Acid, Soy Lecithin, Monoglycerides, Potassium Citrate, Calcium Carbonate, Potassium Chloride, Carrageenan, Ferrous Sulfate, Magnesium Chloride, Choline Chloride, Choline Bitartrate, Taurine, Inositol, Calcium Phosphate, Calcium Phosphate, Alpha-Tocopheryl Acetate, Niacinamide, Calcium Pantothenate, L-Carnitine, Vitamin A Palmitate, Copper Sulfate, Potassium Hydrochloride, Ribotatin, Pyridoxine Hydrochloride, Fiolic Acid, Manganese Sulfate, Phylloquinone, Biotin, Sodium Selenate, Vitamin D₃, Vitamin B₁₂. Salt, Potassium Iodide, Potassium Hydroxide, and Nucleotides (Adenosine 5'-Monophosphate, Cytidine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate). Contains milk and soy ingredients.

	100 Cal	1000 mL		100 Cal	1000 mL
Energy, Cal	100	676	Biotin, mcg	4.4	29.7
Volume, mL	148	1000	Vitamin C, mg	9	60.8
Protein, g	2.07	13.99	Choline, mg	24	162.2
% Total Cal	8	8	Inositol, mg	24	162.2
Source	Nonfat milk and	whey protein concentrate	Minerals		
Fat, g	5.4	36.49	Calcium, mg	78	527
% Total Cal	49	49	Calcium, mEq	3.9	26.4
Source High	oleic safflower, soy, and coconut oils	(0.15% DHA, 0.4% ARA)	Phosphorus, mg	42	284
Oil Ratio	40:30:29		Magnesium, mg	6	40.5
Linoleic Acid, mg	1000	6757	Iron, mg	1.8	12.16
Carbohydrate, g	11	74.32	Zinc, mg	0.75	5.07
% Total Cal	43	43	Manganese, mcg	5	33.8
Source	Lactose		Copper, mcg	90	608
Prebiotics	Human Milk Oligosaccharides		lodine, mcg	15	101
Vitamins			Selenium, mcg	2	14
Vitamin A, IU	300	2027	Sodium, mg	24	162
Vitamin D, IU	60	405	Sodium, mEq	1	6.8
Vitamin E, IU	1.5	10.1	Potassium, mg	105	709
Vitamin K, mcg	8	54.1	Potassium, mEq	2.7	18.2
Thiamin (Vit B1), mo	g 100	676	Chloride, mg	65	439
Riboflavin (Vit B2), r	ncg 150	1014	Chloride, mEq	1.8	12.2
Vitamin B6, mcg	60	405	Other Characteristics		
Vitamin B12, mcg	0.25	1.69	PRSL, mOsm	18.7	126.4
Niacin, mcg	1050	7095	Water, g	133	899
Folic Acid, mcg	15	101.4	Osmolality, mOsm/kg H2O	310	310
Pantothenic Acid, m	ncg 450	3041			

[†]Source of DHA

[‡]Source of ARA

Human milk oligosaccharides not from human milk.

Similac Pro-Advance®

Infant Formula with Iron



Description/Indications

A 20 Cal/fl oz. nutritionally complete, non-GMO[†], milk-based. iron fortified standard infant formula designed to support both immune and cognitive development with key ingredients found in breast milk. For use as a supplement or alternative to breastfeeding.

Features

- Similac Pro-Advance with 2'-FL Human Milk Oligosaccharide* helps strengthen the immune system to be more like the breastfed infant's than ever before 1-3
- Nucleotides to help support the developing immune system^{2,3}
- OptiGRO® is our exclusive blend of DHA. Lutein, and Vitamin E; these important ingredients are found in breast milk
 - DHA for brain and eye development
 - Lutein to support eve health
- Vitamin E. an important nutrient found in breast milk to support developing cells
- · Our exclusive formula has:
 - Calcium for strong bones no palm olein oil
 - Prebiotics to help promote digestive health
 - Carotenoids like those naturally found in breast milk
- Clinically shown to support normal growth and tolerance in newborns^{4,5}

- Clinically shown to support greater calcium absorption[‡] for strong bones^{6,7}
- Non-GMO
- · Gluten-free
- · Kosher, Halal

Precaution

Not for infants or children with galactosemia

- Goehring KC. J Nutr 2016;146(12):2559-2566.
 Schaller JP, et al. Pediatr Res 2004;56(6):883-890.
- 3. Buck RH, et al. Pediatr Res 2004;56(6):891-900. 4.Marriage B, et al. JPGN 2015;61:649-653.
- Kaizer J. et al. FASEB J 2016;30:S1:671.4.
 Nelson SE, et al. J Am Coll Nutr 1998;17(4):327 332.
- Koo W. et al. Pediatrics 2003:111:1017
- * Not from human milk.
- † Ingredients not genetically engineered.
- [‡] Similac® with Iron vs Enfamil® with Iron.

Enfamil is not a trademark of Abbott Laboratories

Availability: Hosp	ital/Institutional	
Size	Container	List No.
Custom Feeding Sy	stem	
Ready To Feed:	plastic bottle; 48/case	67222
2 II 02 (39 IIIL)	. plastic bottle, 46/case	01323
Availability: Retai	il	
Size	Container	List No.
Ready To Feed:		
2 fl oz (59 mL)	. plastic bottle; 12/case	67562
2 fl oz (59 mL)	plastic bottle; 24/case	67621
8 fl oz (237 mL)	plastic bottle; 24/case	67512
32 fl oz (946 mL)	plastic bottle; 6/case	64248
Powder: (with measu	uring scoop)	
20.6 oz (584 a): viel	ds 151 fl oz container; 4/case	68088
30.8 oz (874 a): viel	ds 226 fl oz container; 4/case.	66439
	249 fl oz container; 6/case	
Powder: (Stick Pack))	
	elds 16 4-fl-oz bottles: 16/cas	e66781

Ingredients

Ready to Feed: Water, Nonfat Milk, Lactose, High Oleic Safflower Oil, Soy Oil, Coconut Oil, Whey Protein Concentrate; Less than 0.5% of: C. Cohnii Oil, M. Alpina Oil, 2'-Fucosyllactose, Short-Chain Fructooligosaccharides, Beta-Carotene, Lutein, Ascorbic Acid, Soy Lecithin, Monoglycerides, Potassium Citrate, Calcium Carbonate, Potassium Chloride, Choline Bitartrate, Taurine, Inositol, Calcium Phosphate, Zinc Sulfate, Potassium Phosphate, d-Alpha-Tocopheryl Acetate, Niacinamide, Calcium Pantothenate, L-Carnitine, Vitamin A Palmitate, Copper Sulfate, Thiamine Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Manganese Sulfate, Phylloquinone, Biotin, Sodium Selenate, Vitamin D₃, Vitamin B₁₂, Salt, Potassium Iodide, Potassium Hydroxide and Nucleotides (Adenosine 5'-Monophosphate, Cytidine 5'-Monophosphate). Contains milk and soy ingredients.

NUTRITION	INFORMATION					
	100 Cal	1000 mL		100 Cal	1000 mL	
Energy, Cal	100	676	Biotin, mcg	4.4	29.7	
Volume, mL	148	1000	Vitamin C, mg	9	60.8	
Protein, g	2.07	13.99	Choline, mg	24	162.2	
% Total Cal	8	8	Inositol, mg	24	162.2	
Source	Nonfat milk, whey protein concen	trate	Minerals			
Fat, g	5.4	36.49	Calcium, mg	78	527	
% Total Cal	49	49	Calcium, mEq	3.9	26.4	
Source Hig	h oleic safflower oil, soy, and coconut oils (0.1	5% DHA, 0.40% ARA)	Phosphorus, mg	42	284	
Oil Ratio	40:30:29	40:30:29	Magnesium, mg	6	40.5	
Linoleic Acid, mg	1000	6757	Iron, mg	1.8	12.16	
Carbohydrate, g	11	74.32	Zinc, mg	0.75	5.07	
% Total Cal	43	43	Manganese, mcg	5	33.8	
Source	Lactose		Copper, mcg	90	608	
Prebiotic	2'-FL human milk oligosaccharides, fructoo	ligosaccharides	lodine, mcg	15	101	
Vitamins			Selenium, mcg	2	14	
Vitamin A, IU	300	2027	Sodium, mg	24	162	
Vitamin D, IU	60	405	Sodium, mEq	1	6.8	
Vitamin E, IU	1.5	10.1	Potassium, mg	105	709	
Vitamin K, mcg	8	54.1	Potassium, mEq	2.7	18.2	
Thiamin (Vit B ₁), mo	g 100	676	Chloride, mg	65	439	
Riboflavin (Vit B ₂), m	ncg 150	1014	Chloride, mEq	1.8	12.2	
Vitamin B ₆ , mcg	60	405	Other Characteristics			
Vitamin B ₁₂ , mcg	0.25	1.69	PRSL, mOsm	18.7	126.7	
Niacin, mcg	1050	7095	Water, g	133	899	
Folic Acid, mcg	15	101.4	Osmolality, mOsm/kg H ₂ O	310	310	
Pantothenic Acid, m	ncg 450	3041				

C. Cohnii Oil is a source of DHA.

M. Alpina Oil is a source of ARA.



Description/Indications

A **20 Cal/fl oz**, nutritionally complete, milk-based, iron-fortified infant formula for use as a supplement or alternative to breastfeeding.

Features

- OptiGRO® is our exclusive blend of DHA, Lutein, and Vitamin E; these important ingredients are found in breast milk
 - DHA for brain and eye development
 - Lutein to support eye health
 - Vitamin E, an important nutrient found in breast milk to support developing cells
- · Our exclusive formula has:
 - Calcium for strong bones-no palm olein oil
- Nucleotides to help support the immune system
- Prebiotics to help promote digestive health
- Carotenoids like those naturally found in breast milk
- Clinically proven tolerance in newborns1
- Clinically shown to support greater calcium absorption* for strong bones^{2,3}
- Gluten-free

· Kosher, Halal

Precaution

- Not for infants or children with galactosemia
- 1. Lloyd B, et al. Pediatrics 1999;103:1-6.
- 2. Nelson SE, et al. J Am Coll Nutr 1998;17(4):327-332.
- 3. Koo W, et al. Pediatrics 2003;111:1017.
- * Similac® with Iron vs Enfamil® with Iron. Enfamil is not a trademark of Abbott Laboratories.

Availability: Retail

Size	Container	List No.
Ready to Feed: 32 fl oz (946 mL)	. bottle/6 ct	53363
Concentrated Liquid 13 fl oz (384 mL)	d: . can; 12/case	56973
20.6 oz (584 g); yiel	uring scoop) ds 90 fl oz container; 6/caseds 151 fl oz container; 6/caseds 226 fl oz container; 6/caseds	68084

Ingredients

Ready to Feed: Water, Nonfat Milk, Lactose, High Oleic Safflower Oil, Soy Oil, Coconut Oil, Galactooligosaccharides[†], Whey Protein Concentrate; Less than 0.5% of: C. Cohnii Oil, M. Alpina Oil, Beta-Carotene, Lutein, Ascorbic Acid, Calcium Carbonate, Potassium Citrate, Soy Lecithin, Monoglycerides, Potassium Chloride, Carrageenan, Magnesium Chloride, Ferrous Sulfate, Choline Bitartrate, Choline Chloride, Taurine, Calcium Phosphate, Potassium Phosphate, Inositol, Zinc Sulfate, Niacinamide, d-Alpha-Tocopheryl Acetate, Calcium Pantothenate, L-Carnitine, Riboflavin, Vitamin A Palmitate, Copper Sulfate, Thiamine Hydrochloride, Pyridoxine Hydrochloride, Folic Acid, Manganese Sulfate, Phylloquinone, Biotin, Sodium Selenate, Vitamin D₃, Vitamin B₁₂, Salt, Potassium Iodide, Potassium Hydroxide, and Nucleotides (Adenosine 5'-Monophosphate, Cytidine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate). Contains milk and soy ingredients.

	100 Cal	1000 mL		100 Cal	1000 mL
Energy, Cal	100	676	Biotin, mcg	4.4	29.7
Volume, mL	148	1000	Vitamin C, mg	9	60.8
Protein, g	2.07	13.99	Choline, mg	24	162.2
% Total Cal	8	8	Inositol, mg	24	162.2
Source	Nonfat milk and whey protein concentrate		Minerals		
Fat, g	5.4	36.49	Calcium, mg	78	527
% Total Cal	49	49	Calcium, mEq	3.9	26.4
Source	High oleic safflower, soy, and coconut oils (0.15)	% DHA, 0.4% ARA)	Phosphorus, mg	42	284
Oil Ratio	40:30:29		Magnesium, mg	6	40.5
Linoleic Acid, mg	1000	6757	Iron, mg	1.8	12.16
Carbohydrate, g	11.2	75.68	Zinc, mg	0.75	5.07
% Total Cal	43	43	Manganese, mcg	5	33.8
Source	Lactose		Copper, mcg	90	608
Prebiotic	Galactooligosaccharides		lodine, mcg	15	101
Vitamins			Selenium, mcg	2	14
Vitamin A, IU	300	2027	Sodium, mg	24	162
Vitamin D, IU	60	405	Sodium, mEq	1	7
Vitamin E, IU	1.5	10.1	Potassium, mg	105	709
Vitamin K, mcg	8	54.1	Potassium, mEq	2.7	18.2
Thiamin (Vit B ₁), mcg	100	676	Chloride, mg	65	439
Riboflavin (Vit B ₂), mcg	150	1014	Chloride, mEq	1.8	12.4
Vitamin B ₆ , mcg	60	405	Other Characteristics		
Vitamin B ₁₂ , mcg	0.25	1.69	PRSL, mOsm	18.7	126.7
Niacin, mcg	1050	7095	Water, g	134	905
Folic Acid, mcg	15	101.4	Osmolality, mOsm/kg H ₂ O	310	310
Pantothenic Acid, mcg	450	3041			

C. Cohnii Oil is a source of DHA.

M. Alpina Oil is a source of ARA.

 $^{^\}dagger\mbox{\sc Prebiotics}$ sourced from milk.

Pure Bliss[®] by Similac[®]

Infant Formula with Iron



Description/Indications

A **19 Cal/fl oz**, nutritionally complete, non-GMO,* milk-based, iron-fortified infant formula that starts with fresh milk from grass-fed cows. For use as a supplement or alternative to breastfeeding.

Features

- European crafted; modeled after breast milk
- Has DHA and ARA, key nutrients to support brain & eye development
- No artificial growth hormones[†]
- No antibiotics
- Prebiotics for digestive health (GOS), lutein for eye health, and nucleotides for immune support
- Calcium for strong bones—no palm olein oil
- · Gluten-free
- · Kosher, Halal

Precaution

· Not for infants or children with galactosemia

*Ingredients not genetically engineered.

[†]No significant difference has been shown between milk derived from rbST-treated and non-rbST-treated cows.

Availability: Retail

Ingredients

Powder: Nonfat Milk, Lactose, High Oleic Sunflower Oil, Whey Protein Concentrate, Soy Oil, Coconut Oil, Galactooligosaccharides'. Less than 2% of: C. Cohnii Oil, M. Alpina Oil, Beta-Carotene, Lutein, Potassium Citrate, Calcium Carbonate, Ascorbic Acid, Soy Lecithin, Potassium Chloride, Magnesium Chloride, Ferrous Sulfate, Choline Bitartrate, Choline Chloride, Ascorbyl Palmitate, Salt, Taurine, m-Inositol, Zinc Sulfate, Mixed Tocopherols, d-Alpha-Tocopheryl Acetate, Niacinamide, Calcium Pantothenate, L-Carnitine, Vitamin A Palmitate, Copper Sulfate, Thiamine Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Manganese Sulfate, Phylloquinone, Biotin, Sodium Selenate, Vitamin D₃, Vitamin B₁₂. Calcium Phosphate, Potassium Hydroxide, Potassium Iodide, and Nucleotides (Adenosine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate). Contains milk and soy ingredients.

NUTRITION IN	FORMATION				
	100 Cal (5.3 fl oz, as prepared)	1000 mL (1.1 qt, as prepared)		100 Cal (5.3 fl oz, as prepared)	1000 mL (1.1 qt, as prepared)
Energy, Cal	100	645	Biotin, mcg	4.6	29.7
Volume, mL	156	1000	Vitamin C, mg	9	58
Protein, g	2.07	13.34	Choline, mg	24	154.7
% Total Calories	8	8	Inositol, mg	4.9	31.6
Source	Nonfat milk and whey protein concentrat	e	Minerals		
Fat, g	5.6	36.1	Calcium, mg	82	529
% Total Calories	50	50	Calcium, mEq	4.1	26.4
Source	High oleic sunflower oil, soy, and coconut of	ils (0.2% DHA, 0.5% ARA)	Phosphorus, mg	44	284
Oil Ratio	40:31:27	40:31:27	Magnesium, mg	6	38.7
Linoleic Acid, mg	938	6047	Iron, mg	1.9	12.3
Carbohydrate, g	10.7	68.98	Zinc, mg	0.79	5.09
% Total Calories	42	42	Manganese, mcg	5	32.2
Source	Lactose		Copper, mcg	95	612
Prebiotic	Galactooligosaccharides		lodine, mcg	8	52
Vitamins			Selenium, mcg	2	13
Vitamin A, IU	300	1934	Sodium, mg	25	161
Vitamin D, IU	60	387	Sodium, mEq	1.1	7
Vitamin E, IU	1.5	9.7	Potassium, mg	110	709
Vitamin K, mcg	8	51.6	Potassium, mEq	2.8	18.2
Thiamin (Vit B ₁), mcg	100	645	Chloride, mg	68	438
Riboflavin (Vit B ₂), mcg	160	1031	Chloride, mEq	1.9	12.4
Vitamin B ₆ , mcg	63	406	Other Characteristics		
Vitamin B ₁₂ , mcg	0.26	1.68	PRSL, mOsm	19.1	122.9
Niacin, mcg	1100	7091	Water, g	141	904
Folic Acid, mcg	16	103.1	Osmolality, mOsm/kg H ₂ O	305	305
Pantothenic Acid, mcg	470	3030			

C. Cohnii Oil is a source of DHA.

M. Alpina Oil is a source of ARA.

[†] Prebiotics sourced from milk.

Similac[®] Organic

Infant Formula with Iron



Description/Indications

A 20 Cal/fl oz, nutritionally complete, organic, milk-based, iron-fortified infant formula for use as a supplement or alternative to breastfeeding. Certified USDA Organic. Made with premium ingredients, including lactose like that found in breast milk.

Features

- OptiGRO® is our exclusive blend of DHA, Lutein, and Vitamin E; these important ingredients are found in breast milk
 - DHA for brain and eye development
 - Lutein to support eye health
 - Vitamin E, an important nutrient to support developing cells
- · Our exclusive formula has:
 - Calcium for strong bones-no palm olein oil
 - Nucleotides to help support the immune system
 - Prebiotics to help promote digestive health
 - Carotenoids like those naturally found in breast milk
- Has NO artificial growth hormones,* NO antibiotics, and it's non-GMO[†]

- Gluten-free
- Kosher, Halal

* No significant difference has been shown between milk derived from rbST-treated and non-rbST-treated cows.

Precaution

· Not for infants or children with galactosemia

Availability: Hospital/Institutional

Size Container List No.
Custom Feeding System
Ready To Feed:
2 fl oz (59 mL)....... plastic bottle; 48/case..................56525

Availability: Retail

Size

Ready to Feed: 32 fl oz (946 mL)..... plastic bottle; 6/carton................... 68403

List No.

Container

Powder: (with measuring scoop)

20.6 oz (584 g); yields 152 fl oz container; 4/case............ 68092

[†] Ingredients not genetically engineered.

Ready to Feed: Water, Organic Lactose, Organic Nonfat Milk, Organic High Oleic Sunflower Oil, Organic Soy Oil, Organic Coconut Oil; Less than 0.5% of: C. Cohnii Oil, M. Alpina Oil, Beta Carotene, Lutein, Short-Chain Fructooligosaccharides, Potassium Citrate, Calcium Carbonate, Calcium Citrate, Ascorbic Acid, Organic Soy Lecithin, Ferrous Sulfate, Salt, Choline Chloride, Choline Bitartrate, Taurine, Inositol, Magnesium Chloride, Zinc Sulfate, d-Alpha-Tocopheryl Acetate, Niacinamide, Calcium Pantothenate, L-Camittine, Vitamin A Palmitate, Copper Sulfate, Thiamine Hydrochloride, Fibic Acid, Manganese Sulfate, Phylloquinone, Biotin, Sodium Selenate, Vitamin D₃, Vitamin B₁₂, Potassium Iodide, Potassium Hydroxide, and Nucleotides (Adenosine 5*-Monophosphate, Cytidine 5*-Monophosphate). Contains milk and soy ingredients.

NUTRITION	INFORMATION				
	100 Cal	1000 mL		100 Cal	1000 mL
Energy, Cal	100	676	Vitamin C, mg	9	60.8
Volume, mL	148	1000	Choline, mg	24	162.2
Protein, g	2.07	13.99	Inositol, mg	24	162.2
% Total Cal	8	8	Minerals		
Source	Organic nonfat milk		Calcium, mg	78	527
Fat, g	5.63	38.06	Calcium, mEq	3.9	26.4
% Total Cal	50	50	Phosphorus, mg	42	284
Source	Organic high oleic sunflower, soy, and coconut	oils (0.15% DHA, 0.4% ARA)	Magnesium, mg	6	40.5
Oil Ratio	40:30:29		Iron, mg	1.8	12.16
Linoleic Acid, mg	860	5811	Zinc, mg	0.75	5.07
Carbohydrate, g	10.4	70.27	Manganese, mcg	5	33.8
% Total Cal	42	42	Copper, mcg	90	608
Source	Organic lactose		lodine, mcg	15	101
Prebiotic	Fructooligosaccharides		Selenium, mcg	2	14
Vitamins			Sodium, mg	24	162
Vitamin A, IU	300	2027	Sodium, mEq	1	6.8
Vitamin D, IU	60	405	Potassium, mg	105	709
Vitamin E, IU	1.5	10.1	Potassium, mEq	2.7	18.2
Vitamin K, mcg	8	54.1	Chloride, mg	65	439
Thiamin (Vit B ₁), mcg	100	676	Chloride, mEq	1.8	12.2
Riboflavin (Vit B ₂), me	cg 150	1014	Other Characteristics		
Vitamin B ₆ , mcg	60	405	PRSL, mOsm	18.7	126.4
Vitamin B ₁₂ , mcg	0.25	1.69	Water, g	133	899
Niacin, mcg	1050	7095	Osmolality, mOsm/kg H ₂ O	300	300
Folic Acid, mcg	15	101.4			
Pantothenic Acid, me	cg 450	3041			
Biotin, mcg	4.4	29.7			

C. Cohnii Oil is a source of DHA.

M. Alpina Oil is a source of ARA.

Similac[®] Organic with A2 Milk

Infant Formula with Iron



Description/Indications

A **20 Cal/fl oz**, organic, milk-based, iron-fortified infant formula for use as a supplement or alternative to breastfeeding. Certified USDA Organic. Made with premium ingredients, including organic lactose.

First organic infant formula made with A2 milk.

Features

- OptiGRO® is our exclusive blend of DHA, Lutein, and Vitamin E; these important ingredients are found in breast milk
 - DHA for brain and eye development
 - Lutein to support eye health
 - Vitamin E, an important nutrient to support developing cells
- Complete, gentle nutrition with A2 milk protein
- · Our exclusive formula has:
 - Calcium for strong bones-no palm olein oil
 - Nucleotides to help support the immune system
 - Prebiotics to help promote digestive health
 - Carotenoids like those naturally found in breast milk

- Has NO artificial growth hormones,* NO antibiotics, and it's non-GMO[†]
- · Gluten-free
- Kosher, Halal

Precaution

· Not for infants or children with galactosemia

Availability: Retail

Size Container List No.

Powder: (with measuring scoop)

20.6 oz (584 g); yields 152 fl oz container; 6/case 68099

^{*} No significant difference has been shown between milk derived from rbST-treated and non-rbST-treated cows.

[†] Ingredients not genetically engineered.

Powder: Organic Lactose, Organic Nonfat Milk, Organic High Oleic Sunflower Oil, Organic Soy Oil, Organic Coconut Oil; Less than 0.5% of: C. Cohnii Oil, M. Alpina Oil, Beta-Carotene, Lutein, Short-Chain Fructooligosaccharides, Potassium Citrate, Calcium Carbonate, Ascorbic Acid, Ascorbyl Palmitate, Ferrous Sulfate, Salt, Choline Chloride, Choline Bitartrate, Taurine, Inositol, Magnesium Chloride, Zinc Sulfate, Mixed Tocopherols, d-Alpha-Tocopheryl Acetate, Niacinamide, Calcium Panethe, L-Carnitine, Vitamin A Palmitate, Copper Sulfate, Thiamine Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Manganese Sulfate, Phylloquinone, Biotin, Sodium Selenate, Vitamin B₁₂: Potassium lodide, Potassium Hydroxide, and Nucleotides (Adenosine 5-Monophosphate, Cytidine 5'-Monophosphate). Contains milk ingredients.

	100 Cal	1000 mL		100 Cal	1000 mL
Energy, Cal	100	676	Vitamin C, mg	9	60.8
Volume, mL	148	1000	Choline, mg	24	162.2
Protein, g	2.07	13.99	Inositol, mg	24	162.2
% Total Cal	8	8	Minerals		
Source	Organic nonfat A2 milk		Calcium, mg	78	527
Fat, g	5.63	38.05	Calcium, mEq	3.9	26.4
% Total Cal	50	50	Phosphorus, mg	42	284
	Organic high oleic sunflower, soy, and coconut	oils (0.15% DHA, 0.4% ARA)	Magnesium, mg	6	40.6
Oil Ratio	40:30:29		Iron, mg	1.8	12.17
Linoleic Acid, mg	860	5811	Zinc, mg	0.75	5.07
Carbohydrate, g	10.4	70.29	Manganese, mcg	5	33.8
% Total Cal	42	42	Copper, mcg	90	608
Source	Organic lactose		lodine, mcg	15	101
Prebiotic	Fructooligosaccharides		Selenium, mcg	2	14
Vitamins			Sodium, mg	24	162
Vitamin A, IU	300	2028	Sodium, mEq	1	6.8
Vitamin D, IU	60	406	Potassium, mg	105	710
Vitamin E, IU	1.5	10.1	Potassium, mEq	2.7	18.2
Vitamin K, mcg	8	54.1	Chloride, mg	65	439
Thiamin (Vit B ₁), mcg	100	676	Chloride, mEq	1.8	12.2
Riboflavin (Vit B2), mc		1014	Other Characteristics		
Vitamin B ₆ , mcg	60	406	PRSL, mOsm	18.7	126.4
Vitamin B ₁₂ , mcg	0.25	1.69	Water, g	133	899
Niacin, mcg	1050	7097	Osmolality, mOsm/kg H ₂ O	295	295
Folic Acid, mcg	15	101.4			
Pantothenic Acid, mo		3041			
Biotin, mcg	4.4	29.7			

C. Cohnii Oil is a source of DHA.

M. Alpina Oil is a source of ARA.

Similac[®] Advance[®] Kosher Cholov Yisroel





Description/Indications

A **20 Cal/fl oz**, nutritionally complete infant formula for use as a supplement or alternative to breastfeeding. Cholov Yisroel.

Features

- · Cholov Yisroel Kosher certification
- Acceptable for those who consume kitniyot on Passover
- DHA & ARA for brain and eye development
- Vitamin E, an important nutrient found in breast milk to support development
- · Calcium & vitamin D for strong bones
- · No artificial growth hormones*
- Provides excellent calcium and fat absorption, improved bone mineralization, and softer stools due to no palm olein oil.
- · Gluten-free

Precaution

- · Not for infants with galactosemia
- * No significant difference has been shown between milk derived from rbST-treated and non-rbST-treated cows.

Ready to Feed: Water, Lactose, Milk Protein Concentrate, High Oleic Safflower Oil, Soy Oil, Coconut Oil; Less than 0.5% of: C. Cohnii Oil, M. Alpina Oil, Potassium Chloride, Sodium Citrate, Calcium Carbonate, Potassium Phosphate, Ascorbic Acid, Monoglycerides, Soy Lecithin, Magnesium Phosphate, Carrageenan, Potassium Citrate, Choline Chloride, m-Inositol, Ferrous Sulfate, Taurine, Zinc Sulfate, d-Alpha-Tocopheryl Acetate, L-Carnitine, Niacinamide, Calcium Pantothenate, Riboflavin, Vitamin A Palmitate, Cupric Sulfate, Thiamine Chloride Hydrochloride, Pyridoxine Hydrochloride, Beta-Carotene, Potassium Iodide, Folic Acid, Manganese Sulfate, Phylloquinone, Potassium Hydroxide, Biotin, Sodium Selenate, Vitamin D₃, Cyanocobalamin, Magnesium Chloride, and Calcium Phosphate. Contains milk and soy ingredients.

NUTRITION IN	IFORMATION				
	100 Cal	1000 mL		100 Cal	1000 mL
Energy, Cal	100	676	Choline, mg	16	108.1
Volume, mL	148	1000	Inositol, mg	4.7	31.8
Protein, g	2.15	14.53	Minerals		
% Total Cal	8	8	Calcium, mg	78	527
Source	Milk protein concentrate		Calcium, mEq	3.9	26.4
Fat, g	5.4	36.49	Phosphorus, mg	42	284
% Total Cal	49	49	Magnesium, mg	6	40.5
Source	High oleic safflower, soy, and coconut oils	(0.15% DHA, 0.4% ARA)	Iron, mg	1.8	12.16
Oil Ratio	40:30:29		Zinc, mg	0.75	5.07
Linoleic Acid, mg	1000	6757	Manganese, mcg	5	33.8
Carbohydrate, g	10.7	72.3	Copper, mcg	90	608
% Total Cal	43	43	lodine, mcg	6	41
Source	Lactose		Selenium, mcg	2	14
Vitamins			Sodium, mg	24	162
Vitamin A, IU	300	2027	Sodium, mEq	1	7
Vitamin D, IU	75	507	Potassium, mg	105	709
Vitamin E, IU	1.5	10.1	Potassium, mEq	2.7	18.2
Vitamin K, mcg	8	54.1	Chloride, mg	65	439
Thiamin (Vit B ₁), mcg	100	676	Chloride, mEq	1.8	12.2
Riboflavin (Vit B2), mcg	150	1014	Other Characteristics		
Vitamin B ₆ , mcg	60	405	PRSL, mOsm	19.2	129.8
Vitamin B ₁₂ , mcg	0.25	1.69	Water, g	133	899
Niacin, mcg	1050	7095	Osmolality, mOsm/kg H ₂ O	310	310
Folic Acid, mcg	15	101.4			
Pantothenic Acid, mcg	450	3041			
Biotin, mcg	4.4	29.7			
Vitamin C, mg	9	60.8			

C. Cohnii Oil is a source of DHA.

M. Alpina Oil is a source of ARA.

Similac[®] Advance[®] Lamehadrin

Infant Formula with Iron





Description/Indications

A **20 Cal/fl oz**, nutritionally complete infant formula for use as a supplement or alternative to breastfeeding. Badatz-certified. Cholov Yisroel.

Features

- First and only BADATZ-CERTIFIED infant formula registered for sale in the US
- DHA & ARA for brain and eye development¹⁻³
- Vitamin E, an important nutrient found in breast milk to support development^{4,5}
- Calcium & vitamin D for strong bones
- · Nucleotides for immune support
- No artificial growth hormones*
- · Gluten-free

Precaution

· Not for infants or children with galactosemia

Availability: Retail

Size Container List No.
Powder: (with measuring scoop)

Powder: (with measuring scoop)

24.7 oz (700 g); yields 180 fl oz can; 12/case.....67321

- 1. Auestad N, et al. Pediatrics. 2001;108(2):372-381.
- 2. Auestad N, et al. Pediatrics. 2003;112(3, pt 1):e177-e183.
- 3. Scott DT. et al. Pediatrics. 1998:102(5):E59.
- 4. Gaur S. et al. J Nutr. 2017;147(7):1301-1307.
- 5. Takada T, et al. Mol Nutr Food Res. 2010;54(5):616-622.

^{*} No significant difference has been shown between milk derived from rbST-treated and non-rbST-treated cows.

Powder: Nonfat Milk, Lactose, High Oleic Sunflower Oil, Soy Oil, Coconut Oil, Whey Protein Concentrate; Less than 2% of: C. Cohnii Oil, M. Alpina Oil, Beta-Carotene, Potassium Citrate, Calcium Carbonate, Ascorbic Acid, Soy Lecithin, Potassium Chloride, Magnesium Chloride, Ferrous Sulfate, Choline Bitartrate, Ascorbyl Palmitate, Salt, Taurine, L-Tryptophan, Inositol, Zinc Sulfate, Mixed Tocopherols, d-Alpha-Tocopherols, d-Alpha-Tocopherol Acetate, Niacinamide, Calcium Pantothenate, L-Carnitine, Vitamin A Palmitate, Copper Sulfate, Thiamine Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Manganese Sulfate, Phylloquinone, Biotin, Sodium Selenate, Vitamin D₃, Vitamin B₁₂, Potassium Iodide, Calcium Phosphate, Potassium Hydroxide and Nucleotides (Cytidine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate). Contains milk and soy ingredients.

	100 Cal	1000 mL		100 Cal	1000 mL
Francis Oct	100	675	Diations		
Energy, Cal			Biotin, mcg	2.9	19.6
Volume, mL	148	1000	Vitamin C, mg	15	101.3
Protein, g	2.12	14.31	Choline, mg	12	81
% Total Cal		8	Inositol, mg	4.7	31.7
Source		d whey protein concentrate	Minerals		
Fat, g	5.2	35.11	Calcium, mg	62	419
% Total Cal	47	47	Calcium, mEq	3.1	20.9
Source		coconut oils (0.15% DHA, 0.4% ARA)	Phosphorus, mg	34	230
Oil Ratio	40:30:28		Magnesium, mg	6	40.5
Linoleic Acid, mg	771	5205	Iron, mg	1.4	9.45
Carbohydrate, g	11.2	75.62	Zinc, mg	0.58	3.92
% Total Cal	45	45	Manganese, mcg	16	108
Source	Lactose		Copper, mcg	60	405
Vitamins			lodine, mcg	16	108
Vitamin A, IU	278	1877	Selenium, mcg	2	14
Vitamin D, IU	40	270	Sodium, mg	21	142
Vitamin E, IU	2.3	15.5	Sodium, mEq	0.9	6.1
Vitamin K, mcg	8	54	Potassium, mg	98	662
Thiamin (Vit B,), mcg	93	628	Potassium, mEq	2.5	16.9
Riboflavin (Vit B ₂), mcg	129	871	Chloride, mg	55	371
Vitamin B ₆ , mcg	62	419	Chloride, mEq	1.5	10.1
Vitamin B ₁₀ , mcg	0.23	1.55	Other Characteristics		
Niacin, mcg	835	5637	PRSL, mOsm	18.2	122.9
Folic Acid, mcg	11	74.3	Water, q	135	911
Pantothenic Acid, mca	466	3146	Osmolality, mOsm/kg H ₂ O	298	289

C. Cohnii Oil is a source of DHA.

M. Alpina Oil is a source of ARA.

Similac[®] Soy Isomil[®]

Infant Formula with Iron



Description/Indications

A 20 Cal/fil oz, nutritionally complete, soy-based infant formula for infants with feeding problems, such as fussiness and gas, and for infants whose parents choose a milk-free formula as a first feeding or as a supplement to breastfeeding. Plant-based protein nutritional formula for infants of vegetarian families. A lactose-free feeding for infants with disorders for which lactose should be avoided, including lactase deficiency, lactose intolerance, and galactosemia.

Features

- · Easy-to-digest gentle soy nutrition
- OptiGRO® is our exclusive blend of DHA, Lutein, and Vitamin E; these important ingredients are found in breast milk
 - DHA for brain and eye development
 - Lutein to support eye health
 - Vitamin E, an important nutrient found in breast milk to support developing cells
- · Our exclusive formula has:
- Calcium for strong bones-no palm olein oil
- Nucleotides to help support the immune system*
- Prebiotics to help promote digestive health
- Carotenoids like those naturally found in breast milk
- The first and only soy formula clinically shown to help support the developing immune system, 1.2 similar to infants fed human milk/ milk-based formula[†]
- Clinically shown to reduce fussiness, gas in infants experiencing general intolerance on cow's milk formula³

- Clinically shown to help with dietary management of infants with diarrhea⁴
- Excellent growth during baby's first year⁵
- Soy protein isolate to help manage IgE-mediated cow's milk protein allergy or sensitivity
- A unique blend of carbohydrates using two different absorption pathways to help maximize absorption and minimize malabsorption risks
- Low osmolality (200 mOsm/kg water)
- Gluten-free
- · Kosher, Halal
- 1. Ostrom KM, et al. J Ped Gastroenterol Nutr 2002;34:137-144.
- 2. Cordle CT, et al. J Ped Gastroenterol Nutr 2002;34:145-153.
- 3. Lasekan JB and Baggs GM, Clin Pediatr 2021;60(3):184-192.
- 4. Naidoo BT et al. J Intern Med Res 1981;9: 232-235.
- 5. Lasekan JB, et al. Clin Pediatr 1999;38:563-571.
- * Soy protein isolate is a source of nucleotides.
- [†] Exclusively breastfed for at least 2 months and weaned to milk-based formula without added nucleotides.

Precaution

 Soy formulas are not recommended for premature infants with birth weights less than 1800 g

Availability: Hospital/Institutional	
Size Container	List No.
Custom Feeding System	
Ready To Feed:	
2 fl oz (59 mL) plastic bottle; 48/case	67391
Availability: Retail	
Size Container	List No.
Ready To Feed: (20 Cal/fl oz)	
32 fl oz (946 mL) plastic bottle; 6/carton	67012
Concentrated Liquid: (40 Cal/fl oz)	
13 fl oz (384 mL) can; 12/case	56975
Powder: (with measuring scoop)	
12.4 oz (352 g); yields 90 fl oz can; 6/case	
30.8 oz (873 g); yields 223 fl oz container; 4/case	67699

Ready to Feed: Water (85%), Corn Syrup (6.6%), Soy Protein Isolate (1.9%), High Oleic Safflower Oil (1.4%), Sugar (1.3%), Soy Oil (1.1%), Coconut Oil (1.0%); Less than 0.5% of: C. Cohnii Oil, M. Alpina Oil, Beta-Carotene, Lutein, Short-Chain Fructooligosaccharides, Calcium Citrate, Calcium Phosphate, Potassium Phosphate, Potassium Hydroxide, Ascorbic Acid, Potassium Citrate, Magnesium Chloride, Monoglycerides, Soy Lecithin, Salt, L-Methionine, Carrageenan, Chloride Chloride, Potassium Chloride, Taurine, Ferrous Sulfate, Inositol, Zinc Sulfate, L-Carnitine, Niacinamide, d-Alpha-Tocopheryl Acetate, Calcium Pantothenate, Riboflavin, Vitamin A Palmitate, Copper Sulfate, Thiamine Hydrochloride, Pyridoxine Hydrochloride, Folic Acid, Potassium Iodide, Phylloquinone, Biotin, Sodium Selenate, Vitamin D₃, and Vitamin B₃, Contains soy ingredients.

	400.0	4000		400.0.1	4000 1
	100 Cal	1000 mL		100 Cal	1000 mL
Energy, Cal	100	676	Pantothenic Acid, mcg	750	5068
Volume, mL	148	1000	Biotin, mcg	4.5	30.4
Protein, g	2.45	16.55	Vitamin C, mg	9	60.8
% Total Cal	10	10	Choline, mg	24	162.2
Source	Soy protein isolate, L-methionine		Inositol, mg	24	162.2
Fat, g	5.46	36.89 I	Minerals		
% Total Cal	49	49	Calcium, mg	105	709
Source	High oleic safflower, soy, and coconut oils (0.	15% DHA, 0.4% ARA)	Calcium, mEq	5.25	35.8
Oil Ratio	40:30:29		Phosphorus, mg	75	507
Linoleic Acid, mg	1000	6757	Magnesium, mg	7.5	50.7
Carbohydrate, g	10.4	70.27	Iron, mg	1.8	12.16
% Total Cal	41	41	Zinc, mg	0.75	5.07
Source	Corn syrup, sugar		Manganese, mcg	25	168.9
Ratio	80:20		Copper, mcg	75	507
Prebiotic	Fructooligosaccharides		lodine, mcg	15	101
Vitamins			Selenium, mcg	2	14
Vitamin A, IU	300	2027	Sodium, mg	44	297
Vitamin D, IU	60	405	Sodium, mEq	1.9	12.8
Vitamin E, IU	1.5	10.1	Potassium, mg	108	730
Vitamin K, mcg	11	74.3	Potassium, mEq	2.77	18.9
Thiamin (Vit B ₁), mcg	60	405	Chloride, mg	62	419
Riboflavin (Vit B ₂), mcg	90	608	Chloride, mEq	1.75	12.2
Vitamin B ₆ , mcg	60	405	Other Characteristics		
Vitamin B ₁₂ , mcg	0.45	3.04	PRSL, mOsm	22.8	154.1
Niacin, mcg	1350	9122	Water, g	133	899
Folic Acid, mcg	15	101.4	Osmolality, mOsm/kg H _o O	200	200

C. Cohnii Oil is a source of DHA.

M. Alpina Oil is a source of ARA.

Similac® 360 Total Care® Sensitive

Infant formula with Iron



Description/Indications

A **20 Cal/fl oz**, nutritionally complete, non-GMO, milk-based, iron fortified infant formula designed for use for sensitive tummies as a milk-based alternative to standard milk-based formulas for sensitive tummies. Designed to support the immune system, brain development and digestive health. Has key ingredients found in breast milk including 5 HMO prebiotics. Suitable for infants with lactose sensitivity.

Features

- Designed to support the immune system, brain development, and digestive health.
- 5 HMO Prebiotics. HMOs are immune nourishing prebiotics structurally identical to those in breast milk.¹
- DHA, lutein, vitamin E.
- Similac 360 Total Care Sensitive for fussiness and gas due to lactose sensitivity.
- No artificial growth hormones.*
- Non-GMO.§
- No palm olein oil.
- Kosher/Halal

Precaution

- Not for infants or children with galactosemia
- 1. Hill DR, Chow JM, & Buck RH. Nutrients. 2021;13(10):3364
- * No significant difference has been shown between milk derived from rbST-treated and non-rbST-treated cows.
- § Ingredients not genetically engineered.

Powder: (with measuring scoop)

Availability:	Hospital/Institutional	
Size	Container	List No
Custom Feedin	ng System	
Ready To Feed	: (20 Cal/fl oz)	
2 fl oz (59 mL).	plastic bottle; 48/case	68155
Availability:	Retail	
Size	Container	List No
Ready To Feed	: (20 Cal/fl oz)	
2 fl oz (59 mL).	plastic bottle; 12/case	68149
2 fl oz (59 mL).	plastic bottle; 24/case	68152
8 fl oz (237 mL) plastic bottle; 24/case	68157
32 fl oz (946 m	L) plastic bottle;6/case	68160
,		

20.1 oz (570 g); yields 145 fl oz container; 4/case 68070

29.5 oz (837 g); yields 213 fl oz container; 4/case 68072

Ready To Feed: Water, Maltodextrin, Milk Protein Isolate, High Oleic Safflower Oil, Sugar, Soy Oil, Coconut Oil. Less than 0.5% of: Human Milk Oligosaccharides (2'-fucosyllactose, Lacto-N-Tetraose, 3-Fucosyllactose, 6'-Sialyllactose, 3'-Sialyllactose)*, C. Cohnii Oil*, M. Alpina Oil*, Beta-Carotene, Lutein, Calcium Phosphate, Potassium Citrate, Carrageenan, Potassium Chloride, Ascorbic Acid, Monoglycerides, Soy Lecithin, Magnesium Chloride, Inositol, Magnesium Phosphate, Choline Chloride, Calcium Carbonate, Ferrous Sulfate, Choline Bitartrate, Taurine, Zinc Sulfate, L-Carnitine, d-Alpha-Tocopheryl Acetate, Niacinamide, Calcium Partotherate, Vitamin Palmitate, Copper Sulfate, Thiamine Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Potassium Iodide, Manganese Sulfate, Phylloquinone, Biotin, Sodium Selenate, Vitamin D₃, Vitamin B₁, Sodium Citrate, Potassium Hydroxide, and Nucleotides (Adenosine 5'-Monophosphate, Ottidine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate, Disodium Selenate, Vitamine B, Carrageenan, Potassium Hydroxide, and Nucleotides (Adenosine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate, Disodium Selenate, Vitamine B, Carrageenan, Potassium Hydroxide, and Nucleotides (Adenosine 5'-Monophosphate, Ottidine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate, Ottidine S'-Monophosphate, Disodium Uridine 5'-Monophosphate, Ottidine S'-Monophosphate, Disodium Uridine S'-Monophosphate, Ottidine S'-Monophosphate, Disodium Guanosine S'-Monophosphate, Disodium Uridine S'-Monophosphate, Ottidine S'-Monophosphate, Disodium Guanosine S'-Monophosphate, Disodium Uridine S'-Monophosphate, Disodium

	100 Cal	1000 mL		100 Cal	1000 mL
Energy, Cal	100	676	Pantothenic Acid, mcg	450	3041
Volume, mL	148	1000	Biotin, mcg	4.4	29.7
Protein, g	2.1	14.19	Vitamin C, mg	9	60.8
% Total Cal	8	8	Choline, mg	24	162.2
Source	Milk pro	tein isolate	Inositol, mg	24	162.2
Fat, g	5.4	36.49	Minerals		
% Total Cal	49	49	Calcium, mg	84	568
Source Hig	h oleic safflower, soy, and coconut oils (0.	5% DHA, 0.4% ARA)	Calcium, mEq	4.2	28.4
Oil Ratio	40:30:29		Phosphorus, mg	56	378
Linoleic Acid, mg	1000	6757	Magnesium, mg	6	40.5
Carbohydrate, g	10.9	73.65	Iron, mg	1.8	12.16
% Total Cal	43	43	Zinc, mg	0.75	5.07
Source	Maltodextrin, sugar		Manganese, mcg	5	33.8
Ratio	80:20		Copper, mcg	90	608
Prebiotics	Human Milk Oligosaccharides		lodine, mcg	15	101
Vitamins			Selenium, mcg	2	14
Vitamin A, IU	300	2027	Sodium, mg	30	203
Vitamin D, IU	60	405	Sodium, mEq	1.3	8.8
Vitamin E, IU	1.5	10.1	Potassium, mg	107	723
Vitamin K, mcg	8	54.1	Potassium, mEq	2.7	18.2
Thiamin (Vit B1), m	cg 100	676	Chloride, mg	65	439
Riboflavin (Vit B2),	mcg 150	1014	Chloride, mEq	1.8	12.2
Vitamin B _e , mcg	60	405	Other Characteristics		
Vitamin B ₁₂ , mcg	0.25	1.69	PRSL, mOsm	19.7	133.2
Niacin, mcg	1050	7095	Water, g	133	899
Folic Acid, mcg	15	101.4	Osmolality, mOsm/kg H2O	200	200

[†] Source of DHA

[‡] Source of ARA

Similac Pro-Sensitive®



Description/Indications

A **20 Cal/fl oz**, nutritionally complete, non-GMO,[†] iron-fortified infant formula designed for use as a milk-based alternative to standard milk-based formulas for sensitive tummies. Designed to support both immune and cognitive development with key ingredients found in breast milk. Suitable for infants with lactose sensitivity.[‡]

Features

- Gentle nutrition designed to ease common tummy troubles like fussiness,[§] gas,[§] or mild spit-up
- Similac Pro-Sensitive with 2'-FL Human Milk Oligosaccharide* helps strengthen the immune system to be more like the breastfed infant's than ever before¹-³
- Nucleotides to help support the developing immune system^{2,3}
- A unique blend of two carbohydrates using two different absorption pathways to help maximize absorption and minimize malabsorption risks
- OptiGRO® is our exclusive blend of DHA, Lutein, and Vitamin E; these important ingredients are found in breast milk
 - DHA for brain and eye development
 - Lutein to support eye health
 - Vitamin E, an important nutrient found in breast milk to support developing cells

- · Our exclusive formula has:
 - Calcium for strong bones no palm olein oil
 - Prebiotics to help promote digestive health
 - Carotenoids like those naturally found in breast milk
- Promotes excellent growth, tolerance, and soft stools⁴
- Low osmolality (200 mOsm/kg water)
- Non-GMO
- · Gluten-free
- · Kosher, Halal

Precaution

- · Not for infants or children with galactosemia
- Goehring KC. J Nutr 2016;146(12):2559-2566.
- Schaller JP, et al. Pediatr Res 2004;56(6):883-890.
- Buck RH, et al. Pediatr Res 2004;56(6):891-900.
- Lasekan JB, et al. Clin Pediatr 2011;50:330-337.
- * Not from human milk.
- † Ingredients not genetically engineered.
- [‡] Typical Value 0.1 g lactose/100 kcal compared to Similac® Advance® (11 g lactose/100 kcal).

§ Due to lactose sensitivity.

Availability: Hospital/In:	stitutional
----------------------------	-------------

Size	Container	List No.
Custom Feedin	g System	
Ready To Fe	ed: (20 Cal/fl oz)	
2 fl oz (59 m	L) plastic bottle; 48/case	67325

Availability: Ret	ail	
Size	Container	List No.
Ready to Feed:		
2 fl oz (59 mL)	plastic bottle; 12/case	67564
2 fl oz (59 mL)	plastic bottle; 24/case	67622
8 fl oz (237 mL)	plastic bottle; 24/case	67517
32 fl oz (946 mL)	plastic bottle; 6/case	64253
Powder: (with mea		
20.1 oz (570 g); yie	elds 145 fl oz	
	container; 4/case	
29.8 oz (845 g); yi	elds 225 fl oz container; 4/case	66441

Ready to Feed: Water, Maltodextrin, Milk Protein Isolate, High Oleic Safflower Oil, Sugar, Soy Oil, Coconut Oil; Less than 0.5% of: C. Cohnii Oil, M. Alpina Oil, Short-Chain Fructooligosaccharides, Beta-Carotene, Lutein, Calcium Phosphate, Potassium Citrate, Carrageenan, Potassium Chloride, Ascorbic Acid, Monoglycerides, Soy Lecithin, Magnesium Chloride, Calcium Carbonate, Choline Chloride, Magnesium Phosphate, Sodium Citrate, Ferrous Sulfate, Taurine, Inositol, Choline Bitartrate, Zinc Sulfate, L-Carnitine, Niacinamide, d-Alpha-Tocopheryl Acetate, Calcium Pantothenate, Riboflavin, Copper Sulfate, Vitamin A Palmitate, Thiamine Hydrochloride, Pyridoxine Hydrochloride, Folic Acid, Potassium Iodide, Manganese Sulfate, Phylloquinone, Biotin, Sodium Selenate, Vitamin D₃, Vitamin B₁₂, Potassium Hydroxide, and Nucleotides (Adenosine 5'-Monophosphate, Cytidine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate). Contains milk and soy ingredients.

NUTRITION INF	ORMATION				
	100 Cal	1000 mL		100 Cal	1000 mL
Energy, Cal	100	676	Pantothenic Acid, mcg	450	3041
Volume, mL	148	1000	Biotin, mcg	4.4	29.7
Protein, g	2.1	14.19	Vitamin C, mg	9	60.8
% Total Cal	8	8	Choline, mg	24	162.2
Source	Milk protein isolate		Inositol, mg	24	162.2
Fat, g	5.4	36.49	Minerals		
% Total Cal	49	49	Calcium, mg	84	568
Source	High oleic safflower, soy, and coconut of	ils (0.15% DHA, 0.4% ARA)	Calcium, mEq	4.2	28.4
Oil Ratio	40:30:29		Phosphorus, mg	56	378
Linoleic Acid, mg	1000	6757	Magnesium, mg	6	40.5
Carbohydrate, g	10.9	73.65	Iron, mg	1.8	12.16
% Total Cal	43	43	Zinc, mg	0.75	5.07
Source	Maltodextrin, sugar		Manganese, mcg	5	33.8
Ratio	80:20		Copper, mcg	90	608
Prebiotic 2'-FL hu	man milk oligosaccharides, fructooligo	saccharides	lodine, mcg	15	101
Vitamins			Selenium, mcg	2	14
Vitamin A, IU	300	2027	Sodium, mg	30	203
Vitamin D, IU	60	405	Sodium, mEq	1	8.8
Vitamin E, IU	1.5	10.1	Potassium, mg	107	723
Vitamin K, mcg	8	54.1	Potassium, mEq	2.7	18.2
Thiamin (Vit B ₁), mcg	100	676	Chloride, mg	65	439
Riboflavin (Vit B2), mcg	150	1014	Chloride, mEq	1.8	12.2
Vitamin B ₆ , mcg	60	405	Other Characteristics		
Vitamin B ₁₂ , mcg	0.25	1.69	PRSL, mOsm	19.7	133.2
Niacin, mcg	1050	7095	Water, g	133	899
Folic Acid, mcg	15	101.4	Osmolality, mOsm/kg H ₂ O	200	200

C. Cohnii Oil is a source of DHA.

M. Alpina Oil is a source of ARA.

Similac Sensitive®

Infant Formula with Iron



Description/Indications

A **20 Cal/fl oz**, nutritionally complete infant feeding that is an alternative to standard milk-based formulas. Complete nutrition for fussiness,* gas,* or mild spit-up. Suitable for infants with lactose sensitivity.[†]

Features

- Easy-to-digest gentle carbohydrate for fussiness and gas due to lactose sensitivity
- OptiGRO® is our exclusive blend of DHA, Lutein, and Vitamin E; these important ingredients are found in breast milk
 - DHA for brain and eye development
 - Lutein to support eye health
- Vitamin E, an important nutrient found in breast milk to support developing cells
- Our exclusive formula has:
- Calcium for strong bones no palm olein oil
- Nucleotides to help support the immune system
- Prebiotics to help promote digestive health
- Carotenoids like those naturally found in breast milk
- Clinically shown to support normal growth in infants¹

- A unique blend of carbohydrates using two different absorption pathways to help maximize absorption and minimize malabsorption risks
- Low osmolality (200 mOsm/kg water)
- Gluten-free
- Kosher, Halal

Precaution

- Not for infants or children with galactosemia
- 1. Lasekan JB, et al. Clin Pediatr 2011;50:330-337.
- * Due to lactose sensitivity.
- [†] Typical Value 0.1 g lactose/100 kcal compared to Similac® Advance® (11 g lactose/100 kcal).

Availability: Hospital/Institutional

Size	Container	List No.
Custom Feeding System	1	
Concentrated Liquid	d: (40 Cal/fl oz)	
13 fl oz (384 mL)	. can; 12/case	57535
Availability: Reta	il	
Size	Container	List No.
Ready To Feed: (20	Cal/fl oz)	
32 fl oz (946 mL)	. plastic bottle; 6/case	57533
	,	
Powder: (with measu	3 17	F7500
(0,,,,	ds 90 fl oz container; 6/d	
20.1 oz (570 g); yiel	ds 145 fl oz container; 6	/case 68082
29.8 oz (845 g); yiel	ds 215 fl oz container; 6	/case 62951
34 oz (964 g); yields	245 fl oz container; 2 c	t67550
34.9 oz (989 g); yiel	ds 252 fl oz container; 3	/case 64440

Ready to Feed: Water, Maltodextrin, Milk Protein Isolate, High Oleic Safflower Oil, Sugar, Soy Oil, Coconut Oil. Less than 0.5% of: C. Cohnii Oil, M. Alpina Oil, Short-Chain Fructooligosaccharides, Beta-Carotene, Lutein, Calcium Phosphate, Potassium Citrate, Carrageenan, Potassium Chloride, Ascorbic Acid, Monoglycerides, Soy Lecithin, Magnesium Chloride, Calcium Carbonate, Choline Chloride, Magnesium Phosphate, Sodium Citrate, Ferrous Sulfate, Taurine, Inositol, Choline Bitartrate, Zinc Sulfate, L-Carnitine, Niacinamide, d-Alpha-Tocopheryl Acetate, Calcium Pantothenate, Riboflavin, Copper Sulfate, Vitamin A Palmitate, Thiamine Hydrochloride, Pyridoxine Hydrochloride, Folic Acid, Potassium Iodide, Manganese Sulfate, Phylloquinone, Biotin, Sodium Selenate, Vitamin D₃, Vitamin B₁₂, Potassium Hydroxide and Nucleotides (Adenosine 5'-Monophosphate, Cytidine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate). Contains milk and soy ingredients.

NUTRITION IN	FORMATION				
	100 Cal	1000 mL		100 Cal	1000 mL
Energy, Cal	100	676	Pantothenic Acid, mcg	450	3041
Volume, mL	148	1000	Biotin, mcg	4.4	29.7
Protein, g	2.1	14.19	Vitamin C, mg	9	60.8
% Total Cal	8	8	Choline, mg	24	162.2
Source	Milk protein isolate		Inositol, mg	24	162.2
Fat, g	5.4	36.49	Minerals		
% Total Cal	49	49	Calcium, mg	84	568
Source	High oleic safflower, soy, and coconut oils	(0.15% DHA, 0.4% ARA)	Calcium, mEq	4.2	28.4
Oil Ratio	40:30:29		Phosphorus, mg	56	378
Linoleic Acid, mg	1000	6757	Magnesium, mg	6	40.5
Carbohydrate, g	10.9	73.65	Iron, mg	1.8	12.16
% Total Cal	43	43	Zinc, mg	0.75	5.07
Source	Maltodextrin, sugar		Manganese, mcg	5	33.8
Ratio	80:20		Copper, mcg	90	608
Prebiotic	Fructooligosaccharides		lodine, mcg	15	101
Vitamins			Selenium, mcg	2	14
Vitamin A, IU	300	2027	Sodium, mg	30	203
Vitamin D, IU	60	405	Sodium, mEq	1	8.8
Vitamin E, IU	1.5	10.1	Potassium, mg	107	723
Vitamin K, mcg	8	54.1	Potassium, mEq	2.7	18.5
Thiamin (Vit B ₁), mcg	100	676	Chloride, mg	65	439
Riboflavin (Vit B ₂), mcg	150	1014	Chloride, mEq	1.8	12.2
Vitamin B _e , mcg	60	405	Other Characteristics		
Vitamin B ₁₂ , mcg	0.25	1.69	PRSL, mOsm	19.7	133.2
Niacin, mcg	1050	7095	Water, g	133	899
Folic Acid, mcg	15	101.4	Osmolality, mOsm/kg H ₂ O	200	200

C. Cohnii Oil is a source of DHA.

M. Alpina Oil is a source of ARA.

Similac Pro-Total Comfort®

Infant Formula with Iron



Description/Indications

A **20 Cal/fl oz**, nutritionally complete, non-GMO* infant formula that has partially hydrolyzed whey protein as an alternative to standard intact-protein, milk-based formulas. Designed to support both immune and cognitive development with key ingredients found in breast milk. Suitable for lactose sensitivity. A good choice for infants who may have difficulty tolerating other milk-based formulas.

Features

- Similac Pro-Total Comfort with 2'-FL Human Milk Oligosaccharide[†] helps strengthen the immune system to be more like the breastfed infant than ever before.¹⁻³
- Easy to digest; gentle protein and prebiotics
- OptiGRO[®] is our exclusive blend of DHA, Lutein, and vitamin E; these important ingredients are found in breast milk:
 - DHA for brain and eye development
 - Lutein to support eye health
 - Vitamin E; an important nutrient found in breast milk to support developing cells
- · Our exclusive formula has:
 - Partially hydrolyzed protein for easy digestion
 - A unique blend of carbohydrates using two different

absorption pathways to help maximize absorption and minimize malabsorption risks

- Calcium for strong bones; no palm olein oil
- Nucleotides to help support the immune system
- Prebiotics to help promote digestive health
- Gluten-free
- Halal
- Kosher

Precaution

• Not for infants or children with galactosemia

- 1. Goehring KC, et al. J Nutr. 2016;146(12):2559-2566.
- Schaller JP, et al. Pediatr Res. 2004;56(6):883-890.
- 3. Buck RH, et al. Pediatr Res. 2004;56(6):891-900.

Availability: Hospital/Institutional

Containor

Size	Container	LIST NO.
Ready to Feed:		
2 fl oz (59 mL).	48/case	67132
32 fl oz (946 m	L)6/case	68224
Availability: I	Retail	
Size	Container	List No.
Ready to Feed:		
2 fl oz (59 mL).	24/case	67683
32 fl oz (946 m	L)6/case	68224
20.1 oz (570 g) 29.8 oz (845 g)	neasuring scoop) ; yields 144 fl oz container; 4 ; yields 214 fl oz container; 4 yields 258 fl oz container; 3/	/case 66937

I tot No.

^{*} Ingredients not genetically engineered.

[†] Not from human milk.

Ready to Feed: Water, Corn Syrup Solids, Whey Protein Hydrolysate, High Oleic Safflower Oil, Sugar, Soy Oil, Coconut Oil, Less than 0.5% of: C. Cohnii Oil, M. Alpine Oil, 2'-Fucosyllactose, Short-Chain Fructooligosaccharides, Beta-Carotene, Lutein, Soy Lecithin, Calcium Citrate, Potassium Phosphate, Calcium Phosphate, Carrageenan, Ascorbic Acid, Magnesium Chloride, Monoglycerides, Salt, Inositol, Choline Chloride, Potassium Hydroxide, Potassium Chloride, Ferrous Sulfate, Choline Bitartrate, Taurine, Zinc Sulfate, L-Carnitine, d-Alpha-Tocopheryl Acetate, Niacinamide, Calcium Pantothenate, Potassium Citrate, Vitamin A Palmitate; Cupric Sulfate, Thiamine Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Manganese Sulfate, Potassium Iodide, Phylloquinone, Biotin, Sodium Selenate, Vitamin D_y, Cyanocobalamin, and Nucleotides (Adenosine 5'-Monophosphate, Cytidine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate). Contains milk and soy ingredients.

NUTRITION INF	ORMATION				
	100 Cal	1000 mL		100 Cal	1000 mL
Energy, Cal	100	676	Pantothenic Acid, mcg	450	3041
Volume, mL	148	1000	Biotin, mcg	4.4	29.7
Protein, g	2.32	15.68	Vitamin C, mg	9	60.8
% Total Cal	8	8	Choline, mg	24	162.2
Source	Whey protein hydrolysate		Inositol, mg	24	162.2
Fat, g	5.4	36.49	Minerals		
% Total Cal	49	49	Calcium, mg	105	709
Source	High oleic safflower, soy, and cocor	nut oils (0.15% DHA, 0.4% ARA)	Calcium, mEq	5.25	35.8
Oil Ratio	40:30:29		Phosphorus, mg	75	507
Linoleic Acid, mg	1000	6757	Magnesium, mg	6	40.5
Carbohydrate, g	10.7	72.3	Iron, mg	1.8	12.16
% Total Cal	43	43	Zinc, mg	0.75	5.07
Source	Corn syrup solids, sugar		Manganese, mcg	5	33.8
Ratio	80:20		Copper, mcg	90	608
Prebiotic	2'-FL human milk oligosaccl	harides, fructooligosaccharides	lodine, mcg	15	101
Vitamins			Selenium, mcg	2	14
Vitamin A, IU	300	2029	Sodium, mg	44	297
Vitamin D, IU	60	405	Sodium, mEq	1.9	12.8
Vitamin E, IU	1.5	10.1	Potassium, mg	118	797
Vitamin K, mcg	8	54.1	Potassium, mEq	3	20.3
Thiamin (Vit B ₁), mcg	100	676	Chloride, mg	65	439
Riboflavin (Vit B ₂), mcg	150	1014	Chloride, mEq	1.8	12.2
Vitamin B ₆ , mcg	60	405	Other Characteristics		
Vitamin B ₁₂ , mcg	0.25	1.69	PRSL, mOsm	22.4	151.4
Niacin, mcg	1050	7095	Water, g	133	899
Folic Acid, mcg	15	101.4	Osmolality, mOsm/kg H ₂ O	200	200

C. Cohnii Oil is a source of DHA.

M. Alpina Oil is a source of ARA.

Similac Total Comfort®

Infant Formula with Iron



Description/Indications

A **20 Cal/fl oz**, nutritionally complete infant formula that has partially hydrolyzed whey protein as an alternative to standard intact-protein, milk-based formulas. Suitable for lactose sensitivity; a good choice for infants who may have difficulty tolerating other milk-based formulas.

Features

- Easy to digest; gentle protein and prebiotics
- OptiGRO® is our exclusive blend of DHA, Lutein, and Vitamin E; these important ingredients are found in breast milk
 - DHA for brain and eye development
 - Lutein to support eye health
 - Vitamin E, an important nutrient found in breast milk to support developing cells
- · Our exclusive formula has:
 - Calcium for strong bones no palm olein oil
 - Partially hydrolyzed protein for easy digestion
 - Nucleotides to help support the immune system
 - Prebiotics to help promote digestive health

- A unique blend of carbohydrates using two different absorption pathways to help maximize absorption and minimize malabsorption risks
- Clinically shown to support normal infant growth¹
- · Gluten-free
- · Kosher, Halal

Precaution

- Not for infants or children with galactosemia
- Borschel MW, et al. Clin Pediatr 2014;53:1375-1382.

Availability: Retail

Size	Container	List No.
Powder: (with measuring scoop)	
12.6 oz (3	357 g); yields 90 fl oz can; 6/case	62599

Powder: Corn Maltodextrin, Whey Protein Hydrolysate, High Oleic Safflower Oil, Sugar, Soy Oil, Coconut Oil; Less than 2% of: C. Cohnii Oil, M. Alpina Oil, Short-Chain Fructooligosaccharides, Beta-Carotene, Lutein, Calcium Citrate, Potassium Phosphate, Calcium Phosphate, Potassium Hydroxide, Satt, Magnesium Chloride, Ascorbic Acid, Choline Chloride, Potassium Chloride, Sodium Citrate, Calcium Hydroxide, Potassium Chloride, Incostel, Choline Bitartrate, Zinc Sulfate, Ascorbyl Palmitate, L-Carnitine, Niacinamide, Mixed Tocopherols, d-Alpha-Tocopheryl Acetate, Calcium Pantothenate, Copper Sulfate, Vitamin A Palmitate, Thiamine Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Manganese Sulfate, Potassium Iodide, Phylloquinone, Biotin, Sodium Selenate, Vitamin B 12: and Nucleotides (Adenosine 5'-Monophosphate, Cytidine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate). Sontains milk ingredients.

	100 Cal	1000 mL		100 Cal	1000 mL
Energy, Cal	100	676	Pantothenic Acid, mcg	450	3043
Volume, mL	148	1000	Biotin, mcg	4.4	29.8
Protein, g	2.32	15.69	Vitamin C, mg	9	60.9
% Total Cal	8	8	Choline, mg	24	162.2
Source	Whey protein hydrolysate		Inositol, mg	24	162.2
Fat, g	5.4	36.52	Minerals		
% Total Cal	49	49	Calcium, mg	105	710
Source	High oleic safflower, soy, and coconut oils (0.15%	DHA, 0.4% ARA)	Calcium, mEq	5.25	35.5
Oil Ratio	0:30:29		Phosphorus, mg	75	507
Linoleic Acid, mg	1000	6763	Magnesium, mg	6	40.6
Carbohydrate, g	10.7	72.36	Iron, mg	1.8	12.17
% Total Cal	43	43	Zinc, mg	0.75	5.07
Source	Maltodextrin, sugar		Manganese, mcg	5	33.8
Ratio	80:20		Copper, mcg	90	609
Prebiotic	Fructooligosaccharides		lodine, mcg	15	101
Vitamins			Selenium, mcg	2	14
Vitamin A, IU	300	2029	Sodium, mg	44	298
Vitamin D, IU	60	406	Sodium, mEq	1.9	12.8
Vitamin E, IU	1.5	10.1	Potassium, mg	118	798
Vitamin K, mcg	8	54.1	Potassium, mEq	3	20.3
Thiamin (Vit B,), m	cg 100	676	Chloride, mg	65	440
Riboflavin (Vit B ₂),	mcg 150	1014	Chloride, mEq	1.8	12.2
Vitamin B ₆ , mcg	60	406	Other Characteristics		
Vitamin B ₁₂ , mcg	0.25	1.69	PRSL, mOsm	22.4	151.4
Niacin, mcg	1050	7101	Water, g	133	897
Folic Acid, mcg	15	101.4	Osmolality, mOsm/kg H _o O	200	200

C. Cohnii Oil is a source of DHA.

M. Alpina Oil is a source of ARA.

Similac® For Spit-Up Non-GMO®

Infant Formula with Iron



Description/Indications

A **20 Cal/fl oz**, nutritionally complete, non-GMO, milk-based infant formula with added rice starch to help reduce frequent spit-up; suitable for lactose sensitivity.[†]

Features

- A non-GMO option for complete nutrition for baby's first year
- Easy-to-digest formula with added rice starch
- Clinically shown to reduce frequent spit-up in healthy babies by 54%^{1‡}
- OptiGRO® is our exclusive blend of DHA, LUTEIN, and VITAMIN E; these important ingredients are found in breast milk
 - DHA for brain and eye development
 - Lutein to support eye health
 - Vitamin E, an important nutrient found in breast milk to support developing cells
- Our exclusive formula has:
 - Calcium for strong bones-no palm olein oil
 - Nucleotides to help support the immune system
 - Prebiotics to help promote digestive health
 - Carotenoids like those naturally found in breast milk

- A unique blend of carbohydrates using two different absorption pathways to help maximize absorption and minimize malabsorption risks
- Clinically shown to support normal infant growth¹
- · Gluten-free
- · Kosher, Halal

Precaution

- Not for infants or children with galactosemia
- 1. Lasekan JB, et al. J Am Coll Nutr 2014;33(2):136-146.
- * Ingredients not genetically engineered.
- † Typical Value 0.1 g lactose/100 kcal compared to Similac® Advance® (11 g lactose/100 kcal).
- [‡] Among healthy 2-month-old infants compared to a standard formula.

Availability: Hospital/Institutional

Ready to Feed: Water, Corn Syrup, Rice Starch, Milk Protein Isolate, High Oleic Safflower Oil, Soy Oil, Sugar, Coconut Oil, Galactooligosaccharides; Less than 0.5% of: C. Cohnii Oil, M. Alpina Oil, Beta-Carotene, Lutein, Calcium Phosphate, Calcium Carbonate, Potassium Chloride, Potassium Phosphate, Ascorbic Acid, Magnesium Chloride, Potassium Citrate, Soy Lecithin, Monoglycerides, Carrageenan, Inositol, Choline Chloride, Potassium Hydroxide, Ferrous Sulfate, Taurine, Choline Bitartrate, Zinc Sulfate, L-Carnitine, Niacinamide, d-Alpha-Tocopheryl Acetate, Calcium Pantothenate, Riboflavin, Vitamin A Palmitate, Copper Sulfate, Thiamine Hydrochloride, Pyridoxine Hydrochloride, Potassium Iodide, Folic Acid, Phylloquinone, Biotin, Sodium Selenate, Vitamin D₃, Vitamin B₁₂₂, and Nucleotides (Adenosine 5'-Monophosphate, Cytidine 5'-Monophosphate), Contains milk and soy ingredients.

NUTRITION I	NFORMATION				
	100 Cal	1000 mL		100 Cal	1000 mL
Energy, Cal	100	676	Pantothenic Acid, mcg	450	3041
Volume, mL	148	1000	Biotin, mcg	4.4	29.7
Protein, g	2.1	14.19	Vitamin C, mg	9	60.8
% Total Cal	8	8	Choline, mg	24	162.2
Source	Milk protein isolate		Inositol, mg	24	162.2
Fat, g	5.4	36.49	Minerals		
% Total Cal	49	49	Calcium, mg	84	568
Source	High oleic safflower, soy, and coconut oils (0.15	5% DHA, 0.4% ARA)	Calcium, mEq	4.2	28.4
Oil Ratio	40:30:29		Phosphorus, mg	56	378
Linoleic Acid, mg	1000	6757	Magnesium, mg	6	40.5
Carbohydrate, g	11	74.32	Iron, mg	1.8	12.16
% Total Cal	43	43	Zinc, mg	0.75	5.07
Source	Corn syrup, rice starch, sugar		Manganese, mcg	5	33.8
Ratio	50:30:20		Copper, mcg	90	608
Prebiotic	Galactooligosaccharides		lodine, mcg	15	101
Vitamins			Selenium, mcg	2	14
Vitamin A, IU	300	2027	Sodium, mg	30	203
Vitamin D, IU	60	405	Sodium, mEq	1.3	8.8
Vitamin E, IU	1.5	10.1	Potassium, mg	107	723
Vitamin K, mcg	8	54.1	Potassium, mEq	2.7	18.2
Thiamin (Vit B,), mcg	100	676	Chloride, mg	65	439
Riboflavin (Vit B ₂), mcg	g 150	1014	Chloride, mEq	1.8	12.2
Vitamin B ₆ , mcg	60	405	Other Characteristics		
Vitamin B ₁₂ , mcg	0.25	1.69	PRSL, mOsm	19.7	133.2
Niacin, mcg	1050	7095	Water, g	133	899
Folic Acid, mcg	15	101.4	Osmolality, mOsm/kg H ₂ O	180	180

C. Cohnii Oil is a source of DHA.

M. Alpina Oil is a source of ARA.

Similac® Alimentum®

Infant Formula with Iron



Description/Indications

A **20 Cal/fl oz**, nutritionally complete, hypoallergenic formula for infants, including those with colic symptoms due to cow's milk protein sensitivity, food allergies, or sensitivity to intact protein. Appropriate supplemental beverage for children with food allergies or sensitivity to intact protein.

Features

- First and only hypoallergenic extensively hydrolyzed casein-based infant formula with 2'-FL for immune support.
- Starts reducing excessive crying due to cow's milk protein sensitivity in most infants within 24 hours¹
- Hypoallergenic; contains a predigested protein to virtually eliminate allergic reactions in most babies who are allergic to cow's milk protein
- Hydrolyzed casein supplemented with free amino acids for infants who are sensitive to or unable to digest intact protein
- A unique blend of carbohydrates using two different absorption pathways to help maximize absorption and minimize malabsorption risks
- Made with oils that have been shown to be well absorbed.² Approximately 33% of the fat as medium

- chain triglycerides (MCTs)
- DHA and ARA, special nutrients found in breast milk that are important for brain and eye development³⁻⁵
- Suitable for lactose sensitivity
- · Gluten-free
- · Ready-To-Feed formula is corn-free
- Based on a clinical study with Similac Alimentum Ready To Feed without DHA and ARA, lutein, beta-carotene, and 2'-FL HMO in a small group of infants experiencing colic symptoms due to cow's milk protein sensitivity. Data on file, Study AC84, August 2004. Abbott. Columbus, Ohio.
- Fomon SJ. Nutrition of Normal Infants, St. Louis, Mosby-Year Book, Inc; 1993:152-153.
- 3. Auestad N, et al. Pediatrics 2001;108:372-381.
- Auestad N, et al. Pediatrics 2003;112:e177-e183.
- Scott DT, et al. Pediatrics 1998;102(5):E59.
- *WIC is a service mark of the US Department of Agriculture, and an abbreviation for the Special Supplemental Nutrition Program for Women, Infants, and Children. No endorsement of any brand or product by the USDA is implied or intended. At least one form of Similac® Alimentum® is available through WIC in all 50 states with a valid medical referral.

Availability: Hospital/Institutional

Size	Container	List No.
Custom Feeding Sy		
Ready To Feed: (20	,	
2 fl oz (59 mL)	. plastic bottle; 48/case	59738
Availability: Reta	il	
Size	Container	List No.
Ready To Feed: (20	Cal/fl oz)	
2 fl oz (59 mL)	. plastic bottle; 24/case	67703
32 fl oz (946 mL)	. plastic bottle; 6/case	57512
Powder: (with measu	uring scoop)	
12.1 oz (343 g); yiel	ds 87 fl oz can; 6/case	64715
19.8 oz (561 g); yiel	ds 143 fl oz container; 4/	case64719

Ready To Feed: Water (87%), Sugar (4%), Casein Hydrolysate [Derived from Milk] (2%), Modified Tapioca Starch (2%), Safflower Oil (1%), Medium Chain Triglycerides (1%), Soy Oil (1%). Less than 0.5% of: 2"-Fucosyllactose [Derived from Milk]*, Lutein, Beta-Carotene, C. Cohnii Oil, M. Alpina Oil, Calcium Citrate, Calcium Phosphate, Carrageenan, Ascorbic Acid, Potassium Phosphate, Magnesium Chloride, Potassium Citrate, Salt, Calcium Hydroxide, Potassium Dibrydrochloride, Potassium Chloride, Potassium Chloride, Potassium Chloride, Forous Sulfate, Taurine, Inositol, d-Alpha-Tocopheryl Acetate, Zinc Sulfate, L-Carnitine, Niacinamide, Calcium Pantothenate, Vitamin A Palmitate, Copper Sulfate, Thiamine Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Potassium Iodide, Phylloquinone, Biotin, Sodium Selenate, Vitamin D., and Vitamin B.,...

Powder: Corn Maltodextrin (35%), Casein Hydrolysate [Derived from Milk] (18%), Sugar (15%), High Oleic Safflower Oil (10%), Medium Chain Triglycerides (10%), Soy Oil (8%). Less than 2% of: 2'-Fucosyllactose (Derived from Milk], Lutein, Beta-Carotene, C. Cohnii Oil, M. Alpina Oil, Calcium Phosphate, DATEM, Potassium Citrate, Xanthan Gum, Magnesium Chloride, Monoglycerides, Ascorbic Acid, Salt, L-Cystine Dihydrochloride, Calcium Carbonate, Potassium Chloride, L-Tyrosine, L-Tryptophan, Choline Chloride, Ferrous Sulfate, Talmine, Niacinamide, Mixed Tocopherols, Calcium Pantothenate, Copper Sulfate, Vitamin A Palmitate, Thiamine Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Potassium Iodide, Potassium Hydroxide, Phylloquinone, Biotin, Sodium Selenate, Vitamin D₁, and Vitamin B₁,

	100 Cal	1000 mL		8 fl oz (237mL)	1000 mL
Energy, Cal	100	676	Pantothenic Acid, mcg	750	5072
Volume, mL	148	1000	Biotin, mcg	4.5	30.4
Protein, g	2.75	18.6	Vitamin C, mg	9	61
% Total Cal	11	11	Choline, mg	12	81
Source	Casein hydrolysate, L-cystine, L-tyre	osine, L-tryptophan	Inositol, mg	5	33.8
Fat, g	5.54	37.47	Minerals		
% Total Cal	48	48	Calcium, mg	105	710
	ium chain triglycerides, and soy oil (0.15		Calcium, mEq	5.2	35.4
Oil Ratio	38:33:2	-	Phosphorus, mg	75	507
Linoleic Acid, mg	1900	12,850	Magnesium, mg	7.5	50.7
Carbohydrate, g	10.2	69	Iron, mg	1.8	12.17
% Total Cal	41	41	Zinc, mg	0.75	5.07
Source	Sugar, modified tapioca starch		Manganese, mcg Copper, mcg	8	54
Ratio		70:30:00		75	507
Prebiotics	2'-FL human milk oli	gosaccharides	lodine, mcg	15	101
Vitamins			Selenium, mcg	2	14
Vitamin A, IU	300	2029	Sodium, mg	44	298
Vitamin D, IU	45	304	Sodium, mEq	1.9	12.9
Vitamin E, IU	3	20.3	Potassium, mg	118	798
Vitamin K, mcg	15	101	Potassium, mEq	3	20.3
Thiamin (Vit B1), mcg	60	406	Chloride, mg	80	541
Riboflavin (Vit B2), mcg	90	609	Chloride, mEq	2.3	15.5
Vitamin B6, mcg	60	406	Other Characteristics		
Vitamin B12, mcg	0.45	3.04	PRSL, mOsm	25.3	171.3
Niacin, mcg	1350	9130	Water, g	133	899
Folic Acid, mcg	15	101	Osmolality, mOsm/kg H ₂ O	370	370

EleCare® (for Infants)

Amino Acid-Based Infant Formula with Iron





Description/Indications

A 20 Cal/fl oz, nutritionally complete amino acid-based formula for infants who cannot tolerate intact or hydrolyzed protein. EleCare is indicated for the dietary management of protein maldigestion, malabsorption, severe food allergies, short-bowel syndrome, eosinophilic Gl disorders, Gl-tract impairment, or other conditions in which an amino acid-based diet is required.

- For infants 0-12 months of age
- For oral or tube feeding
- Use under medical supervision

Features

- Hypoallergenic^{1,*}—virtually eliminating the potential for an alleroic reaction to the formula in multiple-food-alleroic children
- Clinically shown to support the growth of exclusively formula fed infants^{2,*}
- 100% free amino acids as nitrogen source
- DHA and ARA, nutrients that help support brain and eye development
- 33% of fat blend as medium-chain triglycerides,

- an easily digested and well-absorbed fat source
- Supported by strict manufacturing standards and ELISA (Enzyme-linked ImmunoSorbent Assay) testing
- Does not contain milk protein, soy protein, fructose, galactose, lactose, or gluten
- Halal
- Kosher-made with pareve ingredients
- Sicherer SH, et al. J Pediatr. 2001;138:688-693.
- 2. Borschel MW, et al. Clin Pediatr. 2013;52(10):910-917.
- * Study conducted with a previous formulation of EleCare unflavored without DHA/ARA.
- † The WIC (Special Supplemental Nutrition Program for Women, Infants, and Children) name is a service mark of the US Department of Agriculture. USDA does not endorse any goods, services, or enterprises. Requires WIC state medical documentation form. WIC state approval is subject to change, check with your WIC agency on availability.

Availability

Powder: Corn Syrup Solids (55%), High Oleic Safflower Oil (9%), Medium-Chain Triglycerides (8%), Soy Oil (7%), L-Glutamine (2%). Less than 2% of: C. Cohnii Oil, M. Alpina Oil, L-Asparagine, L-Lecinie, L-Lysine Acetate, DATEM, Calcium Phosphate, L-Valine, Potassium Dhosphate, L-Isoleucine, L-Arginine, L-Prevallanine, L-Tyrosine, L-Threonine, Potassium Citrate, Sodium Citrate, L-Proline, L-Serine, L-Alanine, Glycine, L-Histidine, L-Methionine, Ascorbic Acid, Magnesium Carbonate, L-Cystine Dihydrochloride, L-Tryptophan, Salt, Choline Chloride, Inositol, Ferrous Sulfate, Taurine, Ascorbyl Palmitate, Zinc Sulfate, dl-Alpha-Tocopheryl Acetate, L-Carnitine, Niacinamide, Calcium Pantothenate, Thiamine Hydrochloride, Copper Sulfate, Manganese Sulfate, Vitamin A Palmitate, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Beta-Carotene, Biotin, Phylloquinone, Chromium Chloride, Potassium Iodide, Sodium Molybdate, Vitamin D, Vitamin B, Vitamin

	100 Cal	100 g powder	1000 mL [†]		100 Cal	100 g powder	1000 mL [†]
Calories	100	475	676	Vitamin C (Ascorbic Acid), mg	9	43	61
Protein, q	3.1	14.5	21	Choline, mg	15	71	101
% Total Cal ¹	15¹	15¹	15¹	Inositol, mg	5.1	24	35
Source		Free L-amino acids		Minerals			
Fat, g	4.8	23	32.5	Calcium, mg	116	550	785
% Total Cal	43	43	43	Calcium, mEq	5.8	27.5	39.2
Linoleic Acid, mg	840	4000	5683	Phosphorus, mg	84.2	400	570
MCT:LCT	33:67	33:67	33:67	Magnesium, mg	8.4	40	56.8
Source	High oleic safflower oi	I, MCTs, soy oil, (0.15% D	HA; 0.40% ARA)	Iron, mg	1.8	8.4	12.2
Carbohydrate, g	10.7	51	72.4	Zinc, mg	1.15	5.5	7.8
% Total Cal	42	42	42	Manganese, mcg	84	400	568
Source		Corn syrup solids		Copper, mcg	126	599	852
Vitamins				lodine, mcg	8.9	42	60
Vitamin A, IU	273	1300	1847	Selenium, mcg	2.6	12.6	17.6
Vitamin A, mcg RE	82	390	554	Chromium, mcg	2.3	11	15.6
Vitamin D, IU	60	285	406	Molybdenum, mcg	2.5	12	17.1
Vitamin D, mcg	1.5	7.1	10.15	Sodium, mg	45	215	304
Vitamin E, IU	2.1	10	14.2	Sodium, mEq	2.0	9.4	13.2
Vitamin K, mcg	13	60	88	Potassium, mg	150	715	1015
Thiamine (Vit B ₄), mcg	210	1000	1421	Potassium, mEq	3.9	18.3	26
Riboflavin (Vit B ₂), mcg	105	500	710	Chloride, mg	60	285	406
Vitamin B ₆ , mcg	84.2	400	570	Chloride, mEq	1.7	8.0	11.5
Vitamin B ₁₂ , mcg	0.4	2.0	2.7	Other Characteristics			
Niacin, mcg	1680	8000	11,366	Water, g	132		895
Folic Acid (Folacin), mcg	29.5	140	200	Osmolality, mOsm/kg H ₂ O			350
Pantothenic Acid, mcg	421	2000	2848	PRSL, mOsm	28	131	187
Biotin, mcg	4.2	20	28.4				

[†] Prepared at 20 Cal/fl oz.

M. Alpina Oil is a source of ARA. www.abbottnutrition.com

^{1.} May ME, et al. Am J Clin Nutr 1990;52:770-776.

C. Cohnii Oil is a source of DHA.

Similac[®] PM 60/40

Low-Iron Infant Formula



Description/Indications

A **20 Cal/fl oz** formula for infants who would benefit from lowered mineral intake, including those with impaired renal function.

Features

- Mineral levels closely approximating the mineral content of human milk (60:40 ratio of whey to casein)
- Calcium-to-phosphorus ratio and content designed to manage serum calcium disorders—both hypercalcemia and hypocalcemia due to hyperphosphatemia
- · Gluten-free
- Kosher, Halal

Precautions

- Additional iron should be supplied from other sources
- In conditions where the infant is losing abnormal quantities
 of one or more electrolytes, it may be necessary to supply
 electrolytes from sources other than the formula
- It may be necessary to supply low-birth-weight infants weighing less than 1500 g at birth additional calcium, phosphorus, and sodium during periods of rapid growth
- Not for infants or children with galactosemia

Availability

Size	Container	List No.
Powder: (with	h measuring scoop)	
14.1 oz (400	g); yields 102 fl oz can; 6/case	00850

Powder: Lactose, High Oleic Safflower Oil, Whey Protein Concentrate, Soy Oil, Coconut Oil, Sodium Caseinate. Less than 2% of: Potassium Citrate, Calcium Phosphate, Calcium Carbonate, Ascorbic Acid, Magnesium Chloride, Potassium Chloride, m-Inositol, Salt, Choline Chloride, Taurine, Ascorbyl Palmitate, Ferrous Sulfate, Zinc Sulfate, Mixed Tocopherols, L-Carnitine, Niacinamide, d-Alpha-Tocopheryl Acetate, Calcium Pantothenate, Cupric Sulfate, Vitamin A Palmitate, Thiamine Chloride Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Beta-Carotene, Folic Acid, Manganese Sulfate, Potassium Iodide, Phylloquinone, Biotin, Sodium Selenate, Vitamin D₃, Cyanocobalamin and Potassium Hydroxide. Contains milk ingredients.

	100 Cal	1000 mL		100 Cal	1000 mL
Francis Oct		***	Dietie men		
Energy, Cal	100	676	Biotin, mcg	5	30.4
Volume, mL	148	1000	Vitamin C, mg	9	61
Protein, g	2.2	15	Choline, mg	12	81
% Total Cal	9	9	Inositol, mg	24	162
Source	Whey protein concentrate and sodium cas		Minerals		
Fat, g	5.6	37.9	Calcium, mg	56	379
% Total Cal	50	50	Calcium, mEq	2.8	18.9
Source	High oleic safflower, soy, and coconut	oils	Phosphorus, mg	28	189
Oil Ratio	41:30:29	41:30:29	Magnesium, mg	6	40.6
Linoleic Acid, mg	1000	6763	Iron, mg	0.7	4.7
Carbohydrate, g	10.2	69.0	Zinc, mg	0.8	5.1
% Total Cal	41	41	Manganese, mcg	5	34
Source	Lactose		Copper, mcg	90	609
Vitamins			lodine, mcg	6	41
Vitamin A, IU	300	2029	Selenium, mcg	2	14
Vitamin D, IU	60	406	Sodium, mg	24	162
Vitamin E, IU	1.5	10.1	Sodium, mEq	1.0	7.1
Vitamin K, mcg	8	54	Potassium, mg	80	541
Thiamin (Vit B.), mcg	100	676	Potassium, mEq	2.1	13.8
Riboflavin (Vit B ₂), mcg	150	1014	Chloride, mg	59	399
Vitamin B _e , mcg	60	406	Chloride, mEa	1.7	11.3
Vitamin B ₁₂ , mcg	0.3	1.7	Other Characteristics		
Niacin, mcg	1050	7101	PRSL, mOsm	18.3	124.1
Folic Acid, mcg	15	101	Water, q	134	899
Pantothenic Acid, mca	450	3043	Osmolality, mOsm/kg H ₂ O	280	280





Nutrient Composition of Preterm Human Milk

Preterm Human Milk*								
	100 Cal	1000 mL		100 Cal	1000 mL 107			
Energy, Cal	100	671	Vitamin C, mg	16				
Volume, mL	149	1000	Choline, mg	14	94			
Protein, g	2.10	14.09	Inositol, mg	22.0	147.7			
% Total Cal	8	8	Minerals					
Source	Preterm Human Milk	Preterm Human Milk	Calcium, mg	37	248			
Fat, g	5.80	38.93	Calcium, mEq	1.9	12.4			
% Total Cal	52	52	Phosphorus, mg	19	128			
Source	Preterm Human Milk	Preterm Human Milk	Magnesium, mg	4.6	30.9			
Linoleic Acid, mg	550	3691	Iron, mg	0.18	1.21			
Carbohydrate, g	9.9	66.4	Zinc, mg	0.51	3.42			
% Total Cal	40	40	Manganese, mcg	1	6			
Source	Preterm Human Milk	Preterm Human Milk	Copper, mcg	96	644			
Vitamins			lodine, mcg	16	107			
Vitamin A, IU	581	3899	Selenium, mcg	2.2	14.8			
Vitamin D, IU	3	20	Sodium, mg	37	248			
Vitamin E, IU	1.6	10.7	Sodium, mEq	1.6	10.8			
Vitamin K, mcg	0.3	2.0	Potassium, mg	85	570			
Thiamin (Vit B ₁), mcg	31	208	Potassium, mEq	2.2	14.6			
Riboflavin (Vit B ₂), mcg	72	483	Chloride, mg	82	550			
Vitamin B ₆ , mcg	22	148	Chloride, mEq	2.3	15.6			
Vitamin B ₁₂ , mcg	0.07	0.47	Other Characteristics					
Niacin, mcg	224	1503	PRSL, mOsm	18.7	125.6			
Folic Acid, mcg	5	33	Water, g	131	879			
Pantothenic Acid, mcg	269	1805	Approx Osmolality,	290	290			
Biotin, mcg	0.6	4.0	mOsm/kg H ₂ O					

^{*} Composition of preterm human milk varies with maternal diet, stage of lactation, within feedings, diurnally, and among mothers. Values for term human milk have been used for linoleic acid, biotin, choline, inositol, manganese, iodine, and selenium.¹ Values represent mature preterm milk (not colostrum or transitional milk).

^{1.} Meeting the Special Nutrient Needs of Low-Birth-Weight and Premature Infants in the Hospital (A8100). Columbus, Ohio: Abbott, Abbott Laboratories, January 1998:56.

Human Milk Handling Guidelines[†]

Expressing and transporting human milk

- Begin mechanical expression as soon as possible after giving birth, with the use of a clean, hospital-grade, electric breast pump; recommend double pumping.
- Wash hands prior to expressing or pumping milk.
- Properly clean the personal collection kit after each use and sanitize it daily, as instructed.
- Store human milk in "food-grade" hard plastic containers or glass; these containers should have an airtight seal.
- Label human milk supplied to the facility with complete and accurate information, including the infant's name, medical record number, and date and time of pumping to prevent errors in human milk delivery.
- Maintain human milk transported to and from the hospital at proper temperatures (<4°C, <39°F) to prevent loss of nutrients and to minimize bacterial growth.

Storing human milk

- Provide dedicated freezers and refrigerators for storing human milk; store food in a separate refrigerator.
- Store milk from each mother in a separate plastic bin or sealed plastic bag to prevent misadministration and crosscontamination.
- Maintain refrigerator temperatures at ≤4°C (≤39°F); freezer temperatures at ≤-20°C (≤-4°F). Use thawed human milk within 24 hours. Do not refreeze thawed human milk.
- Fresh human milk can be safely stored in the refrigerator for 48 to 72 hours.
- Fortified human milk should be stored in the refrigerator and used within 24 hours.
- Frozen human milk can be safely stored in a home freezer for 3 to 6 months at ≤-18°C (≤0°F) and in a freezer with lower temperatures of ≤-20°C (≤-4°F) for 6 to 12 months.

Feeding expressed human milk

- Bottles, nipples, and graduated feeders should be for single use.
- Expect that the milk will separate during storage because it is not homogenized. The cream will rise to the top of the milk and look thicker and whiter.
- Before feeding, gently swirling the container of milk will mix the cream back through again. Avoid vigorously shaking the milk.
- The color of milk may vary from day to day, depending on maternal diet. It may look bluish, yellowish, or brownish. Frozen human milk may also smell different than fresh human milk.
- Use frozen human milk in the order in which it was expressed (oldest milk first).
- Thaw containers of human milk in the refrigerator (if frozen) or under warm running water. Never use a microwave oven or hot water to warm or thaw human milk.
- Do not add warm human milk to frozen milk because it will partially thaw the frozen milk.
- Warming is not recommended for continuous feedings. Warming time for oral or bolus feedings should be limited to no more than 15 minutes.
- Acceptable methods for warming include electric warming units and warm running water. Water level should not reach the level of the nipple ring or submerge the lid.
- Discard any feeding remaining in the bottle after 1 hour for infants being nipple fed.

†Adapted from:

Pediatric Nutrition Practice Group, Steele C, Collins E, eds. Infant and Pediatric Feedings: Guidelines for Preparation of Human Milk and Formula in Health Care Facilities, 3rd ed. Chicago: Academy of Nutrition and Dietetics. 2019, Pp. 6-7, 17.

Lawrence RA, Lawrence RM. Breastfeeding: A Guide for the Medical Profession, 8th ed. Philadelphia: Elsevier, 2016, Pp. 315, 712-742.

Premature Infant Fortifiers and Formulas

Comprehensive Nutrition Options for Preterm Infants

Low-birth-weight and preterm infants arrive with unique feeding challenges, which can continue in hospital and post-discharge.

Insufficient amounts of nutrients, an imbalance of nutrients, or their poor bioavailability may prevent premature infants from achieving their full growth potential.

Human milk is the optimal form of infant nutrition and the preferred source of enteral nutrition for preterm infants, primarily due to its immunologic properties. However, despite its documented advantages, human milk alone does not meet all the nutritional needs of preterm infants, who often require supplementation to meet their unique nutritional needs.¹

 American Academy of Pediatrics Committee on Nutrition. Nutritional Needs of the Preterm Infant. In: Kleinman RE, Greer FR, eds. Pediatric Nutrition. 8th ed. Itasca, IL: American Academy of Pediatrics; 2019:142.

Abbott: A Tradition of Innovation Supporting the Preterm Infant

Abbott provides a number of solutions to support the nutritional needs of low-birth-weight and preterm infants, including a flexible strategy to help human milk meet the high nutrient demands of preterm infants.

Nutritional products specifically designed for preterm infants:

- Over 50 years of research
- Over 100 scientific publications

Extensive product line for individualized feedings:

- Similac® Human Milk Fortifiers (Concentrated Liquids and Powder) for use as a nutritional supplement to add to human milk
- Similac® Special Care® family of milkbased liquid formulas 20, 24, 24 High Protein, and 30 Cal/fl oz suitable for mixing using the Liqui-Mix® System
- Similac® NeoSure®, a nutrient-enriched (increased protein, vitamins, and minerals compared to term infant formula), post-discharge, milk-based, preterm infant formula available in powder and ready-to-feed forms
- Liquid Protein Fortifier, a commercially sterile liquid protein to fortify NICU formula and human milk

These formulations are specially designed to meet the unique nutritional requirements of most preterm infants in the NICU and can be fed with confidence.

Similac® Human Milk Fortifier Hydrolyzed Protein Concentrated Liquid



Description/Indications

Intended for premature and low-birth-weight infants as a nutritional supplement to add to human milk.

• Use under medical supervision

Features

- Clinical study shows improved growth for your littlest babies¹⁻³
- Extensively hydrolyzed protein for easy digestion and absorption
- Non-acidified
- . DHA for brain and eye development
- Lutein to support eye development
- When added to human milk, meets expert recommendations for protein^{4,5*} and other nutrients for the premature infant⁴
- · Well tolerated
- Small, convenient packet is designed for easy mixing
- Commercially sterile and meets the AND and CDC recommendation to use liquid for NICU feedings^{6,7} †
- · Low iron level provides flexibility to add iron as needed
- Gluten-free

Precautions

- Add only to human milk—do not add water
- This product is nutritionally incomplete by itself and is designed to be added to human breast milk
- · Additional iron may be necessary
- Tolerance to enteral feedings should be confirmed by offering small volumes of unfortified human milk
- Once enteral feeding is well established, Similac Human Milk Fortifier Hydrolyzed Protein Concentrated Liquid can be added to human milk
- Not intended for low-birth-weight infants after they reach a weight of 3600 g (approximately 8 lb) or as directed by a physician

Preparation and Use

Follow directions as specified on carton. Improper dilution may be harmful.

- 1. Kim JH, et al. J Pediatr Gastroenterol Nutr. 2015;61:665-671.
- 2. Schanler RJ, et al. J Pediatr. 2018;202:31 27.
- 3. Kumar N, et al. Am J Perinatol. 2017;34:1411 1416.
- 4. Klein CJ. J Nutr. 2002;132:1395S-1577S.
- 5. 5. Agostoni C, et al.J Pediatr Gastroenterol Nutr.2010;58:85-91.
- Centers for Disease Control and Prevention. Enterobacter sakazakii infections associated with the use of powdered infant formula—Tennessee, 2001. Available at http://www.cdc.gov/mmw/ preview/mmw/html/mm5114a1.htm. Published April 11, 2002. Accessed November 27, 2019.
- Pediatric Nutrition Practice Group.Steele C, Collins E, eds. Infant and Pediatric Feedings: Guidelines for Preparation of Human Milk and Formula in Health Care Facilities. 3rd ed. Chicago, IL: Academy of Nutrition and Dietetics; 2019:12.

Availability: Hospital/Institutional

Concentrated Liquid	trated Liquid List N .17-fl-oz) packet144/case630	
5-mL (0.17-fl-oz) packet	144/case	63010

^{*} One packet mixed with 25 mL of human milk.

[†] Academy of Nutrition and Dietetics and Centers for Disease Control and Prevention.

Concentrated Liquid: Water, Corn Maltodextrin, Casein Hydrolysate. Less than 2% of: M. Alpina Oilt; C. Cohnii Oilš, Lutein, Beta-Carrotene, Calcium Phosphate, Magnesium Chloride, Potassium Citrate, Sodium Citrate, Sodium Chloride, Potassium Phosphate, Zinc Sulfate, Ferrous Sulfate, Copper Sulfate, Manganese Sulfate, Sodium Selenite, Medium Chain Triglycerides, Soy Oil, Coconut Oil, Modified Corn Starch, Vitamin A Palmitate, Vitamin D₃, d-Alpha-Tocopheryl Acetate, Phylloquinone, Thiamine Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Vitamin B₁₂, Niacinamide, Folic Acid, Calcium Pantothenate, Biotin, Ascorbic Acid, m-Inositol and Choline Chloride, Monoglycerides, L-Tyrosine, L-Leucine, L-Tryptophan, L-Carnitine, Gellan Gum and Potassium Hydroxide. Contains milk ingredients.

Pe	r 5 mL packet	Per 30 mL‡	Per 100 mL	Per 100 Cal [§]		Per 5 mL packet	Per 30 mL‡	Per 100 mL	Per 100 Cal [§]
Energy, Cal	7	24	79	100	Vitamin C, mg	7.7	10.4	34.6	43.7
Volume, mL	5	30	100	126	Choline, mg	0.9	3.3	10.8	13.7
Protein, g	0.5	0.85	2.84	3.58	Inositol, mg	1.7	5.4	18	22.7
% Total Cal	-	15	15	15	Minerals				
Source Preterm human milk, casein hydrolysate					Calcium, mg	30	36	121	152
Fat, g	0.21	1.18	3.94	4.98	Calcium, mEq	1.5	1.8	6	7.6
% Total Cal	-	44	44	44	Phosphorus, mg	17	20	67	85
Source	Preterm huma	an milk, medium-c	hain triglycerides, so	y and coconut oils	Magnesium, mg	2.1	2.8	9.6	12.1
Linoleic Acid, mg	18	110	368	464	Iron, mg	0.11	0.14	0.47	0.59
Carbohydrate, g	0.75	2.41	8.04	10.14	Zinc, mg	0.31	0.39	1.32	1.66
% Total Cal	-	41	41	41	Manganese, mcg	2.2	2.4	7.8	9.9
Source Preterm human milk, maltodextrin, modified cornstarch					Copper, mcg	15	31	104	131
Vitamins					lodine, mcg	0.4	3.1	10	13
Vitamin A, IU	197	295	982	1238	Selenium, mcg	0.2	0.5	2	2
Vitamin D, IU	35	36	118	149	Sodium, mg	5	12	37	47
Vitamin E, IU	1	1.3	4.2	5.3	Sodium, mEq	0.2	0.5	1.6	2
Vitamin K, mcg	2.4	2.5	8.2	10.3	Potassium, mg	21	35	118	148
Thiamin (Vit. B,), mcg	48	53	177	224	Potassium, mEq	0.5	0.9	2.9	3.6
Riboflavin (Vit. B ₂), mo	cg 74	86	287	362	Chloride, mg	13	27	89	113
Vitamin B ₆ , mcg	50	53	179	226	Chloride, mEq	0.4	0.8	2.6	3.3
Vitamin B ₁₂ , mcg	0.13	0.14	0.47	0.6	Other Characterist	tics			
Niacin, mcg	980	1010	3392	4279	PRSL, mOsm	5.1	8.4	25.6	32.2
Folic Acid, mcg	7	8	26.1	32.9	Water, g	3.3	25.3	84	106
Pantothenic Acid, mc	g 310	355	1184	1494	Approx. Osmolality	у,			
Biotin, mcg	5.8	5.9	19.7	24.8	mOsm/kg H₂O	NA	450	450	450

[‡]One packet (5 mL) Similac Human Milk Fortifier Concentrated Liquid in 25 mL preterm human milk. Preterm human milk values from Meeting the Special Needs of Low-Birth Weight and Premature Infants in the Hospital. Columbus, OH: Abbott, Abbott Laboratories, January 1998.

[§] Per 100 Calories, as mixed, 4 packets added to 100 mL preterm human milk (24 Cal/fil oz). Preterm human milk values from Meeting the Special Needs of Low-Birth Weight and Premature Infants in the Hospital. Columbus, OH: Abbott, Abbott Laboratories, January 1998.

Values based on term human milk from American Academy of Pediatrics Committee on Nutrition: Pediatric Nutrition Handbook, 4th ed. Elk Grove Village, American Academy of Pediatrics, 1998;40:132-135, 217, 258, 655-658.

M. Alpina Oil is a source of ARA.C. Cohnii Oil is a source of DHA.

Similac® Human Milk Fortifier Concentrated Liquid





Description/Indications

Intended for premature and low-birth-weight infants as a nutritional supplement to add to human milk.

• Use under medical supervision

Features

- Small, convenient packet is designed for easy mixing
- When added to human milk, meets the nutrient recommendations for the premature infant¹
- Commercially sterile and meets the AND and CDC recommendation to use liquid for NICU feedings^{2,3*}
- Packet is simple to open and mixes easier with human milk than powder⁴
- Low iron level provides flexibility to add iron as needed
- · Kosher, Halal

Precautions

- Add only to human milk—do not add water
- This product is nutritionally incomplete by itself and is designed to be added to human breast milk
- · Additional iron may be necessary

- Tolerance to enteral feedings should be confirmed by offering small volumes of unfortified human milk
- Once enteral feeding is well established, Similac Human Milk Fortifier Concentrated Liquid can be added to human milk
- Not intended for low-birth-weight infants after they reach a weight of 3600 g (approximately 8 lb) or as directed by a physician

Preparation and Use

Follow directions as specified on carton. Improper dilution may be harmful.

- 1. Klein CJ. J Nutr. 2002;132(6):1395S-1577S.
- Centers for Disease Control and Prevention. Enterobacter sakazakii infections associated with the use of powdered infant formula – Tennessee, 2001. Available at http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5114a1.htm. Published April 11, 2002. Accessed November 27, 2019.
- Pediatric Nutrition Practice Group. Steele C, Collins E, eds. Infant and Pediatric Feedings: Guidelines for Preparation of Human Milk and Formula in Health Care Facilities. 3rd ed. Chicago, IL: Academy of Nutrition and Dietetics; 2019:12.
- 4. Data on file, 2010. Abbott Market Research, Abbott Laboratories, Columbus. Ohio.

^{*} Academy of Nutrition and Dietetics and Centers for Disease Control and Prevention.

Ingredients

Concentrated Liquid: Water, Nonfat Milk, Corn Syrup Solids, Medium-Chain Triglycerides, Whey Protein Concentrate. Less than 2% of: Minerals (Calcium Phosphate, Magnesium Chloride, Potassium Citrate, Salt, Calcium Carbonate, Potassium Phosphate, Sodium Citrate, Magnesium Phosphate, Zinc Sulfate, Ferrous Sulfate, Cupric Sulfate, Manganese Sulfate and Sodium Selenate), Vitamins (Vitamin A Palmitate, Vitamin D₃, d-Alpha-Tocopheryl Acetate, Phylloquinone, Thiamine Chloride Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Cyanocobalamin, Niacinamide, Folic Acid, Calcium Pantothenate, Biotin, Ascorbic Acid, m-Inositol and Choline Chloride), Soy Lecithin, M. Alpina Oil, C. Cohnii Oil and Potassium Hydroxide.

Contains milk and soy Ingredients.

P	er 5 mL packet	Per 30 mL [†]	Per 100 mL	Per 100 Cal [‡]		Per 5 mL packet	Per 30 mL [†]	Per 100 mL	Per 100 Cal [‡]
Energy, Cal	7	24	79	100	Vitamin C, mg	7.7	10.4	34.6	43.7
Volume, mL	5	30	100	126	Choline, mg	0.9	3.3 [§]	10.8	13.7
Protein, g	0.35	0.70	2.34	2.95	Inositol, mg	1.7	5.4 [§]	18	22.7
% Total Cal	-	12	12	12	Minerals				
Source	Preterm human	milk, nonfat milk,	whey protein concen	trate	Calcium, mg	35	41	137	173
Fat, g	0.27	1.24	4.14	5.23	Calcium, mEq	1.8	2.1	7	8.9
% Total Cal	-	46	46	46	Phosphorus, mg	20	23	77	98
Source	Preterm hu	man milk, medium	-chain triglycerides		Magnesium, mg	2.2	2.9	9.9	12.5
Linoleic Acid, mg	1	93 [§]	311	392	Iron, mg	0.11	0.14	0.47	0.59
Carbohydrate, g	0.81	2.47	8.24	10.4	Zinc, mg	0.30	0.39	1.3	1.62
% Total Cal	-	42	42	42	Manganese, mcg	2.1	2.3 [§]	7.5	9.5
Source	Pret	erm human milk, c	orn syrup solids		Copper, mcg	15	31	104	131
Vitamins					lodine, mcg	0.5	3.2 [§]	11	13
Vitamin A, IU	197	294	982	1238	Selenium, mcg	0.2	0.6§	2	2
Vitamin D, IU	35	35	118	149	Sodium, mg	5	12	37	47
Vitamin E, IU	1.0	1.2	4.2	5.3	Sodium mEq	0.23	0.5	1.7	2.1
Vitamin K, mcg	2.4	2.5	8.2	10.3	Potassium, mg	21	35	118	148
Thiamin (Vit. B,), mc	g 48	53	177	224	Potassium, mEq	0.53	0.9	3	3.8
Riboflavin (Vit. B ₂), m	ncg 123	135	450	568	Chloride, mg	13	27	89	113
Vitamin B ₆ , mcg	49	53	176	222	Chloride, mEq	0.38	0.8	2.6	3.2
Vitamin B ₁₂ , mcg	0.08	0.09	0.31	0.39	Other Characterist	ics			
Niacin, mcg	1040	1080	3592	4532	PRSL, mOsm	3.8	6.9	23.1	29.3
Folic Acid, mcg	7	8	26.1	32.9	Water, g	3.6	25.6	85	108
Pantothenic Acid, m	cg 309	354	1180	1489	Osmolality,				
Biotin, mcg	7.6	7.7§	25.7	32.4	mOsm/kg H₂O	NA	385	385	385

[†] One packet (5 mL) Similac Human Milk Fortifier Concentrated Liquid in 25 mL preterm human milk. Preterm human milk values from Meeting the Special Needs of Low-Birth Weight and Premature Infants in the Hospital. Columbus, OH: Abbott, Abbott Laboratories, January 1998.

[‡] Per 100 Calories, as mixed, 4 packets added to 100 mL preterm human milk (24 Cal/fi oz). Preterm human milk values from Meeting the Special Needs of Low-Birth Weight and Premature Infants in the Hospital. Columbus, OH: Abbott, Abbott Laboratories, January 1998.

[§] Values based on term human milk from America Academy of Pediatrics Committee on Nutrition: Pediatric Nutrition Handbook, 4th ed. Elk Grove Village, American Academy of Pediatrics, 1998:40; 132-135, 217, 258, 655-658.

M. Alpina Oil is a source of ARA.

C. Cohnii Oil is a source of DHA.

		Mul	Helit Golli	parison of Pretern		ommac opecial	Jaie 30 (336 30)		
					Per 100 Cal		Per 100 mL		
Nutrients		lutrient ements† Max	Preterm Human Milk (PTHM)‡	PTHM + SSC 30 (2:1 ratio) [§]	PTHM + SSC 30 (1:1 ratio)§	PTHM + SSC 30 (1:2 ratio) [§]	PTHM + SSC 30 (2:1 ratio) [§]	PTHM + SSC 30 (1:1 ratio)§ +	PTHM + SSC 30 (1:2 ratio) [§]
Cal/fl oz				23	25	27	23	25	27
Energy, Cal	100	100	100	100	100	100	79	84	90
Volume, mL	-	100	149	127	119	111	100	100	100
Protein, q	2.5	3.6	2.1	2.5	2.6	2.8	2.0	2.2	2.5
Source	2.0	0.0	2.1	2.0		eterm human milk, nonfat n			2.0
Fat, q	4.4	5.7	5.8	6.1	6.3	6.4	4.8	5.3	5.8
Source	4.4	5.7	5.0	0.1	Preterm	human milk, medium chair			3.0
Carbohydrate, q	9.6	12.5	9.9	9.0	8.6	8.3	7.0	7.2	7.4
Source Source	3.0	12.0	5.5	3.0	0.0		orn syrup solids, lactose	1.2	7.4
Vitamins				i	ı	I	I syrup sonus, ruotosc		
Vitamin A, IU	700	1254	581	869	984	1084	683	829	975
Vitamin D. IU	75	270	3	66	91	113	52	77	102
Vitamin E, IU	2.0	8.0	1.6	2.6	3.0	3.4	2.1	2.6	3.1
Vitamin E, IO Vitamin K, mcg	4.0	25	0.3	5	3.0 7	9	4	6	8
	30	250	31	125	163	196	98	137	176
Thiamin (Vit B ₁), mcg			72	308	163 402	196 484	242	339	435
Riboflavin (Vit B ₂), mcg	80 30	620	22	120			94	134	174
Vitamin B ₆ , mcg		250			159	193			
Vitamin B ₁₂ , mcg	0.08 550	0.7	0.07	0.28	0.36 3098	0.43	0.22	0.30	0.39
Niacin, mcg		5000	224	2280		3813	1791	2611	3432
Folic Acid, mcg	30	45	5	19	24	29	15	20	26
Pantothenic Acid, mcg	300	1900	269	971	1251	1495	763	1054	1345
Biotin, mcg	1.0	37.0	0.6	16.3	22.5	28.0	12.8	19.0	25.2
Vitamin C, mg	8	37	16	25.0	28.6	31.8	19.7	24.1	28.6
Choline, mg	7	23	14 22	12	12	11 36	10 23	10	10
Inositol, mg	4	44	22	30	33	36	23	28	32
Minerals									
Calcium, mg	123	185	37	99	123	144	77	104	130
Phosphorus, mg	82	109	19	54	68	80	42	57	72
Magnesium, mg	6.8	17.0	4.6	7.8	9.1	10.2	6.1	7.6	9.1
Iron, mg	1.7	3.0	0.18	0.9	1.15	1.4	0.7	1.0	1.3
Zinc, mg	1.1	1.5	0.51	0.94	1.11	1.25	0.74	0.93	1.13
Manganese, mcg	6	25	1	6	8	9	4	6	8
Copper, mcg	100	250	96	162	189	212	127	159	191
lodine, mcg	6	35	16	12	10	8	9	8	8
Selenium, mcg	2	5	2	2	2	2	2	2	2
Sodium, mg (mEq)	39	63	37 (1.61)	40 (1.72)	41 (1.77)	42 (1.80)	31 (1.35)	34 (1.49)	37 (1.62)
Potassium, mg (mEq)	60	160	85 (2.17)	104 (2.66)	111 (2.85)	118 (3.02)	82 (2.09)	94 (2.40)	106 (2.72)
Chloride, mg (mEq)	60	160	82 (2.32)	82 (2.30)	81 (2.30)	81 (2.30)	64 (1.81)	69 (1.94)	73 (2.07)
Other Characteristics									
PRSL. mOsm	-	-	18.7	22.6	24.2	25.6	17.8	20.4	23.0
Approx Osmolality.	-	-	290	302	310	313	302	310	313
mOsm/kg H ₂ O					- 70				-10

[†] Life Sciences Research Office recommendations for preterm and low-birth-weight infants.² Visit http://www.lsro.org/articles/lowbirthweight_rpt.pdf for a summary of the report. **NOTE:** These guidelines are designed for preterm infant formulas, NOT fortified human milk. Thus, these recommendations can serve only as a general guideline relative to fortified human milk.

[‡] The nutrient concentrations listed for preterm human milk are mean human milk levels. The range of nutrients in preterm human milk is highly variable. Therefore, actual nutrient levels may be higher or lower than the levels listed. Human milk values were obtained from: Meeting the Special Nutrient Needs of Low-Birth-Weight and Premature Infants in the Hospital (A8100). Columbus, OH-L Abbott, Abbott, Laboratories, January 1998.

[§] When combined with human milk, does not increase concentration of nutrients to levels achieved with Similac® Human Milk Fortifier.

High-Calorie Human Milk Fortification Options

Similac® Human Milk Fortifier Hydrolyzed Protein Concentrated Liquid (SHMF HP CL) + Similac® Special Care® 30 (SSC 30)

Nutrients	Preterm Human Milk	PTHM + SHMF HP CL = Fortified	FHM + SSC 30	FHM + SSC 30	FHM + SSC 30	SSC 30
Per 100 Cal	(PTHM) ¹	Human Milk (FHM)	+ SSC 30 26 Cal	+ SSC 30 27 Cal +	28 Cal + 1	<u> </u>
Cal/fl oz	20	24	26	27	28	30
Ratios		Mix 1 packet SHMF HP CL + 25 mL PTHM	Mix 2 parts FHM + 1 part SSC 30	Mix 1 part FHM + 1 part SSC 30	Mix 1 part FHM + 2 parts SSC 30	
Energy, Cal	100	100	100	100	100	100
Volume, mL	149	126	115	111	106	99
Protein, g	2.1	3.58	3.36	3.26	3.16	3
Fat, g	5.8	4.98	5.61	5.89	6.15	6.61
Carbohydrate, g	9.9	10.14	9.2	8.79	8.41	7.73
Vitamins						
Vitamin A, IU	581	1238	1243	1245	1247	1250
Vitamin D, IU	3	149	150	150	150	150
Vitamin E, IU	1.6	5.3	4.8	4.6	4.4	4
Vitamin K, mcg	0.3	10.3	11	11.3	11.5	12
Thiamin (Vit B ₁), mcg	31	224	234	238	243	250
Riboflavin (Vit B ₂), mcg	72	362	462	507	548	620
Vitamin B ₆ , mcg	22	226	235	239	243	250
Vitamin B ₁₂ , mcg	0.07	0.6	0.58	0.57	0.56	0.55
Niacin, mcg	224 4.9	4279 32.9	4560 34.5	4683 35.2	4798 35.8	5000 37
Folic Acid, mcg						
Pantothenic Acid, mcg	269 0.6	1494 24.8	1652 29.6	1721 31.6	1786 33.6	1900 37
Biotin, mcg Vitamin C, mg	16	24.8 43.7	29.6 41.1	31.6	33.6	37
	14	13.7	12.2	11.6	36.9	10
Choline, mg	22	22.7	29.4	32.4	35.1	40
Inositol, mg	22	22.1	29.4	32.4	35.1	40
Minerals						
Calcium, mg	37	152	163	168	172	180
Phosphorus, mg	19	85	91	93	96	100
Magnesium, mg	4.6	12.1	12	12	12	12
Iron, mg	0.18	0.59	1.06	1.27	1.46	1.8
Zinc, mg	0.51	1.66	1.6	1.57	1.55	1.5
Manganese, mcg	0.9	9.9	10.7	11.1	11.4	12
Copper, mcg	96	131	177	198	217	250
lodine, mcg	16	13	10	9	8	6
Selenium, mcg	2	2	2	2	2	2
Sodium, mg (mEq)	37 (1.6)	47 (2)	46 (1.9)	45 (1.9)	44 (1.9)	43 (1.9)
Potassium, mg (mEq)	85 (2.2)	148 (3.6)	141 (3.5)	137 (3.4)	134 (3.4)	129 (3.3)
Chloride, mg	82 (2.3)	113 (3.3)	100 (2.9)	95 (2.7)	90 (2.6)	81 (2.3)
Other Characteristics						
PRSL, mOsm	18.7	32.2	30.7	30	29.3	27.8
Approx Osmolality, mOsm/kg H ₂ O	290	450	410	390	365	325

^{1.} Meeting the Special Nutrient Needs of Low-Birth-Weight and Premature Infants in the Hospital (A8100). Columbus, Ohio: Abbott, Abbott Laboratories, January 1998.

High-Calorie Human Milk Fortification Options Similac® Human Milk Fortifier Hydrolyzed Protein Concentrated Liquid (SHMF HP CL) + Similac® Special Care® 30 (SSC 30)

Ollilliao II	aman mik rorumor	niyardiy2ea i roteiii oo	ilioolitiatoa Eigala (oli	ini in or, i onimuo	opeoidi odio oo (
Nutrients Per 100 mL	Preterm Human Milk (PTHM)¹	PTHM + SHMF HP CL = Fortified Human Milk (FHM)	FHM + SSC 30 26 Cal +	FHM + SSC 30 27 Cal +	FHM + SSC 30 28 Cal +	SSC 30
Cal/fl oz	20	24	26	27	28	30
Ratios		Mix 1 packet SHMF HP CL + 25 mL PTHM	Mix 2 parts FHM + 1 part SSC 30	Mix 1 part FHM + 1 part SSC 30	Mix 1 part FHM + 2 parts SSC 30	
Energy, Cal	67	79	87	90	94	101
Volume, mL	100	100	100	100	100	100
Protein, g	1.41	2.84	2.9	2.94	2.98	3.04
Fat, g	3.89	3.94	4.85	5.31	5.78	6.71
Carbohydrate, q	6.64	8.04	7.96	7.92	7.91	7.84
Vitamins						
Vitamin A, IU	390	982	1075	1122	1173	1268
Vitamin D, IU	2	118	129	135	141	152
Vitamin E, IU	1.1	4.2	4.2	4.1	4.1	4.1
Vitamin K, mcg	0.2	8.2	9.5	10.1	10.8	12.2
Thiamin (Vit B.), mcg	21	177	202	215	228	254
Riboflavin (Vit B.), mcg	48	287	400	457	515	629
Vitamin B _{et} mcg	15	179	203	216	229	254
Vitamin B,,, mcg	0.05	0.47	0.5	0.51	0.53	0.56
Niacin, mcg	150	3392	3945	4221	4512	5072
Folic Acid, mcg	3.3	26.1	29.8	31.7	33.7	37.5
Pantothenic Acid, mcg	181	1184	1429	1551	1680	1927
Biotin, mcg	0.4	19.7	25.6	28.5	31.6	37.5
Vitamin C, mg	10.7	34.6	35.5	36	36.6	37.5
Choline, mg	9.4	10.8	10.6	10.5	10.4	10.1
Inositol, mg	14.8	18	25.4	29.2	33	40.6
Minerals						
Calcium, mg	25	121	141	151	162	183
Phosphorus, mg	13	67	79	84	90	101
Magnesium, mg	3.1	9.6	10.4	10.8	11.3	12.2
Iron, mg	0.12	0.47	0.92	1.14	1.37	1.83
Zinc, mg	0.34	1.32	1.38	1.42	1.45	1.52
Manganese, mcg	0.6	7.8	9.3	10	10.7	12.2
Copper, mcg	64	104	153	178	204	254
lodine, mcg	11	10	9	8	7	6
Selenium, mcg	1	2	2	2	2	2
Sodium, mg (mEq)	25 (1.1)	37 (1.6)	39 (1.7)	40 (1.7)	42 (1.8)	44 (1.9)
Potassium, mg (mEq)	57 (1.5)	118 (2.9)	122 (3)	124 (3.1)	126 (3.2)	131 (3.3)
Chloride, mg (mEq)	55 (1.6)	89 (2.6)	87 (2.5)	86 (2.5)	85 (2.4)	82 (2.3)
Other Characteristics						
PRSL, mOsm	12.6	25.6	26.5	26.9	27.3	28.2
Approx Osmolality,	290	450	410	390	365	325
mOsm/kg H ₂ O						

^{1.} Meeting the Special Nutrient Needs of Low-Birth-Weight and Premature Infants in the Hospital (A8100). Columbus, Ohio: Abbott, Abbott Laboratories, January 1998.

Liquid Protein Fortifier

For Human Milk, Fortified Human Milk or Infant Formula



Description/Indications

The first and only commercially sterile, extensively hydrolyzed liquid protein fortifier for preterm infants who may require additional protein.

Features

- Designed to be mixed with human milk, fortified human milk, or formula feedings with minimal displacement for any infant who needs additional protein
- Helps customize feedings so preterm infants can receive the optimal protein level to meet their specific needs
- Liquid product eliminates the need for powder mixing and meets the AND and CDC* recommendations to reduce risk of contamination^{1,2}
- Extensively hydrolyzed liquid protein fortifier for easy digestion and absorption (the same protein source as Similac® Alimentum®)
- 1 g of protein/6 mL (16.7 g protein/100 mL)
- · Convenient, multi-serving bottle
- Gluten-free
- · Lactose-free

Precaution

 If signs of intolerance develop, consult with infant's healthcare professional.

Preparation and Use

- · Use as directed by a physician
- This product is nutritionally incomplete. Must be mixed with human milk, fortified human milk, or formula before feeding
- Do not attach nipple and ring to Liquid Protein Fortifier bottle
- Do not add water
- Shake gently
- Do not use if breakaway ring is missing or broken
- Enteral use only; not for IV use
- Discard unused dispensed fortifier

Storage

- Store unopened at room temperature; avoid extreme temperatures.
- Fortifier: Refrigerate after opening: use within 72 hours or discard.
- Fortified human milk or formula: Once mixed, refrigerate and use within 24 hours or discard.
- Pediatric Nutrition Practice Group, Steele C, Collins E, eds. Infant and Pediatric Feedings: Guidelines for Preparation of Human Milk and Formula in Health Care Facilities. 3rd ed. Chicago, IL: Academy of Nutrition and Dietetics; 2019:12.
- Centers for Disease Control and Prevention. Enterobacter sakazakii infections
 associated with the use of powdered infant formula—Tennessee, 2001. Available at
 http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5114a1.htm. Published April 11,
 2002. Accessed November 27. 2019.
- * Academy of Nutrition and Dietetics and Centers for Disease Control and Prevention.

Availability: Hospital/Institutional

Size	Container	List No.
1.8 fl oz (54 mL)	. plastic bottle; 48/case	62317

Liquid Protein Fortifier is easy to distinguish from other NICU products

Bottle has several distinguishing characteristics for differentiation:

- Black cap for easy product identification
- Stickers displaying important usage information
- Label has clinical look to stand out from formulas on shelf
- Liquid is opaque and darker than infant formula

Ingredients

Water and Casein Hydrolysate (milk).

NUTRIENTS PER 6 ML	
Calories	4
Protein	1 g
Water	5 g

NUTRITIONAL CHARACTERISTICS OF LIQUID PROTFIN FORTIFIEF

Protein: Source and Concentration → 100% extensively hydrolyzed casein 16.7 g protein/100 mL; 1 g protein/6 mL

Calories → 4 Cal/1 g protein

Osmolality \rightarrow 1 mL contributes \sim 12 mOsm/kg water when added to make 100 mL of the feeding

Displacement → 1 mL will displace 1 mL of feeding





Similac® Special Care® 20

Premature Infant Formula with Iron



Description/Indications

A **20 Cal/fl oz**, iron-fortified feeding for growing, low-birth-weight infants and premature infants.

Features

- OptiGRO® is our exclusive blend of DHA, Lutein, and Vitamin E; these important ingredients are found in breast milk
 - DHA for brain and eye development
 - Lutein to support eye health
 - Vitamin E, an important nutrient found in breast milk to support developing cells
- Lutein, a carotenoid naturally found in colostrum and human milk, has been shown to help support eye development of the preterm infant¹⁻⁴
- Nucleotides for immune support
- Calcium:phosphorus content and ratio (1.8:1) supports intrauterine accretion rates⁵
- Approximately 2 mg iron/kg body weight per day when fed at 120 Cal/kg body weight per day
- Gluten-free
- · Kosher, Halal

Precautions

- Very low-birth-weight infants are particularly susceptible to gastrointestinal complications; therefore, feeding should be initiated cautiously
- Tolerance to enteral feedings should be confirmed by initially offering small volumes of formula followed by cautious progression to higher caloric feedings
- Spitting up, abdominal distention, abnormal stools or stool
 patterns, excessive gastric residuals, or other signs of intestinal
 dysfunction have been associated with enteral feeding before
 the intestinal tract is ready to accommodate the regimen.
 At the first sign of these problems, enteral feeding should be
 slowed or discontinued
- Not intended for feeding low-birth-weight infants after they reach a weight of 3600 g (approximately 8 lb) or as directed by a physician

1. Canfield LM, et al. Eur J Nutr 2003;42:133-141.

2. Schweigert FJ, et al. Eur J Nutr 2004;43:39-44.

3. Patton S, et al. Lipids 1990;25:159-165

Size

4. Rubin LP. et al. J Perinatol 2012; 32(6):418-424.

5. Mize CE, et al. Am J Clin Nutr 1995;62:385-391.

Availability: Hospital/Institutional

Container

List No.

Availability: Retail

Size	Container	List No.
Ready To	Feed: (20 Cal/fl oz)	
2 fl oz (59	mL) plastic bottle; 48/case	56265

Preparation



Ready To Feed:

Do not dilute unless directed by a physician. Can be mixed to a variety of caloric densities using the Liqui-Mix® System (see pages 90-98). Follow physician's instructions.

Ingredients

Ready To Feed: Water, Nonfat Milk, Corn Syrup Solids, Medium Chain Triglycerides, Lactose, Whey Protein Concentrate, Soy Oil, Coconut Oil. Less than 0.5% of: C. Cohnii Oil, M. Alpina Oil, Beta-Carotene, Lutein, Calcium Phosphate, Ascorbic Acid, Magnesium Chloride, Soy Lecithin, Monoglycerides, Potassium Citrate, m-Inositol, Carrageenan, Calcium Carbonate, Sodium Citrate, Potassium Hydroxide, Ferrous Sulfate, Choline Bitartrate, Taurine, Niacinamide, Choline Chloride, L-Carnitine, Zinc Sulfate, Potassium Chloride, Salt, Potassium Phosphate, d-Alpha-Tocopheryl Acetate, Calcium Pantothenate, Vitamin A Palmitate, Cupric Sulfate, Riboflavin, Thiamine Chloride Hydrochloride, Pyridoxine Hydrochloride, Folic Acid, Manganese Sulfate, Biotn, Phylloquinone, Sodium Selenate, Vitamin D., Cyanocobalamin, and Nucleotides (Cytidine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate, Adenosine 5'-Monophosphate). Contains milk and soy ingredients.

	100 Cal	1000 mL		100 Cal	1000
	100 Cal			100 Cal	1000 mL
Energy, Cal	100	676	Biotin, mcg	37	250
Volume, mL	148	1000	Vitamin C, mg	37	250
Protein, g	3.0	20.29	Choline, mg	10	68
% Total Cal	12	12	Inositol, mg	40	271
Source	Nonfat milk, whey protein concent	rate	Minerals		
Fat, g	5.43	36.72	Calcium, mg	180	1217
% Total Cal	47	47	Calcium, mEq	9.0	60.7
	triglycerides, soy, and coconut oils (0.2	25% DHA; 0.40% ARA)	Phosphorus, mg	100	676
Oil Ratio	50:30:18	50:30:18	Magnesium, mg	12	81.2
Linoleic Acid, mg	700	4734	Iron, mg	1.8	12.2
Carbohydrate, g	10.3	69.7	Zinc, mg	1.5	10.14
% Total Cal	41	41	Manganese, mcg	12	81
Source	Corn syrup solids, lactose		Copper, mcg	250	1691
Ratio	50:50	50:50	lodine, mcg	6	41
Vitamins			Selenium, mcg	2	14
Vitamin A, IU	1250	8454	Sodium, mg	43	291
Vitamin D, IU	150	1014	Sodium, mEq	1.9	12.6
Vitamin E, IU	4.0	27.1	Potassium, mg	129	872
Vitamin K, mcg	12	81.2	Potassium, mEq	3.3	22.3
Thiamin (Vit B ₁), mcg	250	1691	Chloride, mg	81	548
Riboflavin (Vit B ₂), mcg	620	4193	Chloride, mEq	2.3	15.5
Vitamin B _s , mcg	250	1691	Other Characteristics		
Vitamin B ₁₂ , mcg	0.55	3.72	PRSL, mOsm	27.8	188.2
Niacin, mcg	5000	33,815	Water, g	133	899
Folic Acid, mcg	37	250	Approx Osmolality,	235	235
Pantothenic Acid, mcg	1900	12,850	mOsm/kg H ₂ O		

C. Cohnii Oil is a source of DHA.

M. Alpina Oil is a source of ARA.

Similac® Special Care® 24

Premature Infant Formula with Iron



Description/Indications

A **24 Cal/fl oz**, iron-fortified feeding for growing, low-birth-weight infants and premature infants.

Features

- OptiGRO® is our exclusive blend of DHA, Lutein, and Vitamin E; these important ingredients are found in breast milk
 - DHA for brain and eye development
 - Lutein to support eye health
 - Vitamin E, an important nutrient found in breast milk to support developing cells
- Clinically shown to improve early language development,^{1*†} early visual development,^{1*‡} and body composition^{2*}
- Lutein, a carotenoid naturally found in colostrum and human milk, has been shown to help support eye development of the preterm infant³⁻⁶
- Nucleotides for immune support
- Calcium:phosphorus content and ratio (1.8:1) supports intrauterine accretion rates⁷
- Approximately 2 mg iron/kg body weight per day when fed at 120 Cal/kg body weight per day
- Gluten-free
- · Kosher, Halal

Precautions

- Very low-birth-weight infants are particularly susceptible to gastrointestinal complications; therefore, feeding should be initiated cautiously
- Tolerance to enteral feedings should be confirmed by initially offering small volumes of formula followed by cautious progression to higher caloric feedings
- Spitting up, abdominal distention, abnormal stools or stool
 patterns, excessive gastric residuals, or other signs of intestinal
 dysfunction have been associated with enteral feeding before
 the intestinal tract is ready to accommodate the regimen.
 At the first sign of these problems, enteral feeding should be
 slowed or discontinued
- Not intended for feeding low-birth-weight infants after they reach a weight of 3600 g (approximately 8 lb) or as directed by a physician
- 1. O'Connor DL, et al. Pediatrics 2001;108:359-371.
- 2. Groh-Wargo S, et al. Pediatr Res 2005;57:712-718.
- 3. Canfield LM, et al. Eur J Nutr 2003;42:133-141.
- 4. Schweigert FJ, et al. Eur J Nutr 2004;43:39-44.
- Patton S, et al. Lipids 1990;25:159-165.
- 6. Rubin LP. et al. *J Perinatol* 2012:32(6):418-424.
- Hubin LP, et al. J Perinatol 2012;32(6):418-424.
 Mize CE, et al. Am J Clin Nutr 1995;62:385-391.
- * Compared to infants fed a formula without DHA and ARA in a clinical trial with Similac Special Care and Similac® NeoSure® infant formulas with iron.
- † Based on a post-hoc analysis of English-speaking singleton premature infants using the MacArthur Communicative Developmental Inventories.

.

‡ Visual acuity measured at 4 and 6 months corrected age and assessed by VEP (visual evoked potential).

Availability: Hospital/Institutional

S	ıze	Container	LIST NO.
С	ustom Feeding Sy	stem	
R	eady to Feed:		
2	fl oz	. plastic bottle; 48/case	67368
Α	vailability: Reta	il	
S	ize	Container	List No.
R	eady To Feed: (24	Cal/fl oz)	
2	fl oz (59 mL)	. plastic bottle; 48/case	67446

Preparation



Ready To Feed:

Do not dilute unless directed by a physician.

Can be mixed to a variety of caloric densities using the Liqui-Mix® System (see pages 90-98). Follow physician's instructions.

Ingredients

Ready To Feed: Water, Nonfat Milk, Corn Syrup Solids, Medium Chain Triglycerides, Lactose, Whey Protein Concentrate, Soy Oil, Coconut Oil. Less than 0.5% of: C. Cohnii Oil, M. Alpina Oil, Beta-Carotene, Lutein, Calcium Phosphate, Ascorbic Acid, Potassium Citrate, Calcium Carbonate, Soy Lecithin, Monoglycerides, Magnesium Chloride, m-Inositol, Sodium Citrate, Carrageenan, Potassium Hydroxide, Ferrous Sulfate, Choline Bitartrate, Taurine, Choline Chloride, Niacinamide, L-Carnitine, Zinc Sulfate, Potassium Chloride, Salt, Potassium Photosphate, d-Alpha-Tocopheryl Acetate, Calcium Pantothenae, Vitamin A Palmitate, Cupric Sulfate, Riboflavin, Thiamine Chloride Hydrochloride, Pyridoxine Hydrochloride, Folic Acid, Manganese Sulfate, Biotin, Phylloquinone, Sodium Selenate, Vitamin D₃, Cyanocobalamin, and Nucleotides (Cytidine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate, Adenosine 5'-Monophosphate). Contains milk and soy ingredients.

	100 Cal	1000 mL		100 Cal	1000 mL
Energy, Cal	100	812	Biotin, mcg	37	300
Volume, mL	124	1000	Vitamin C, mg	37	300
Protein, g	3.0	24.34	Choline, mg	10	81
% Total Cal	12	12	Inositol, mg	40	325
Source	Nonfat milk, whey protein concentra	ate	Minerals		
Fat, g	5.43	44.07	Calcium, mg	180	1461
% Total Cal	47	47	Calcium, mEq	9.0	72.9
	glycerides, soy, and coconut oils (0.2	25% DHA; 0.40% ARA)	Phosphorus, mg	100	812
Oil Ratio	50:30:18	50:30:18	Magnesium, mg	12	97.4
Linoleic Acid, mg	700	5681	Iron, mg	1.8	14.6
Carbohydrate, g	10.3	83.6	Zinc, mg	1.5	12.17
% Total Cal	41	41	Manganese, mcg	12	97
Source	Corn syrup solids, lactose		Copper, mcg	250	2029
Ratio	50:50	50:50	lodine, mcg	6	49
Vitamins			Selenium, mcg	2	16
Vitamin A, IU	1250	10,144	Sodium, mg	43	349
Vitamin D, IU	150	1217	Sodium, mEq	1.9	15.2
Vitamin E, IU	4.0	32.5	Potassium, mg	129	1047
Vitamin K, mcg	12	97.4	Potassium, mEq	3.3	26.8
Thiamin (Vit B ₁), mcg	250	2029	Chloride, mg	81	657
Riboflavin (Vit B ₂), mcg	620	5032	Chloride, mEq	2.3	18.6
Vitamin B ₆ , mcg	250	2029	Other Characteristics		
Vitamin B ₁₂ , mcg	0.55	4.46	PRSL, mOsm	27.8	225.8
Niacin, mcg	5000	40,578	Water, g	109	885
Folic Acid, mcg	37	300	Approx Osmolality,	280	280
Pantothenic Acid, mcg	1900	15,419	mOsm/kg H ₂ O		

C. Cohnii Oil is a source of DHA.

M. Alpina Oil is a source of ARA.

Similac® Special Care® 24 High Protein

Premature Infant Formula with Iron



Description/Indications

A **24 Cal/fl oz**, iron-fortified feeding for growing, low-birthweight infants and premature infants who may need extra protein to help support growth.

Features

- OptiGRO® is our exclusive blend of DHA, Lutein, and Vitamin E; these important ingredients are found in breast milk
 - DHA for brain and eye development
 - Lutein to support eye health
 - Vitamin E, an important nutrient found in breast milk to support developing cells
- 3.3 g of protein/100 Cal, 10% more protein than Similac[®] Special Care[®] 24
- When fed at 120 Cal/kg/day, meets the average recommended 4 g protein/kg/day for very low-birthweigh infants.¹
- Lutein, a carotenoid naturally found in colostrum and human milk, has been shown to help support eye development of the preterm infant²⁻⁵
- Nucleotides for immune support
- Calcium:phosphorus content and ratio (1.8:1) supports intrauterine accretion rates⁶

- Approximately 2 mg iron/kg body weight per day when fed at 120 Cal/kg body weight per day
- Gluten-free
- · Kosher, Halal

Precautions

- Very low-birth-weight infants are particularly susceptible to gastrointestinal complications; therefore, feeding should be initiated cautiously
- Tolerance to enteral feedings should be confirmed by initially offering small volumes of formula followed by cautious progression to higher caloric feedings
- Spitting up, abdominal distention, abnormal stools or stool
 patterns, excessive gastric residuals, or other signs of intestinal
 dysfunction have been associated with enteral feeding before
 the intestinal tract is ready to accommodate the regimen.
 At the first sign of these problems, enteral feeding should be
 slowed or discontinued
- Not intended for feeding low-birth-weight infants after they reach a weight of 3600 g (approximately 8 lb) or as directed by a physician
- American Academy of Pediatrics Committee on Nutrition. Nutritional needs of the preterm Infant. In: Kleinman RE, Greer FR, eds. Pediatric Nutrition. 8th ed. Itasca, IL: American Academy of Pediatrics: 2019:129.
- 2. Canfield LM, et al. Eur J Nutr 2003;42:133-141.
- 3. Schweigert FJ, et al. Eur J Nutr 2004;43:39-44.
- 4. Patton S, et al. Lipids 1990;25:159-165.
- 5. Rubin LP, et al. J Perinatol 2012;32(6):418-424.
- 6. Mize CE, et al. Am J Clin Nutr 1995;62:385-391.

Availability: Hospital/Institutional

 Size
 Container
 List No.

 Custom Feeding System
 Ready to Feed: (24 Cal/fl oz)

 2 fl oz
 plastic bottle; 48/case
 67370

Preparation



Ready To Feed:

Do not dilute unless directed by a physician.

Can be mixed to a variety of caloric densities using the Liqui-Mix® System (see pages 90-98). Follow physician's instructions.

Ingredients

Ready To Feed: Water, Nonfat Milk, Corn Syrup Solids, Medium Chain Triglycerides, Whey Protein Concentrate, Lactose, Soy Oil, Coconut Oil. Less than 0.5% of: C. Cohnii Oil, M. Alpina Oil, Beta-Carotene, Lutein, Calcium Phosphate, Ascorbic Acid, Potassium Citrate, Calcium Caronate, Soy Lecithin, Monoglycerides, Magnesium Chloride, m-Inositol, Sodium Citrate, Carrageenan, Potassium Hydroxide, Ferrous Sulfate, Chloine Bitartrate, Taurine, Choline Chloride, Niacinamide, L-Carnitine, Zinc Sulfate, Potassium Chloride, Salt, Potassium Phosphate, d-Alpha-Tocopheryl Acetate, Calcium Pantothenate, Vitamin A Palmitate, Cupric Sulfate, Riboflavin, Thiamine Chloride Hydrochloride, Pyridoxine Hydrochloride, Folic Acid, Manganese Sulfate, Biotin, Phylloquinone, Sodium Selenate, Vitamin D₃, Oyanocobalamin, and Nucleotides (Cytidine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate). Contains milk and soy ingredients.

NUTRITION INF	01 111111111111111111111111111111111111				
	100 Cal	1000 mL		100 Cal	1000 mL
Energy, Cal	100	812	Biotin, mcg	37	300
Volume, mL	124	1000	Vitamin C, mg	37	300
Protein, g	3.3	26.8	Choline, mg	10	81
% Total Cal	13	13	Inositol, mg	40	325
Source	Nonfat milk, whey protein concentra	ate	Minerals		
Fat, g	5.43	44.07	Calcium, mg	180	1461
% Total Cal	47	47	Calcium, mEq	9.0	72.9
	iglycerides, soy, and coconut oils (0.2	25% DHA; 0.40% ARA)	Phosphorus, mg	100	812
Oil Ratio	50:30:18	50:30:18	Magnesium, mg	12	97.4
Linoleic Acid, mg	700	5681	Iron, mg	1.8	14.6
Carbohydrate, g	10	81	Zinc, mg	1.5	12.17
% Total Cal	40	40	Manganese, mcg	12	97
Source	Corn syrup solids, lactose		Copper, mcg	250	2029
Ratio	50:50	50:50	lodine, mcg	6	49
Vitamins			Selenium, mcg	2	16
Vitamin A, IU	1250	10,144	Sodium, mg	43	349
Vitamin D, IU	150	1217	Sodium, mEq	1.9	15.2
Vitamin E, IU	4.0	32.5	Potassium, mg	129	1047
Vitamin K, mcg	12	97.4	Potassium, mEq	3.3	26.8
Thiamin (Vit B ₁), mcg	250	2029	Chloride, mg	81	657
Riboflavin (Vit B ₂), mcg	620	5032	Chloride, mEq	2.3	18.6
Vitamin B ₆ , mcg	250	2029	Other Characteristics		
Vitamin B ₁₂ , mcg	0.55	4.46	PRSL, mOsm	29.5	240
Niacin, mcg	5000	40,578	Water, g	109	885
Folic Acid, mcg	37	300	Approx Osmolality,	280	280
Pantothenic Acid, mcg	1900	15,419	mOsm/kg H ₂ O		

C. Cohnii Oil is a source of DHA.

M. Alpina Oil is a source of ARA.

Similac® Special Care® 30

Premature Infant Formula with Iron



Description/Indications

A **30 Cal/fl oz**, iron-fortified feeding for growing, low-birth-weight infants and premature infants. First 30 Cal/fl oz ready-to-feed liquid preterm infant formula.

Features

- OptiGRO® is our exclusive blend of DHA, Lutein, and Vitamin E; these important ingredients are found in breast milk
 - DHA for brain and eye development
 - Lutein to support eye health
 - Vitamin E, an important nutrient found in breast milk to support developing cells
- Suitable for use as a human milk fortifier^{1-3*}
- Can be used as a breast milk extender for mothers whose milk supply is low^{1,2*}
- Lutein, a carotenoid naturally found in colostrum and human milk, has been shown to help support eye development of the preterm infant⁴⁻⁷
- Nucleotides for immune support
 - Suitable for use as a human milk fortifed
 - Can be mixed with human milk to a variety of caloric densities (see pages 74-76)
- · Gluten-free; Kosher, Halal

Precautions

- Very low-birth-weight infants are particularly susceptible to gastrointestinal complications; therefore, feeding should be initiated cautiously
- Use this product only after feedings of lower caloric density are well-established. For improved tolerance, it is best to increase caloric density slowly, by 2- to 4-Cal/fl oz increments
- · Hydration status should be monitored
- Spitting up, abdominal distention, abnormal stools or stool
 patterns, excessive gastric residuals, or other signs of intestinal
 dysfunction have been associated with enteral feeding before
 the intestinal tract is ready to accommodate the regimen.
 At the first sign of these problems, enteral feeding should be
 slowed or discontinued
- Not intended for feeding low-birth-weight infants after they reach a weight of 3600 g (approximately 8 lb) or as directed by a physician
- 1. Pillai A, et al. Nutrients 2018;10:1433.
- Willeitner A, et al. JPGN 2017; 65:574.
- 3. Lin Y-C, et al. Nutrients 2020 12:2229.
- Canfield LM, et al. Eur J Nutr 2003;42:133-141.
 Schweigert FJ, et al. Eur J Nutr 2004;43:39-44.
- 6. Patton S, et al. Lipids 1990;25:159-165.
- 7. Rubin LP, et al. J Perinatol 2012;32(6):418-424.

Availability: Hospital/Institutional

Availability: Retail

Size	Container	List No.
Ready To Fe	ed: (30 Cal/fl oz)	
2 fl oz (59 m	L) plastic bottle: 48/case	67448

^{*} Use once feeding tolerance is established.

Preparation



Ready To Feed:

Do not dilute unless directed by a physician.

Can be mixed to a variety of caloric densities using the Liqui-Mix® System (see pages 90-98). Follow physician's instructions.

Ingredients

Ready To Feed: Water, Nonfat Milk, Corn Syrup Solids, Medium Chain Triglycerides, Soy Oil, Whey Protein Concentrate, Coconut Oil, Lactose. Less than 0.5% of: C. Cohnii Oil, M. Alpina Oil, Beta-Carotene, Lutein, Calcium Phosphate, Ascorbic Acid, Calcium Carbonate, Soy Lecithin, Monoglycerides, Magnesium Chloride, Sodium Citrate, m-Inositol, Potassium Hydroxide, Carrageenan, Ferrous Sulfate, Choline Bitartrate, Taurine, Choline Chloride, Niacinamide, L-Carnitine, Zinc Sulfate, d-Alpha-Tocopheryl Acetate, Calcium Pantothenate, Vitamin A Palmitate, Cupric Sulfate, Riboflavin, Potassium Citrate, Pyridoxine Hydrochloride, Thiamine Chloride Hydrochloride, Folic Acid, Manganese Sulfate, Biotin, Phylloquinone, Sodium Selenate, Vitamin D., Cyanocobalamin, Potassium Chloride, Potassium Phosphate, and Nucleotides (Cytidine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate, Adenosine 5'-Monophosphate, Contains milk and soy ingredients.

	100.01			100.0.1	
	100 Cal	1000 mL		100 Cal	1000 mL
Energy, Cal	100	1014	Biotin, mcg	37	375.3
Volume, mL	99	1000	Vitamin C, mg	37	375
Protein, g	3.0	30.4	Choline, mg	10	101
% Total Cal	12	12	Inositol, mg	40	406
Source	Nonfat milk, whey protein concentra		Minerals		
Fat, g	6.61	67.1	Calcium, mg	180	1826
% Total Cal	57	57	Calcium, mEq	9.0	91.3
	triglycerides, soy, and coconut oils (0.2	5% DHA; 0.40% ARA)	Phosphorus, mg	100	1014
Oil Ratio	50:30:18	50:30:18	Magnesium, mg	12	122
Linoleic Acid, mg	700	7101	Iron, mg	1.8	18.3
Carbohydrate, g	7.73	78.4	Zinc, mg	1.5	15.22
% Total Cal	31	31	Manganese, mcg	12	122
Source	Corn syrup solids, lactose		Copper, mcg	250	2536
Ratio	50:50	50:50	lodine, mcg	6	61
Vitamins			Selenium, mcg	2	20
Vitamin A, IU	1250	12,681	Sodium, mg	43	436
Vitamin D, IU	150	1522	Sodium, mEq	1.9	19.0
Vitamin E, IU	4.0	40.6	Potassium, mg	129	1308
Vitamin K, mcg	12	122	Potassium, mEq	3.3	33.5
Thiamin (Vit B ₁), mcg	250	2536	Chloride, mg	81	821
Riboflavin (Vit B ₂), mcg	620	6290	Chloride, mEq	2.3	23.2
Vitamin B ₆ , mcg	250	2536	Other Characteristics		
Vitamin B ₁₂ , mcg	0.55	5.58	PRSL, mOsm	27.8	282.3
Niacin, mcg	5000	50,722	Water, g	84	852
Folic Acid, mcg	37	375	Approx Osmolality,	325	325
Pantothenic Acid, mcg	1900	19.274	mOsm/kg H ₃ O		

C. Cohnii Oil is a source of DHA.

M. Alpina Oil is a source of ARA.

Similac° Liqui-Mix° System

Description

Eliminate most powder mixing—the AND and the CDC* recommend that powdered infant formulas not be used in the NICU unless there is no nutritionally appropriate alternative available.^{1,2} The Liqui-Mix System is a commercially sterile, all-liquid solution that helps

deliver a high-calorie, nutrient-dense feeding.

- NICU babies are often immunocompromised and may be at high risk for developing infections
- Powdered formulas are not commercially sterile
- Reconstituted powder infant formulas have potential for microbial growth and mixing errors

Similac® Special Care® liquid formulas eliminate the need for most powder mixing.

Features

All-Liquid

 Liquid formulas eliminate the need for most powder mixing and meet the AND and CDC recommendations to reduce risk of contamination

Versatile

- Easy mixing to create a variety of calorically dense formulas

Simple

- Easy to use 2:1, 1:1, 1:2 system

NOTE: Proper hygiene, handling, and storage are important when preparing infant formulas. Always follow your hospital's policies and procedures regarding safe handling practices when preparing infant feedings to prevent the possibility of contamination.

- Pediatric Nutrition Practice Group. Steele C, Collins E, eds. Infant and Pediatric Feedings: Guidelines for Preparation of Human Milk and Formula in Health Care Facilities. 3rd ed. Chicago, IL: Academy of Nutrition and Dietetics; 2019:12.
- Centers for Disease Control and Prevention. Enterobacter sakazakii infections
 associated with the use of powdered infant formula—Tennessee, 2001. Available at
 http://www.cdc.gov/mmwr/preview/mmwr/tml/mmS114a1.htm. Published April 11,
 2002. Accessed November 27, 2019.

^{*}Academy of Nutrition and Dietetics and Centers for Disease Control and Prevention.



Individualized Care

The Liqui-Mix System offers flexible solutions to customize feedings to meet the unique nutrient needs of preterm infants. All Similac Special Care Premature Infant Formulas can be used in the Liqui-Mix System.

Recipes for various formula concentrations using the Liqui-Mix® System

(See pages 94-98 for nutrient information)



[†] **EXAMPLE:** To make 1 fl oz (approximately 30 mL) of 22-Cal formula, mix approximately 15 mL of SSC 20 and 15 mL of SSC 24. For convenience, use a Volu-Feed® bottle.



EXAMPLE: To make 1 fl oz (approximately 30 mL) of 22-Cal formula, mix approximately 15 mL of SSC 20 and 15 mL of SSC 24 HP. For convenience, use a Volu-Feed bottle.



Recipes for various formula concentrations using the Liqui-Mix® System


(See pages 94-98 for nutrient information)



Measure liquid formulas by volume when mixing, as bottle fill weights can vary Similac Special Care 20 (SSC 20) + Similac Special Care 30 (SSC 30)

	To Prepare:	Mix the Following:
Э	23 Cal/fl oz formula	2 parts SSC 20 + 1 part SSC 30
	25 Cal/fl oz formula	1 part SSC 20 + 1 part SSC 30
	27 Cal/fl oz formula*	1 part SSC 20 + 2 parts SSC 30

* EXAMPLE: To make 1 fl oz (approximately 30 mL) of 27-Cal formula, mix approximately 10 mL of SSC 20 and 20 mL of SSC 30. For convenience, use a Volu-Feed® bottle.




† EXAMPLE: To make 1 fl oz (approximately 30 mL) of 28-Cal formula, mix approximately 10 mL of SSC 24 and 20 mL of SSC 30. For convenience, use a Volu-Feed bottle.



Recipes for various formula concentrations using the Liqui-Mix® System

(See pages 94-98 for nutrient information)



[‡] EXAMPLE: To make 1 fl oz (approximately 30 mL) of 28-Cal formula, mix approximately 10 mL of SSC 24 HP and 20 mL of SSC 30. For convenience, use a Volu-Feed bottle.



Liqui-Mix® System

Similac® Special Care® 20 + Similac® Special Care® 24 Per 100 Cal Per 100 mL **Nutrients** Similac Special Similac Special Similac Special Similac Special Care 20 (SSC 20) + 1 part SSC 24 Care 20 (SSC 20) + 1 part SSC 24 Care 24 (SSC 24) Care 24 (SSC 24) Cal/fl oz 20 22 24 20 22 24 Energy, Cal 100 100 100 68 74 81 100 Volume, ml. 148 134 124 100 100 Protein, g 3.00 3.00 3.00 2.03 2.23 2.43 Fat, g 5.43 5.43 5.43 3.67 4.04 4.41 Linoleic Acid, mg 700 700 700 473 521 568 Carbohydrate, q 10.3 10.3 10.3 7.7 8.4 Vitamins Vitamin A, IU 1250 1250 1250 845 930 1014 Vitamin D. IU 150 150 150 101 112 122 Vitamin E. IU 4.0 4.0 4.0 2.7 3.0 3.3 Vitamin K, mcg 12 12 12 9 9.7 8.1 186 Thiamin B., mcg 250 250 250 169 203 Riboflavin B., mca 620 620 620 419 461 503 250 250 250 169 Vitamin B., mcg 186 203 Vitamin B,, mcg 0.55 0.55 0.55 0.37 0.41 0.45 Niacin, mca 5000 5000 5000 3382 3720 4058 Folic Acid, mcg 37 37 37 25 28 30 Pantothenic Acid, mcg 1900 1900 1900 1285 1413 1542 Biotin, mcg 37 37 37 25.0 27.5 30.0 Vitamin C, mg 37 37 37 25 28 30 10 10 10 8 Choline, mg 7 Inositol, ma 40 40 40 27 30 33 Minerals 180 122 134 146 Calcium, mg 180 180 68 Phosphorus, ma 100 100 100 74 81 12.0 12.0 Magnesium, mg 12.0 8.1 8.9 9.7 Iron, mg 1.8 1.8 1.8 1.2 1.3 1.5 Zinc. ma 1.50 1.50 1.50 1.01 1.12 1.22 12 12 12 Manganese, mcg 8 9 10 186 Copper, mcg 250 250 250 169 203 lodine, mca 6 6 6 5 Selenium, mcg 1.4 1.5 1.6 Sodium, mg (mEg) 43 (1.9) 43 (1.9) 43 (1.9) 29 (1.26) 32 (1.39) 35 (1.52) Potassium, mg (mEg) 129 (3.3) 129 (3.3) 129 (3.3) 87 (2.23) 96 (2.45) 105 (2.68) Chloride, mg (mEg) 81 (2.3) 81 (2.3) 81 (2.3) 55 (1.55) 60 (1.70) 66 (1.86) Other Characteristics PRSL, mOsm 27.8 27.8 27.8 18.8 20.7 22.6 Water, g 121 88.5 133 109 89.9 89.2 Approx Osmolality, 235 258 280 235 258 280 mOsm/kg water



Liqui-Mix® System
Similac® Special Care® 20 + Similac® Special Care® 24 High Protein (3.3 g per 100 Cal)

WIX" 51	ommac opeci	ai bare 20 + Sillillac	opediar date 24	riigii i roteiii (s.s g		
		Per 100 Cal			Per 100 mL	
Nutrients	Similac Special Care 20 (SSC 20)	1 part SSC 20 + 1 part SSC 24 HP	Similac Special Care 24 High Protein (SSC 24 HP)	Similac Special Care 20 (SSC 20)	1 part SSC 20 + 1 part SSC 24 HP	Similac Special Care 24 High Protein (SSC 24 HP)
Cal/fl oz	20	22	24 20		22	24
Energy, Cal	100	100	100	68 74		81
Volume, mL	148	134	124	100	100	100
Protein, q	3.0	3.16	3.3	2.03	2.35	2.68
Fat, q	5.43	5.43	5,43	3.67	4.04	4.4
Linoleic Acid, mg	700	700	700	473	521	568
Carbohydrate, q	10.3	10.15	10.0	7	7.5	8.1
Vitamins	1212					
Vitamin A, IU	1250	1250	1250	845	930	1014
Vitamin D. IU	150	150	150	101	112	122
Vitamin E. IU	4.0	4.0	4.0	2.7	3.0	3.3
Vitamin K, mcg	12	12	12	8.1	9	9.7
Thiamin B., mcg	250	250	250	169	186	203
Riboflavin B, mcg	620	620	620	419	461	503
Vitamin B _a , mcg	250	250	250	169	186	203
Vitamin B,,, mcq	0.55	0.55	0.55	0.37	0.41	0.45
Niacin, mcq	5000	5000	5000	3382	3720	4058
Folic Acid, mcg	37	37	37	25	28	30
Pantothenic Acid, mcg	1900	1900	1900	1285	1413	1542
Biotin, mcg	37	37	37	25.0	27.5	30.0
Vitamin C. ma	37	37	37	25	28	30
Choline, ma	10	10	10	7	7	8
Inositol, mg	40	40	40	27	30	33
Minerals	.0		.0	2,	55	
Calcium, mg	180	180	180	122	134	146
Phosphorus, mg	100	100	100	68	74	81
Magnesium, mg	12.0	12.0	12.0	8.1	8.9	9.7
Iron, mg	1.8	1.8	1.8	1.2	1.3	1.5
Zinc, mg	1.50	1.50	1.50	1.01	1.12	1.22
Manganese, mcg	12	12	12	8	9	9.7
Copper, mcq	250	250	250	169	186	203
lodine, mcg	6	6	6	4	4	5
Selenium, mcg	2	2	2	1.4	1.5	1.6
Sodium, mg (mEg)	43 (1.9)	43 (1.9)	43 (1.9)	29 (1.26)	32 (1.39)	35 (1.52)
Potassium, mg (mEq)	129 (3.3)	129 (3.3)	129 (3.3)	87 (2.23)	96 (2.45)	105 (2.68)
Chloride, mg (mEq)	81 (2.3)	81 (2.3)	81 (2.3)	55 (1.55)	60 (1.70)	66 (1.86)
Other Characteristics	0. (2.0)	0. (2.0)	0. (2.0)	00 (1.00)	55 (6)	55 (1.50)
PRSL, mOsm	27.8	28.8	29.5	18.8	21.4	24.0
Water, q	133	121	109	89.9	89.2	88.5
Approx Osmolality.	235	258	280	235	258	280
mOsm/kg water	200	250	200	200	200	200
IIIOaiii/ng wdlei		l .				<u> </u>



Liqui-Mix® System Similac® Special Care® 20 + Similac® Special Care® 30

MIX" 51		Simi	iiac° Specia	i Gare° 20 +	Similac [®] S	jeciai Gare [®]	30			
			Per 100 Cal					Per 100 mL	L	
Nutrients	Similac Special Care 20 (SSC 20)	2 parts SSC 20 + 1 part SSC 30	1 part SSC 20 + 1 part SSC 30	1 part SSC 20 + 2 parts SSC 30	Similac Special Care 30 (SSC 30)	Similac Special Care 20 (SSC 20)	2 parts SSC 20 + 1 part SSC 30	1 part SSC 20 + 1 part SSC 30	1 part SSC 20 + 2 parts SSC 30	Similac Specia Care 30 (SSC 3
			a + a	2 + 22			99+9	<u> </u>	E + EE	
Cal/fl oz	20	23	25	27	30	20	23	25	27	30
Energy, Cal	100	100	100	100	100	68	79	85	90	101
Volume, mL	148	127	118	111	99	100	100	100	100	100
Protein, a	3.0	3.0	3.0	3.0	3.0	2.03	2.37	2.54	2.71	3.04
Fat, q	5.43	5,94	6.14	6.32	6.61	3.67	4.68	5.19	5.69	6.71
Linoleic Acid, mg	700	700	700	700	700	473	552	592	631	710
Carbohydrate, q	10.3	9.2	8.8	8.4	7.73	7	7.3	7.4	7.5	7.8
Vitamins										
Vitamin A, IU	1250	1250	1250	1250	1250	845	986	1057	1127	1268
Vitamin D. IU	150	150	150	150	150	101	118	127	135	152
Vitamin E. IU	4.0	4.0	4.0	4.0	4.0	2.7	3.2	3.4	3.6	4.1
Vitamin K, mcg	12	12	12	12	12	8.1	9	10	11	12.2
Thiamin B., mcg	250	250	250	250	250	169	197	211	225	254
Riboflavin B., mcg	620	620	620	620	620	419	489	524	559	629
Vitamin B., mcq	250	250	250	250	250	169	197	211	225	254
Vitamin B ₁₂ , mcg	0.55	0.55	0.55	0.55	0.55	0.37	0.43	0.46	0.50	0.56
Niacin, mcg	5000	5000	5000	5000	5000	3382	3945	4227	4509	5072
Folic Acid, mcg	37	37	37	37	37	25	29	31	33	37.5
Pantothenic Acid, mcg	1900	1900	1900	1900	1900	1285	1499	1606	1713	1927
Biotin, mcq	37	37	37	37	37	25.0	29.2	31.3	33.4	37.5
Vitamin C, mg	37	37	37	37	37	25	29	31	33	38
Choline, mg	10	10	10	10	10	7	8	8	9	10
Inositol, mg	40	40	40	40	40	27	32	34	36	41
Minerals				10			O.E	0.		- ''
Calcium, ma	180	180	180	180	180	122	142	152	162	183
Phosphorus, mg	100	100	100	100	100	68	79	85	90	101
Magnesium, mg	12	12	12	12	12	8.1	9.5	10.1	10.8	12.2
Iron, mg	1.8	1.8	1.8	1.8	1.8	1,2	1.4	1.5	1.6	1.83
Zinc, mg	1.5	1.5	1.5	1.5	1.5	1.01	1.18	1.27	1.35	1.52
Manganese, mcg	12	12	12	12	12	8	9	10	11	12
Copper, mcg	250	250	250	250	250	169	197	211	225	254
lodine, mcg	6	6	6	6	6	4	5	5	5	6
Selenium, mcg	2	2	2	2	2	1.4	1.6	1.7	1.8	2
Sodium, mg (mEg)	43 (1.9)	43 (1.9)	43 (1.9)	43 (1.9)	43 (1.9)	29 (1.26)	34 (1.48)	36 (1.58)	39 (1.69)	44 (1.90)
Potassium, mg (mEg)	129 (3.3)	129 (3.3)	129 (3.3)	129 (3.3)	129 (3.3)	87 (2.23)	102 (2.60)	109 (2.79)	116 (2.97)	131 (3.35)
Chloride, mg (mEg)	81 (2.3)	81 (2.3)	81 (2.3)	81 (2.3)	81 (2.3)	55 (1.55)	64 (1.81)	68 (1.93)	73 (2.06)	82 (2.32)
Other Characteristics		,	, .,	(-/	,		, , ,	,		, ,
PRSL, mOsm	27.8	27.8	27.8	27.8	27.8	18.8	22.0	23.5	25.1	28.2
Water, q	133	112	104	96	84	89.9	88.3	87.6	86.8	85.2
Approx Osmolality,	235	265	280	295	325	235	265	280	295	325
mOsm/kg water										



Liqui-Mix® System Similac Special Care 24 + Similac Special Care 30

		J	iiiiiau opeui	ai vaic 24 T	· Jililiac Jp	scial vale si	,				
			Per 100 Cal			Per 100 mL					
Nutrients	Similac Special Care 24 (SSC 24)	2 parts SSC 24 + 1 part SSC 30	1 part SSC 24 + 1 part SSC 30	1 part SSC 24 + 2 parts SSC 30	Similac Special Care 30 (SSC 30)	Similac Special Care 24 (SSC 24)	2 parts SSC 24 + 1 part SSC 30	1 part SSC 24 + 1 part SSC 30	1 part SSC 24 + 2 parts SSC 30	Similac Specia Care 30 (SSC 30	
Nutrients		B B + B	D + D			Ē	P P + E	P + P	B + BB		
Cal/fl oz	24	26	27	28	30	24	26	27	28	30	
Energy, Cal	100	100	100	100	100	81	88	91	95	101	
Volume, mL	123	114	110	106	99	100	100	100	100	100	
Protein, q	3.0	3.0	3.0	3.0	3.0	2.43	2.64	2.74	2.84	3.04	
Fat. a	5,43	5.88	6.09	6.27	6.61	4.41	5.17	5,56	5.94	6.71	
Linoleic Acid, mg	700	700	700	700	700	568	615	639	663	710	
Carbohydrate, q	10.3	9.3	8.9	8.5	7.73	8.4	8.2	8.1	8.0	7.8	
Vitamins											
Vitamin A. IU	1250	1250	1250	1250	1250	1014	1099	1141	1184	1268	
Vitamin D, IU	150	150	150	150	150	122	132	137	142	152	
Vitamin E. IU	4.0	4.0	4.0	4.0	4.0	3.2	3.5	3.7	3.8	4.1	
Vitamin K. mcg	12	12	12	12	12	9.7	10.6	11.0	11.4	12.2	
Thiamin B., mcg	250	250	250	250	250	203	220	228	237	254	
Riboflavin B ₀ , mcg	620	620	620	620	620	503	545	566	587	629	
Vitamin B _e , mcq	250	250	250	250	250	203	220	228	237	254	
Vitamin B ₁₀ , mcg	0.55	0.55	0.55	0.55	0.55	0.45	0.48	0.50	0.52	0.56	
Niacin, mcg	5000	5000	5000	5000	5000	4058	4396	4565	4734	5072	
Folic Acid, mcg	37	37	37	37	37	30	32.5	33.8	35	37.5	
Pantothenic Acid, mcg	1900	1900	1900	1900	1900	1542	1670	1735	1799	1927	
Biotin, mcg	37	37	37	37	37	30	32.5	33.8	35	37.5	
Vitamin C, mg	37	37	37	37	37	30	33	34	35	38	
Choline, mg	10	10	10	10	10	8.0	9.0	9.0	9.0	10	
Inositol, mg	40	40	40	40	40	32	35	37	38	41	
Minerals	40	40	40	40	40	32	30	31	30	41	
Calcium, mg	180	180	180	180	180	146	158	164	170	183	
Phosphorus, mg	100	100	100	100	100	81	88	91	95	101	
Magnesium, mg	12	12	12	12	12	9.7	10.6	11.0	11.4	12.2	
Iron, mg	1.8	1.8	1.8	1.8	1.8	1.46	1.58	1.64	1.70	1.83	
Zinc. ma	1.5	1.5	1.5	1.5	1.5	1.22	1.32	1.37	1.42	1.52	
Manganese, mcg	12	1.5	1.5	1.5	1.5	1.22	1.32	1.37	1,42	1.52	
Copper, mcg	250	250	250	250	250	203	220	228	237	254	
	6.0	6.0	6.0	6.0	6.0	5.0	5.0	5.0	6.0	6.0	
lodine, mcg Selenium, mcg	2	0.0	2	2	2	1.6	1.7	1.8	1.9	0.0	
Selenium, mcg Sodium, mg (mEg)	43 (1.9)	43 (1.9)	43 (1.9)	43 (1.9)	43 (1.9)	35 (1.5)	38 (1.6)	39 (1.7)	1.9 41 (1.8)	44 (1.9)	
Potassium, mg (mEq)	129 (3.3)	129 (3.3)	129 (3.3)	129 (3.3)	129 (3.3)	105 (2.7)	113 (2.9)	118 (3.0)	122 (3.1)	131 (3.4)	
	129 (3.3) 81 (2.3)		129 (3.3) 81 (2.3)	129 (3.3) 81 (2.3)							
Chloride, mg (mEq) Other Characteristics	61 (2.3)	81 (2.3)	61 (2.3)	81 (2.3)	81 (2.3)	66 (1.9)	71 (2.0)	74 (2.1)	77 (2.2)	82 (2.3)	
PRSL, mOsm	27.0	27.0	97.0	27.0	07.0	22.6	24.5	25.4	26.2	20.2	
	27.8 109	27.8 100.7	27.8 96.5	27.8 92.3	27.8 84	22.6 88.5	24.5 87.4	25.4 86.9	26.3 86.3	28.2 85.2	
Water, g	109 280	100.7	96.5 305	92.3 310	325	88.5 280	87.4 295		86.3 310	325	
Approx Osmolality, mOsm/kg water	280	295	305	310	325	∠80	295	305	310	325	



Liqui-Mix® System
Similac® Special Care® 24 High Protein (3.3 g per 100 Cal) + Similac® Special Care® 30

	Jillillat	opeciai o	are 24 mgm	i Linfelli (9'9	y per rou Ga	i) + Jilillac	Special Gal	IC 30		
			Per 100 Cal			Per 100 mL				
Nutrients	Similac Special Care 24 HP (SSC 24 HP)	2 parts SSC 24 HP + 1 part SSC 30	1 part SSC 24 HP + 1 part SSC 30 +	1 part SSC 24 HP + 2 parts SSC 30 +	Similac Special Care 30 (SSC 30)	Similac Special Care 24 HP (SSC 24 HP)	2 parts SSC 24 HP + 1 part SSC 30 + 1	1 part SSC 24 HP + 1 part SSC 30 +	1 part SSC 24 HP + 2 parts SSC 30 +	Similac Specia Care 30 (SSC 30
Cal/fl oz	24	26	27	28	30	24	26	27	28	30
Energy, Cal	100	100	100	100	100	81	88	91	95	101
Volume, mL	123	114	110	106	99	100	100	100	100	100
Protein. a	3.30	3.18	3.13	3.09	3.0	2.68	2.80	2.86	2.92	3.04
Fat, q	5.43	5.88	6.09	6.27	6.61	4.41	5.17	5.56	5.94	6.71
Linoleic Acid, ma	700	700	700	700	700	568	615	639	663	710
Carbohydrate, q	10.0	9.1	8.7	8.4	7.73	8.1	8.0	8.0	7.9	7.8
Vitamins	. 5.0	5.1	5.7	5.4	0	5.1	5.0	5.0	7.0	7.0
Vitamin A. IU	1250	1250	1250	1250	1250	1014	1099	1141	1184	1268
Vitamin D. IU	150	150	150	150	150	122	132	137	142	152
Vitamin E. IU	4.0	4.0	4.0	4.0	4.0	3.2	3.5	3.7	3.8	4.1
Vitamin K, mcg	12	12	12	12	12	9.7	10.6	11.0	11.4	12.2
Thiamin B., mcg	250	250	250	250	250	203	220	228	237	254
Riboflavin B., mcg	620	620	620	620	620	503	545	566	587	629
Vitamin B, mcg	250	250	250	250	250	203	220	228	237	254
Vitamin B,, mcg	0.55	0.55	0.55	0.55	0.55	0.45	0.48	0.50	0.52	0.56
Niacin, mcg	5000	5000	5000	5000	5000	4058	4396	4565	4734	5072
Folic Acid, mcg	37	37	37	37	37	30	32.5	33.8	35	37.5
Pantothenic Acid, mcg	1900	1900	1900	1900	1900	1542	1670	1735	1799	1927
Biotin, mcg	37	37	37	37	37	30	32.5	33.8	35	37.5
Vitamin C, mg	37	37	37	37	37	30	33	34	35	38
Choline, ma	10	10	10	10	10	8.0	9.0	9.0	9.0	10
Inositol, mg	40	40	40	40	40	32	35	37	38	41
Minerals										
Calcium, mg	180	180	180	180	180	146	158	164	170	183
Phosphorus, mg	100	100	100	100	100	81	88	91	95	101
Magnesium, mg	12	12	12	12	12	9.7	10.6	11.0	11.4	12.2
Iron, mg	1.8	1.8	1.8	1.8	1.8	1.46	1.58	1.64	1.70	1.83
Zinc, mg	1.5	1.5	1.5	1.5	1.5	1.22	1.32	1.37	1.42	1.52
Manganese, mcg	12	12	12	12	12	10	11	11	11	12
Copper, mcg	250	250	250	250	250	203	220	228	237	254
lodine, mcg	6.0	6.0	6.0	6.0	6.0	5.0	5.0	5.0	6.0	6.0
Selenium, mcg	2	2	2	2	2	1.6	1.7	1.8	1.9	2
Sodium, mg (mEq)	43 (1.9)	43 (1.9)	43 (1.9)	43 (1.9)	43 (1.9)	35 (1.5)	38 (1.6)	39 (1.7)	41 (1.8)	44 (1.9)
Potassium, mg (mEq)	129 (3.3)	129 (3.3)	129 (3.3)	129 (3.3)	129 (3.3)	105 (2.7)	113 (2.9)	118 (3.0)	122 (3.1)	131 (3.4)
Chloride, mg (mEq)	81 (2.3)	81 (2.3)	81 (2.3)	81 (2.3)	81 (2.3)	66 (1.9)	71 (2.0)	74 (2.1)	77 (2.2)	82 (2.3)
Other Characteristics										
PRSL, mOsm	29.5	28.9	28.6	28.3	27.8	24.0	25.4	26.1	26.8	28.2
Water, g	109	100.7	96.5	92.3	84	88.5	87.4	86.9	86.3	85.2
Approx Osmolality,	280	295	305	310	325	280	295	305	310	325
mOsm/kg water										

Similac[®] NeoSure[®]

Infant Formula With Iron



Description/Indications

A **22 Cal/fl oz**, nutrient-enriched† formula for babies who were born prematurely. Designed to be used as a preterm post-discharge formula.

Features

- OptiGRO® is our exclusive blend of DHA, Lutein, and Vitamin E; these important ingredients are found in breast milk
- DHA for brain and eye development
- Lutein to support eye health
- Vitamin E, an important nutrient found in breast milk to support developing cells
- Supports excellent catch-up growth during the first year¹
 - Increased protein, vitamins, and minerals compared to term infant formula
 - Increased calories compared to term infant formula for growth
- Calcium and phosphorus for baby's growing bones
- Supports better gains in weight, length, and head circumference when compared to term infant formula¹
- Preterm infants fed Similac NeoSure through the first year of life showed significantly increased lean body mass and less fat mass at 12 months corrected age^{2‡}
- Improved visual acuity at 6 months corrected age^{3‡§}

- Improved early language development at 14 months corrected age^{3‡||}
- Lutein, a carotenoid naturally found in colostrum and human milk, has been shown to help support eye development of the preterm infant⁴⁻⁷
- 25% of the fat blend as medium-chain triglycerides, an easily digested and well-absorbed fat source
- Gluten-free
- · Kosher, Halal
- 1. Carver JD, et al. Pediatrics 2001;107:683-689.
- 2. Groh-Wargo S, et al. Pediatr Res 2005;57:712-718.
- 3. O'Connor DL, et al. Pediatrics 2001;108:359-371.
- 4. Canfield LM, et al. Eur J Nutr 2003;42:133-141.
- 5. Schweigert FJ, et al. Eur J Nutr 2004;43:39-44.
- 6. Patton S, et al. Lipids 1990;25:159-165.
- 7. Rubin LP, et al. J Perinatol 2011; 32(6):418-424.
- * WIC is a service mark of the US Department of Agriculture, and an abbreviation for the Special Supplemental Nutrition Program for Women, Infants, and Children. No endorsement of any brand or product by the USDA is implied or intended. At least one form of Similac® NeoSure® is available through WIC in all 50 states with a valid medical referral.
- † Increased protein, vitamins, and minerals compared to term infant formula.
- ‡ Compared to infants fed a formula without DHA and ARA in a clinical trial with Similac® Special Care® and Similac NeoSure infant formulas with iron; prior to the addition of Lutein.
- § Visual acuity measured at 4 and 6 months corrected age and assessed by VEP (visual evoked potential).
- Il Based on a subset of infants in a post hoc analysis

Availability: Hospital/Institutio	nal List No.
Custom Hospital Feeding System Ready to Feed: (22 Cal/fl oz)	
2 fl oz (59 mL) plastic bottle; 4	8/case 56177
Availability: Retail	
Size Container	List No.
Ready to Feed: (22 Cal/fl oz)	
2 fl oz (59 mL) plastic bottle; 2	4/case 67705
32 fl oz (946 mL) plastic bottle; 6	/case 57455
Powder: (with measuring scoop)	
13.1 oz (371 g); yields 87 fl oz conta	iner; 6/case 57430
22.8 oz (646 g); yields 151 fl oz cont	ainer; 4/case 66923

Ingredients

Ready To Feed: Water, Nonfat Milk, Corn Syrup Solids, Soy Oil, Lactose, Coconut Oil, Whey Protein Concentrate, Medium Chain Triglycerides. Less than 0.5% of: C. Cohnii Oil, M. Alpina Oil, Beta-Carotene, Lutein, Potassium Citrate, Calcium Carbonate, Soy Lecithin, Monoglycerides, Inositol, Ascorbic Acid, Magnesium Chloride, Calcium Phosphate, Carrageenan, Taurine, Choline Chloride, Ferrous Sulfate, Choline Bitartrate, L-Carnitine, Potassium Chloride, d-Alpha-Tocopheryl Acetate, Zinc Sulfate, Niacinamide, Potassium Phosphate, Calcium Pantothenate, Salt, Copper Sulfate, Vitamin A Palmitate, Thiamine Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Manganese Sulfate, Phylloquinone, Biotin, Sodium Selenate, Vitamin D₃, Vitamin B₁₂₂, Potassium Hydroxide and Nucleotides (Adenosine 5'-Monophosphate, Cytidine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate). Contains milk and soy ingredients.

NUTRITION INI	ORMATION				
	100 Cal	1000 mL		100 Cal	1000 mL
Energy, Cal	100	744	Vitamin C, mg	15	112
Volume, mL	134	1000	Choline, mg	16	119
Protein, g	2.8	20.83	Inositol, mg	35	260.0
% Total Cal	11	11	Minerals		
Source	Nonfat milk, whey protein cor	centrate	Calcium, mg	105	781
Fat, g	5.50	40.92	Calcium, mEq	5.2	39.0
% Total Cal	49	49	Phosphorus, mg	62	461
Source Soy oil, co	conut oil, medium chain triglycerides	(0.25% DHA, 0.40% ARA)	Magnesium, mg	9	67
Oil Ratio#	45:29:25	45:29:25	Iron, mg	1.8	13.4
Linoleic Acid, mg	750	5579	Zinc, mg	1.2	8.9
Carbohydrate, g	10.1	75.1	Manganese, mcg	10	74
% Total Cal	40	40	Copper, mcg	120	893
Source	Corn syrup solids, lacto	se	lodine, mcg	15	112
Ratio	50:50	50:50	Selenium, mcg	2.3	17.1
Vitamins			Sodium, mg	33	245
Vitamin A, IU	350	2604	Sodium, mEq	1.4	10.7
Vitamin D, IU	70	521	Potassium, mg	142	1056
Vitamin E, IU	3.6	26.8	Potassium, mEq	3.6	27.0
Vitamin K, mcg	11	81.8	Chloride, mg	75	558
Thiamin (Vit B ₁), mcg	175	1302	Chloride, mEq	2.1	15.7
Riboflavin (Vit B ₂), mcg	150	1116	Other Characteristics		
Vitamin B _e , mcg	100	744	PRSL, mOsm	25.2	187.4
Vitamin B ₁₂ , mcg	0.40	2.98	Water, g	120	893
Niacin, mcg	1950	14,506	Approx Osmolality,	250	250
Folic Acid, mcg	25	186	mOsm/kg H ₂ O		
Pantothenic Acid, mcg	800	5951			
Biotin, mcg	9	67			

[#] Represents 2-fl-oz bottle.

C. Cohnii Oil is a source of DHA.

M. Alpina Oil is a source of ARA.



DIETARY SUPPLEMENT

Similac® Probiotic Tri-Blend Dietary Supplement

Helps Support the Health of the Developing GI Tract



Description/Indications

- Intended for use in infants to support the health of the developing GI tract.*
- Probiotics are not intended for infants with intestinal permeability.

Features

- Administration: Orally or Enterally
- 1 sachet contains 1 billion CFU per packet of an exclusive probiotic blend:
- Bifidobacterium infantis (BB-02[™])
- Streptococcus thermophilus (TH-4®)
- Bifidobacterium lactis (BB-12®)
- Similac Probiotic Tri-Blend is shipped using a cold chain transport system to ensure that the probiotics are maintained at or below 25°C (77°F) from production to delivery
- Potency is guaranteed through date on carton when stored as directed
- Similac Probiotic Tri-Blend is stable at room temperature

Storage

- Store sachets unopened at or below 25°C (77°F)
- Do not warm mixed product beyond 37°C (98.6°F)
- Use by date on the bottom of the carton
- * This statement has not been evaluated by the Food and Drug Administration.

 This product is not intended to diagnose, treat, cure, or prevent any disease.

Availability: Hospital/Institutional

SUPPLEMENT FACTS, POWDER						
	Per 0.5-g packet	% DV†				
Energy, Cal	2					
Carbohydrate, g	0.4	Ť				
Probiotic Blend (1 billion CFU [‡])	88 mg					
Bifidobacterium infantis (BB-02™)						
Streptococcus thermophilus (TH-	-4®)					
Bifidobacterium lactis (BB-12®)						

[†] The % Daily Value (DV) for infants.

[‡] Daily Value not established.

^{1.} Schimmoeller ME, et al. J Neonatal Perinatal Med 2022;15:351-356.

Similac® Probiotic Tri-Blend Mixing Instructions

Please follow your institution's guidelines for mixing protocols

Directions for Preparation and Use

Proper hygiene, handling, and storage are important when preparing Similac Probiotic Tri-Blend.



Pour 3 mL breast milk, donor milk, infant formula, sterile water, or 5% glucose water (commercially sterile) into disposable mixing container¹



Gently shake sachet to ensure contents are settled to bottom



To open sachet, tear top off completely along perforation



Slowly add 1 sachet of Similac Probiotic Tri-Blend powder to mixing container



Using a single-use oral syringe, gently swirl until uniformly distributed



Draw up contents from container into syringe



Administer with oral syringe; mixed product also flows easily through feeding tube



Administer mixed product within 60 minutes

Suggested Use:

1 sachet per day

Preparation:

Start preparation on a clean surface. Similac Probiotic Tri-Blend mixes easily with 3 mL breast milk, donor milk, infant formula, sterile water, or 5% glucose water (commercially sterile).

Storage:

Store sachets unopened at or below 25°C (77°F). Use immediately once sachet is opened.

Warning:

Do not warm mixed product beyond 37°C (98.6°F).

Similac Probiotic Tri-Blend should be mixed outside of patient areas.

A small amount of sediment is normal. Gently shake to resuspend.

Similac Probiotic Tri-Blend is a dietary supplement and should be mixed with a small volume as directed (3 mL) and given separately rather than mixed with a full feeding or fortified feeding to ensure the entire dose is received.

USE BY DATE ON BOTTOM OF CARTON





Custom Feeding System

Description

Abbott provides a comprehensive line of products and services for optimal nutrition of infants.

- Complete line of formulas and support for feeding choices
- A history of excellence and dedication to nutritional innovations
- Health care professional and parent education

Features

- Complete ready-to-use line of infant formulas and water products—all in 2-fl-oz, lightweight, easy-to-open bottles
 - Twist: Break the tamper-evident seal, which is easy with just a few turns
 - Click: Listen for the clicks to assure safety
 - Open and Pour: Into feeding container
- Complete line of ready-to-feed premature infant formulas including Similac® Special Care® 30, the highest caloric density preterm formula and suitable for use as a human milk fortifier
- Liqui-Mix® Preparation Similac Special Care products mix easily using the Liqui-Mix system for the delivery of high-calorie, nutrient-dense formulas
 - All-Liquid: Liquid feedings eliminate the need for most powder mixing and meet the AND and CDC* recommendations to reduce risk of contamination^{1,2}
 - Versatile: Easy mixing to create a variety of calorically dense feedings
 - Simple: Easy to use 2:1, 1:1, 1:2 system
- Human milk fortifiers
 - Similac® Human Milk Fortifier Concentrated Liquid
 - Similac® Human Milk Fortifier Hydrolyzed Protein Concentrated Liquid
 - Liquid Protein Fortifier
- Complete line of nipples designed for versatility and reliability; home delivery available
- Volu-Feed® Volumetric Feeding System—for accurate measurement of formula intake

- Pediatric Nutrition Practice Group, Steele C, Collins E, eds. Infant and Pediatric Feedings: Guidelines for Preparation of Human Milk and Formula in Health Care Facilities. 3rd ed. Chicago, IL: Academy of Nutrition and Dietetics; 2019:12.
- Centers for Disease Control and Prevention. Enterobacter sakazakii infections
 associated with the use of powdered infant formula—Tennessee, 2001. Available at
 http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5114a1.htm. Published April 11,
 2002. Accessed November 27, 2019.

Availability

See individual product listings.

^{*} Academy of Nutrition and Dietetics and Centers for Disease Control and Prevention.









Similac® Special Care® 20	(20 Cal/fl oz)	48/Case	#56265
Similac® Special Care® 24	(24 Cal/fl oz)	48/Case	#56267
Similac® Special Care® 24 High Protein	(24 Cal/fl oz)	48/Case	#56271
Similac [®] Special Care [®] 30*	(30 Cal/fl oz)	48/Case	#56275
Similac® NeoSure®	(22 Cal/fl oz)	48/Case	#56177
Similac® Probiotic Tri-Blend	3/50-sachets/carton	150/Case	#67397



Similac® Human Milk Fortifier Hydrolyzed Protein Concentrated Liquid	(6/24-pkt ctn)	144/Case	#63010
Liquid Protein Fortifier	1.8 fl oz (54 mL)/bottle	48/Case	#62317
Similac® Human Milk Fortifier Concentrated Liquid	(6/24-pkt ctn)	144/Case	#56649

" Can also be used as a human milk fortifier

 $\dagger 301\,\text{NICU professionals} \text{ and healthcare providers who spend most of their time in the hospital; } 90\% confidence level.$

TERM



Similac [®] 360 Total Care [®]	(20 Cal/fl oz)	48/Case	#68136
Similac® Organic	(20 Cal/fl oz)	48/Case	#56525
Similac® With Iron 24	(20 Cal/fl oz)	48/Case	#63075



Similac [®] 360 Total Care [®] Sensitive	(20 Cal/fl oz)	48/Case	#68155
Similac® For Spit-Up	(20 Cal/fl oz)	48/Case	#67389
Similac® Soy Isomil®	(20 Cal/fl oz)	48/Case	#67391
Similac Pro-Total Comfort®	(20 Cal/fl oz)	48/Case	#67132
Similac® Alimentum®	(20 Cal/fl oz)	48/Case	#59738



Similac® Water (Sterlized)	2 fl oz	48/Case	#51000
Similac® 5% Glucose Water	(6 Cal/fl oz)	48/Case	#51002
Similac® 10% Glucose Water	(12 Cal/fl oz)	48/Case	#51004
Similac® Water (Sterilized)	33.8 fl oz (1.05 qt) 1L	8/Case	#58037
Pedialyte [®] Unflavored	(3 Cal/fl oz)	48/Case	#59892

EXTRA SUPPORT

- BREASTFEEDING



CustomFeed[®]
Breast Milk Storage
60 mL-Bottles
100/case
#64125



Breast Milk Storage 4-fl-oz Plastic Bottles 3/pkg #51950



Breast Milk Storage Bottle Caps 250/Case #54080



8-fl-oz Plastic Bottle 144/Case #00875



Volu-Feed® 60-mL Nurser 100/Case #00180



Volu-Feed® Dispensing Cap 250/Case #00081

- NIPPLES



Similac® Premature Nipple & Ring 250/Case #00094



Similac® Orthodontic Nipple & Ring 250/Case #50512



Similac[®] Slow Flow Nipple & Ring 250/Case #53894



Similac[®] Infant Nipple & Ring 250/Case #00079

FEEDING SUPPORT RESOURCES









SKU 68250
SKU 68264
SKU 63329
SKU 56512
SKU 62773

Pediatric PROCONNECT. com

Stay on top of the latest research and product updates with our 24/7 nutrition resource web portal.





Abbott

For more information and patient support, visit us at abbottnutrition.com or contact your Abbott Nutrition Representative

Similac® With Iron 24

Infant Formula



Description/Indications

A **24 Cal/fl oz**, milk-based, iron-fortified formula for term infants needing an increased caloric-density feeding

Features

- Calcium for strong bones-no palm olein oil
- OptiGRO® is our exclusive blend of DHA, Lutein, and Vitamin E; these important ingredients are found in breast milk:
 - DHA for brain and eye development
 - Lutein to support eye health
 - Vitamin E, an important nutrient found in breast milk to support developing cells
- Nucleotides to help support the immune system
- · Carotenoids like those naturally found in breast milk
- · Gluten-free
- Kosher, Halal

Availability: Hospital/Institutional

Custom Feeding System

Size Container List No.

Ready To Feed: (24 Cal/fl oz)

2 fl oz...... plastic bottle; 48/case 63075

Preparation

Ready To Feed: Do not dilute.

Ready to Feed: Water, Lactose, Nonfat Milk, High Oleic Safflower Oil, Soy Oil, Coconut Oil, Whey Protein Concentrate. Less than 0.5% of: C. Cohnii Oil, M. Alpina Oil, Beta-Carotene, Lutein, Ascorbic Acid, Calcium Carbonate, Potassium Citrate, Monoglycerides, Soy Lecithin, Sodium Citrate, Potassium Chloride, Magnesium Chloride, Calcium Phosphate, Potassium Phosphate, Carrageenan, Choline Chloride, Choline Bitartrate, Ferrous Sulfate, Taurine, Inositol, L-tryptophan, d-Alpha-Tocopheryl Acetate, Zinc Sulfate, Niacinamide, Calcium Pantothenate, L-Carnitine, Vitamin A Palmitate, Thiamine Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Copper Sulfate, Manganese Sulfate, Folic Acid, Phylloquinone, Biotin, Sodium Selenite, Vitamin D₃, Vitamin B₁₂. Magnesium Sulfate, Salt, Potassium Iodide, Potassium Hydroxide, and Nucleotides (Adenosine 5'-Monophosphate, Cytidine 5'-Monophosphate), Disodium Guanosine 5'-Monophosphate).

	100 Cal	1000 mL		100 Cal	1000 mL
Energy, Cal	100	806	Vitamin C, mg	15	121
Volume, mL	124	1000	Choline, mg	16	129
Protein, g	2.03	16.37	Inositol, mg	5.9	47.6
% Total Cal	8	8	Minerals		
Source	Nonfat milk, whey protein concentrate		Calcium, mg	79	637
Fat, g	5.5	44.35	Calcium, mEq	3.95	31.9
% Total Cal	50	50	Phosphorus, mg	42	339
Source	High oleic safflower oil, coconut and soy oils (DHA, ARA)		Magnesium, mg	7	56.5
Oil Ratio	40:30:29		Iron, mg	1.8	14.52
Linoleic Acid, mg	870	7016	Zinc, mg	0.69	5.56
Carbohydrate, g	10.6	85.48	Manganese, mcg	5	40.3
% Total Cal	42	42	Copper, mcg	69	556
Source	Lactose		lodine, mcg	14	113
Vitamins			Selenium, mcg	2	16
Vitamin A, IU	296	2387	Sodium, mg	27	218
Vitamin D, IU	52	419	Sodium, mEq	1.2	9.5
Vitamin E, IU	1.5	12.1	Potassium, mg	117	944
Vitamin K, mcg	12	96.8	Potassium, mEq	3	24.1
Thiamin (Vit B,), mcg	96	774	Chloride, mg	65	524
Riboflavin (Vit B ₂), m	cg 173	1395	Chloride, mEq	1.8	14.8
Vitamin B ₆ , mcg	70	565	Other Characteristics		
Vitamin B ₁₂ , mcg	0.35	2.82	PRSL, mOsm	19	153
Niacin, mcg	670	5403	Water, g	104	839
Folic Acid, mcg	12	96.8	Osmolality, mOsm/kg H ₂ O	380	380
Pantothenic Acid, m	cg 590	4758			
Biotin, mcg	4.4	35.5			

Pedialyte®

Oral Electrolyte Solution



Description/Indications

Therapeutic hydration that replenishes fluid and vital minerals lost during diarrhea and vomiting to help prevent mild to moderate dehydration in infants and children; for maintenance of water and electrolytes following corrective parenteral therapy for diarrhea.

Features

- · Ready to use
- Promotes fluid absorption
- Provides glucose to promote sodium and water absorption
- · Helps replenish zinc lost during diarrhea
- · Low osmolality
- · Unflavored liquid available for infants
- · Kosher, Halal

Dosage

Please refer to page 173 for the dosage administration guide for infants and young children.

Precautions

• No mixing or diluting is necessary or recommended

 Use under medical supervision for the dietary management of mild to moderate dehydration during diarrhea and vomiting

Availability: Hospital/Institutional

Custom Feeding System Ready To Use: (3 Cal/fl oz)

Ingredients

Unflavored: Water, Dextrose. Less than 2% of: Potassium Citrate, Salt, Sodium Citrate, Citric Acid and Zinc Gluconate.

NUTRITION INFORMATION	ON	
	2 fl oz	
Energy, Cal	6	
Dextrose, g	1.5	
Sodium, mEq	2.7	
Potassium, mEq	1.2	
Chloride, mEq	2.1	
Zinc, mg	0.5	
Osmolality, mOsm/kg H ₂ O	250	

Similac® 5% Glucose Water

Ready To Feed



Similac® 10% Glucose Water

Ready To Feed



Description/Indications

- For initial or supplemental feeding as directed by physician
- For oral use only
- · Kosher, Halal

Availability

Custom Feeding System

Size	Container	List No.
2 fl 07	plastic bottle: 12/4-packs.	51002

Ingredients

Water and dextrose solution.

Approximate Analysis (wt/vol)

Cal/fl oz 6	Water 96.6%
Cal/100 mL 20	Dextrose 5.0%

Description/Indications

- For initial or supplemental feeding as directed by physician
- For oral use only
- · Kosher, Halal

Availability

Custom Feeding System

Size	Container	LIST NO.
2 fl oz	plastic bottle; 12/4-p	oacks51004

Ingredients

Water and dextrose solution.

Approximate Analysis (wt/vol)

Cal/fl oz 12	Water 93.7%
Cal/100 mL 40	Dextrose 10.0%

Similac® Water (Sterilized) Ready To Feed

Description/Indications

- As an initial feeding
- Use as directed by physician
- For oral use only
- Kosher, Halal
- Can be used to reconstitute concentrated liquid and powdered formula

Availability

Custom Feeding System

Size	Container	List No.
2 fl oz	plastic bottle; 48/case	51000
1 L	reclosable plastic bottle; 8/case	58037

Nipples









Description/Indications

Provides the flexibility to choose a ready-for-use nipple and ring that match each infant's sucking strength or need.

- Infant (latex-free) for infants with developed sucking strength
- Premature (latex-free) for preterm infants with less developed sucking strength
- Orthodontic (contains latex)—for newborn infants, to stimulate oral exercise while feeding
- Slow Flow (latex-free)—to support the underdeveloped "suck-swallow-breathe" coordination of preterm infants. Provides a more controlled flow rate vs faster-flow nipples for increased physiological stability and a more efficient sucking pattern¹

Features

- Easy to use
- · Nipples and rings individually prepackaged and ready for use
- Easy, effective venting system
- · Facilitates latching and consistent flow
- Disposable-not for reuse

Warning

- DO NOT use as a pacifier, as it presents a choking hazard
- DO NOT reuse, as reuse can cause deterioration or separation of the rubber, presenting a choking hazard
- · Keep out of reach of children

1. Chang YJ, et al. J Nurs Res 2007;15(3):215-223.

Availability: Hospital/Institutional Custom Feeding System

Item	Quantity	List No.
Similac® Infant Nipple & Ring	250/case	00079
Similac® Premature Nipple & Ring	250/case	00094
Similac® Orthodontic Nipple & Ring	250/case	50512
Similac® Slow Flow Nipple & Ring	250/case	53894

Similac® CustomFeed® Breast Milk Storage Bottle

Description/Indications

Individually packaged (polypropylene) plastic bottle and lid for use when storing breast milk.

Features

- · Ready to use
- Reusable
- Freezable
- Dishwasher safe*
- Individually packed

Availability

Instructions for use of the CustomFeed™ Breast Milk Storage Bottle

Bottles are ready to use and do not require sterilization before first use.

First use:

- 1. Wash hands and remove bottle from package.
- 2. Unscrew bottle lid and retain in package for later use.
- Screw bottle to breast pump and express milk, leaving some empty space at the top of the bottle for milk to expand as it freezes.
- 4. When finished, unscrew from pump and screw on lid.
- 5. Label bottle with name, date, and time.
- Filled bottles should be stored immediately in refrigerator or freezer and used as directed by your health care professional.

^{*}Top rack only.

After first use:

- Wash bottles and caps in hot, soapy water and rinse thoroughly; or wash in top rack of dishwasher.
- 2. Your health care professional may recommend that you sterilize bottles before filling them with breast milk.

Helpful tips:

- Rinse bottle and cap following use to prevent milk from sticking to bottle.
- When ready to use frozen breast milk, swirl bottle under warm tap water to thaw.
- Feed thawed breast milk right away or refrigerate; do not refreeze.
- WARNING: Never use a microwave oven to thaw or heat milk.
 Serious burns can result.



Description/Indications

Plastic (polypropylene) bottles designed for use in storing human milk. Standard (40-mm) plastic bottle caps fit on 4-fl-oz human milk storage bottles, the 8-fl-oz plastic bottles, and Volu-Feed® Nurser.

Features

- Bottles are ready for use—do not require sterilization before first use
- · Bottles and caps are food-grade quality
- Dishwasher safe (top rack)
- · Packaging material not made with BPA

Cleanliness is important. Before preparing and filling bottles, wash your hands thoroughly.

Bottles DO NOT require sterilization before first use.
AFTER FIRST USE, follow these directions for preparation and use:

- Wash bottles and caps in hot, soapy water and rinse thoroughly; or wash in top rack of dishwasher.
- Your healthcare professional may recommend that you sterilize bottles before filling them with breast milk.
- Fill bottles with breast milk.
 Cap and refrigerate or freeze them promptly. If bottles are not filled immediately, replace caps until needed.

Rinse bottle and cap following use to prevent milk from sticking to bottle.

Filled bottles should be stored in refrigerator and used as directed by your healthcare professional. Discard after 72 hours. If breast milk is to be frozen, chill and freeze it immediately after filling bottles. Leave at least a quarter of the bottle as air space, because the milk will expand. Label all bottles with name. date, and time. When ready to use frozen breast milk, swirl bottle under warm tap water to thaw. Feed thawed breast milk right away or refrigerate; do not refreeze

WARNING: Never use a microwave oven to thaw or heat milk. Serious burns can result.

Availability

Custom Feeding System Size Container List No. Bottles 4 fl oz plastic bottle; 3 bottles with caps/bag; 48 bags; 144 bottles/case 51950 8 fl oz plastic bottle; 144/case 00875 Bottle Caps 250/case 54080

Volu-Feed®



Description/Indications

Plastic (polypropylene) bottle for use when accurate measurement of feeding intake is needed (for example, with premature and low-birth-weight infants).

Features

- · Easy to read
 - Even numbers on one side and odd numbers on the other side
- Numbers centered on the graduation marks
- Large, easy-to-read font
- . Easy to write
 - Large writing panel
- Lightweight plastic
 - Packaging material not made with BPA
- \bullet $\mbox{\bf Ready}$ for use in a clean-peel blisterpack
 - Nipple ready

Availability: Hospital/Institutional

Custom Feeding System		
Item	Container	List No.
Volu-Feed Nurser	. 100/case	00180
Volu-Feed Dispensing Caps .	. 250/case	00081





EleCare® Jr

Amino Acid-Based Nutrition Powder (For Ages 1 and Up)



Description/Indications

A **30 Cal/fl oz**, nutritionally complete amino acid-based nutrition powder for children age 1 and older who cannot tolerate intact or hydrolyzed protein. EleCare Jr is for the dietary management of protein maldigestion, malabsorption, food allergies, short-bowel syndrome, eosinophilic Gl disorders, Gl tract impairment, or other conditions in which an amino acid-based diet is required.

- For ages 1 year and up
- For oral or tube feeding

Features

- Hypoallergenic^{1,*}—virtually eliminating the potential for an allergic reaction to the formula in multiple-foodallergic children
- Shown to be effective in maintaining growth of children with cow's milk protein allergy when used as a primary source of calories^{1,*}
- 100% free amino acids as nitrogen source
- 33% of fat blend as medium-chain triglycerides, an easily digested and well-absorbed fat source
- Supported by strict manufacturing standards and ELISA (Enzyme-linked ImmunoSorbent Assay) testing

- Does not contain milk protein, soy protein, fructose, galactose, lactose, or gluten
- Halal
- Kosher—with pareve ingredients
- Sicherer SH. et al. J Pediatr. 2001;138:688-693.

Availability

Powder:

14.1-oz (400-g) can (measuring scoop enclosed); yields 62 fl oz; 6/case

Flavor	List No.
Unflavored	55253
Vanilla	56585
Chocolate	66273
Banana	66275

^{*} Study conducted with a previous formulation of EleCare unflavored without DHA/ARA.

Unflavored Powder: Corn Syrup Solids (55%), High Oleic Safflower Oil (9%), Medium-Chain Triglycerides (8%), Soy Oil (7%), L-Glutamine (2%). Less than 2% of: L-Asparagine, L-Leucine, DATEM, L-Lysine Acetate, L-Valine, Calcium Phosphate, L-Isoleucine, Potassium Phosphate, L-Arginine, L-Phenylalanine, L-Tyrosine, Potassium Citrate, Sodium Citrate, Sodium Citrate, Sodium Citrate, L-Triveonine, L-Proline, L-Serine, L-Alanine, Magnesium Chloride, Glycine, L-Histidine, L-Methionine, Ascorbic Acid, Calcium Carbonate, L-Cystine Dihydrochloride, L-Tryptophan, Magnesium Phosphate, Choline Chloride, Inositol, Ferrous Sulfate, Taurine, Ascorbyl Palmitate, Zinc Sulfate, dl-Alpha-Tocopheryl Acetate, L-Carnitine, Niacinamide, Salt, Calcium Pantotthenate, Manganese Sulfate, Thiamine Hydrochloride, Copper Sulfate, Vitamin A Palmitate, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Beta-Carotene, Biotin, Phylloquinone, Chromium Chloride, Potassium Iodide, Sodium Selenate, Sodium Molybdate, Vitamin D₃, Vitamin B₁₂.

	100 Cal	100 g powder	1000 mL [†]		100 Cal	100 g powder	1000 mL [†]
Energy, Cal	100	469	1014	Pantothenic Acid, mg	421	1974	4270
Protein Equivalent, g	3.1	14.3	31	Inositol, mg	5	23.6	51
% Total Cal	15 [‡]	15 [‡]	15 [‡]	Minerals			
Source		Free L-amino acids		Calcium, mg	116	543	1174
Fat, g	4.8	22.7	49.1	Calcium, mEq	5.8	27.2	58.7
% Total Cal	43	43	43	Phosphorus, mg	84	395	854
MCT:LCT	33:67	33:67	33:67	Magnesium, mg	16	74	160
Linoleic acid, mg	840	3939	8520	Iron, mg	1.8	8.3	18
Carbohydrate, g	10.5	49.3	106.7	Zinc, mg	1.15	5.4	11.7
% Total Cal	42	42	42	Manganese, mcg	128.2	601	1300
Source		Corn syrup solids		Copper, mcg	128.2	601	1300
Vitamins				lodine, mcg	9	41.6	90
Vitamin A, IU	273	1200	2769	Selenium, mcg	2.7	12.5	27
Vitamin A, mcg	82	384	830	Chromium, mcg	2.3	10.9	23.5
Vitamin D, IU	60	281	608	Molybdenum, mcg	2.6	12.3	26.5
Vitamin D, mcg	1.5	7.0	15.2	Sodium, mg	45	212	459
Vitamin E, mg	2.1	9.71	21	Sodium, mEq	2	9.2	20
Vitamin K, mcg	13	60	130	Sodium, mEq	2	9.2	20
Vitamin C, mg	9	42.5	92	Potassium, mEq	3.8	18	39
Folic Acid, mcg	29.6	139	300	Chloride, mg	60	281	608
Thiamin (Vit B₁), mcg	210	985	2130	Chloride, mEq	1.7	7.9	17
Riboflavin (Vit B ₂), mcg	106	495	1070	Other Characteristics			
Vitamin B ₆ , mcg	84	393	850	PRSL, mOsm	28	129.7	280
Vitamin B ₁₂ , mcg	0.4	2.0	4.3	Water, g	83		839
Niacin, mcg	1680	7878	17,040	Osmolality, mOsm/kg H ₂ O		(Unflavored &	Vanilla) 590
Choline, mg	29.9	140	303				
Biotin, mcg	4.2	19.9	43				

Percentages refer to the weight of the ingredient per total product weight. Vanilla, Chocolate, and Banana contain Artificial Flavor, Aspartame, Acesulfame K, and Sucralose.

[†] Prepared at 30 Cal/fl oz.

[‡] May ME, et al. Am J Clin Nutr. 1990;52:770-776.

PediaSure® Grow & Gain

Complete, Balanced Nutrition®

WIC®-eligible in nearly all 50 states*



Description/Indications

PediaSure Grow & Gain is a source of complete, balanced nutrition especially designed for children 2-13 years of age.

- May be used as the sole source of nutrition or as a supplement
- · Formulated for oral feeding; may also be tube fed

Features

- Clinically proven[†] to help kids grow¹⁻⁶
- Provides 100% or more of the Dietary Reference Intakes (DRIs) for protein and 25 vitamins and minerals in a complete feeding:
 In 1000 mL for children 2 to 8 years of age
 - In 1500 mL for children 9 to 13 years of age
- Non-GMO^{II} and no artificial growth hormones[‡]
- #1 pediatrician recommended brand
- DHA omega-3§ to support the brain and eyes^{7,8}
- Excellent source of calcium, vitamin D, and phosphorus for strong bones
- 7 Nutrients to support immune system health: protein, vitamins A & D, zinc, and antioxidants (vitamins C & E and selenium)
- Kosher, Halal, gluten-free, suitable for lactose intolerance

Precautions

- Consult healthcare professional for use with children under 2 years of age.
- · Not for children with galactosemia
- 1. Akram DS, et al. J Pak Med Assoc. 2000;50:377-380.
- 2. Alarcon PA, et al. Clin Pediatr. 2003;42:209-217.
- 3. Fisberg M, et al. Int Pediatr. 2002;17:216-222.
- Morales E, et al. J Am Diet Assoc. 1991;91:1233-1238.
- Ramstack M. et al. JPEN. 1991:15:89-92.
- 6. Huvnh. et al. *J Am Hum Nutr Diet.* 2015:28:623-635.
- 7. Lauritzen L, et al. Prog Lipid Res. 2001;40:1-94.
- 8. Krinsky NI, et al. Annu Rev Nutr. 2003; 23:171-201.
- * Excludes Arkansas. The WIC (Special Supplemental Nutrition Program for Women, Infants, and Children) name is a service mark of the US Department of Agriculture. USDA does not endorse any goods, services, or enterprises. Requires WIC state medical documentation form. WIC state approval is subject to change, check with your WIC agency on availability.
- † Studied in children at risk for malnutrition, 2 servings per day.
- $^{\ddagger}\text{No}$ significant difference has been shown between milk derived from rbST-treated and non-rbST-treated cows.

Hosnital/

 \S 32 mg of DHA per 8-fl-oz serving.

Hosnital/

	Inst	Inst 8-fl-oz reclosable bottle; 24/case	8-fl-oz bottle;	
Ready To Fee	ed			
Flavor	List No.	List No.	List No.	List No.
Vanilla	67522	67533	58049	62082
Chocolate	67523	67535	58058	–
Strawberry	67525	67537	58055	–
Banana	67527	–	58052	–
Berry	–	–	53818	–
S'mores	–	–	66918	–

Ingredients not genetically engineered.

Vanilla Ready to Feed: Water, Corn Maltodextrin, Blend Of Vegetable Oils (Soy, High Oleic Safflower), Sugar, Milk Protein Concentrate, Soy Protein Isolate; Less than 0.5% of: Vitamins and Minerals*, Natural and Artificial Flavors, Cellulose Gel, Tuna Oil, Cellulose Gum, Monoglycerides, Soy Lecithin, Carrageenan, Potassium Hydroxide, Inositol, Salt, Taurine, Stevia Leaf Extract, L-carnitine, and Lutein. *Vitamins And Minerals*. Potassium Citrate, Calcium Phosphate, Potassium Citrate, Calcium Phosphate, Potassium Citrate, Calcium Phosphate, Potassium Phosphate, Potassium Phosphate, Calcium Carbonate, Ascorbic Acid, Choline Chloride, Ferrous Sulfate, di-Alpha-Tocopheryl Acetate, Zinc Sulfate, Niacinamide, Calcium Pantothenate, Manganese Sulfate, Thiamine Hydrochloride, Pyridoxine Hydrochloride, Riboflavin, Vitamin A Palmitate, Copper Sulfate, Folic Acid, Chromium Chloride, Potassium Iodide, Sodium Molybdate, Sodium Selenate, Biotin, Phylloquinone, Vitamin B., Menaquinone-7. Contains milk and soy ingredients.

	0.0								
	8 fl oz	1L	1.5L	%DV*		8 fl oz	1L	1.5L	%DV [*]
Energy, Cal	240	1000	1500		Riboflavin (Vit B ₂), mg	0.33	1.4	2.1	25
Protein, g	7	30	44	14	Vitamin B _s , mg	0.34	1.4	2.1	20
% Total Cal	12	12	12		Vitamin B ₁₂ , mcg	0.48	2	3	20
Source			y Protein Isolate, No	onfat Milk	Niacin, mg NE	3.2	14	20	20
L-Carnitine, mg	4	17	25		Choline, mg	80	338	506	15
Taurine, mg	18	76	114		Biotin, mcg	8	34	50	25
Fat, g	9	38	57	12	Pantothenic Acid, mg	1.3	5.5	8.2	25
Saturated Fat, g	1.5	6	10	8	Inositol, mg	20	84	126	
Trans Fat, g	0	0	0		Minerals				
Polyunsaturated Fat, g	3.5	14.8	22.2		Calcium, mg	330	1390	2080	25
Monounsaturated Fat, g	3.5	14.8	22.2		Calcium, mEq	16.5	69.5	104	
% Total Cal	34	34	34		Phosphorus, mg	250	1050	1580	20
Source		Soy Oil, High O	leic Safflower Oil		Magnesium, mg	40	169	253	10
Oil Ratio		60:40			Iron, mg	2.7	11	17	15
Cholesterol, mg	5	21	31.6	2	Zinc, mg	1.7	7.2	10.8	15
Carbohydrate, g	33	139	209	12	Manganese, mg	0.46	1.9	2.9	20
% Total Cal	54	54	54		Copper, mg	0.14	0.59	0.89	15
Source		Corn Maltodext	rin, Sugar, Lactose		lodine, mcg	23	97	145	15
Ratio			5:24		Selenium, mcg	8	34	51	15
Total Sugars,† g	9	38	57		Chromium, mcg	9	38	57	25
Added Sugars,† g	9	38	56	18	Molybdenum, mcg	9	38	57	20
Fiber, q	<1	<4	<6	2	Sodium, mg	90	380	570	4
Vitamins					Sodium, mEq	3.9	16.5	24.7	
Vitamin A, mcg RAE	140	590	886	15	Potassium, mg	470	1980	2970	10
Vitamin D, mcg	6	25	38	30	Potassium, mEq	12.1	51	76	
Vitamin E, mg	3	13	19	20	Chloride, mg	230	970	1450	10
Vitamin K, mcg	18	76	114	15	Chloride, mEq	6.5	27	41	
Vitamin C, mg	23	97	146	25	Other Characteristics				
Folate, mcg DFE	60	253	380	15	PRSL, mOsm/L	297	297	297	
Folic Acid, mcg	36	152	227		Water, q	199	840	1260	
Thiamin (Vit B ₁), mg	0.3	1.3	1.9	25	Osmolality, mOsm/kg H ₂ O	460	460	460	

^{*} Percent Daily Values (%DV) apply to the 8-fl-oz serving and are based on a 2.000 Calorie diet.

[†] Sugar content my vary depending on item number.

PediaSure® Grow & Gain Shake Mix



Description/Indications

PediaSure Grow & Gain Shake Mix provides nutrition for growth; for children 2 to 13 years of age.

• Formulated for oral feeding; may also be tube fed

Features

- Clinically proven* to help kids grow¹⁻⁶
- 6g protein and 23 vitamins and minerals
- Non-GMO[†]
- No artificial growth hormones
- #1 pediatrician recommended brand
- · Kosher, Halal, gluten-free

Precautions

- · Use according to package directions
- Consult healthcare professional for use with children under 2 years of age
- Not intended for sole-source of nutrition; may be used as a supplement
- · Not for children with galactosemia

Preparation and Use

• Mix 5 scoops of powder with 3/4 cup water for 1 serving

Availability: Retail

14.1-oz (400-g) can; 6/case

Flavor	List No.
Vanilla	66959
Chocolate	66960
Strawberry	66961

- 1. Akram DS, et al. J Pak Med Assoc. 2000;50:377-380.
- 2. Alarcon PA, et al. Clin Pediatr. 2003;42:209-217.
- 3. Fisberg M, et al. Int Pediatr. 2002;17:216-222.
- 4. Morales E, et al. J Am Diet Assoc. 1991;91:1233-1238.
- 5. Ramstack M, et al. JPEN. 1991;15:89-92.
- 6. Huynh, et al. J Am Hum Nutr Diet. 2015;28:623-635.

^{*} Studied in children at risk for malnutrition, 2 servings per day.

[†] Ingredients not genetically engineered.

[‡] No significant difference has been shown between milk derived from rbST-treated and non-rbST-treated cows.

Vanilla Powder: Corn Syrup, Blend Of Vegetable Oils (High Oleic Sunflower, Canola, Corn), Corn Maltodextrin, Sugar, Milk Protein Concentrate, Soy Protein Isolate, Short-Chain Fructooligosaccharides; Less than 2% of: Vitamins and Minerals! Medium Chain Triglycerides, Natural and Artificial Flavors, Salt, Soy Lecithin, Taurine, Inositol, C. Cohnii Oil, and L-Carnitine Tartrate. *Vitamins And Minerals: Potassium Polaritate, Potassium Citrate, Calcium Phosphate, Potassium Citrate, Calcium Phosphate, Potassium Citrate, Sodium Citrate, Calcium Phosphate, Potassium Citrate, Calcium Phosphate, Choline Chloride, Potassium Hydroxide, Ascorbyl Palmitate, Ascorbic Acid, Ferrous Sulfate, Mixed Tocopherols, Zinc Sulfate, dl-Alpha-Tocopheryl Acetate, Niacinamide, Calcium Pantothenate, Thiamine Hydrochloride, Manganese Sulfate, Pyridoxine Hydrochloride, Riboflavin, Copper Sulfate, Vitamin A Palmitate, Folic Acid, Potassium Iodide, Chromium Chloride, Sodium Selenite, Phylloquinone, Sodium Molybdate, Biotin, Vitamin D., Menaquinone-7, Vitamin B., Contains soy and milk ingredients.

NUTRITION INFO	RMATION, VANILLA				
	Amount	(%DV) ⁻		Amount	(%DV) ⁻
Serving Size	5 scoops (49 g)				
Energy, Cal	220		Thiamin (Vit B ₁), mg	0.54	45
Protein, g	6	112	Riboflavin (Vit B ₂), mg	0.38	30
% Total Cal	11		Vitamin B _e , mg	0.47	25
Source	Milk Protein Concentrate, Soy Protein Isola	te	Vitamin B ₁₂ , mcg	0.54	20
L-Carnitine, mg	3		Niacin, mg NE	2.7	15
Taurine, mg	13		Choline, mg	60	10
Fat, g	9	12	Biotin, mcg	4	10
Saturated Fat, g	2	10	Pantothenic Acid, mg	1.3	25
Trans Fat, g	0		Inositol, mg	14	
Polyunsaturated Fat, g	2		Minerals		
Monounsaturated Fat, g	4.5		Calcium, mg	180	15
% Total Cal	37		Calcium, mEq	9	
Source	High Oleic Sunflower, Canola Oil, Corn Oil, Medium Triglycerides	Chain	Phosphorus, mg	150	10
Oil Ratio	45:27:18:10		Magnesium, mg	30	6
Cholesterol, mg	10	3	Iron, mg	2.5	15
Carbohydrate, g	29	11	Zinc, mg	1.2	10
% Total Cal	52		Manganese, mg	0.27	10
Source	Corn Syrup, Sugar, Corn Maltodextrin		Copper, mg	0.13	15
Ratio	50:33:17		lodine, mcg	17	10
Total Sugars, g	9		Selenium, mcg	6	10
Added Sugars, g	8	16	Chromium, mcg	5	15
Fiber, g	<1	3	Molybdenum, mcg	7	15
Prebiotic	Short-Chain Fructooligosaccharides		Sodium, mg	130	6
Vitamins			Sodium, mEq	5.7	
Vitamin A, mcg RAE	110	10	Potassium, mg	280	6
Vitamin D, mcg	3.6	20	Potassium, mEq	7.2	
Vitamin E, mg	1.5	10	Chloride, mg	180	8
Vitamin K, mcg	11	8	Chloride, mEq	5.1	
Vitamin C, mg	18	20	Other Characteristics		
Folate, mcg DFE	75	20	PRSL, mOsm/L	291	
Folic Acid, mcg	45		Osmolality, mOsm/kg H ₂ O	417	

^{*} Percent Daily Values (%DV) are based on a 2,000 Calorie diet.

PediaSure® Grow & Gain with Fiber

Complete, Balanced Nutrition®



Description/Indications

PediaSure Grow & Gain with Fiber is a source of complete, balanced nutrition especially designed for children 2-13 years of age.

- May be used as the sole source of nutrition or as a supplement
- Formulated for oral feeding; may also be tube fed

Features

- Clinically proven[†] to help kids grow¹⁻⁶
- · Good source of fiber to help maintain regularity
- Provides 100% or more of the Dietary Reference Intakes (DRIs) for protein and 25 vitamins and minerals in a complete feeding:
 - In 1000 mL for children 2 to 8 years of age
- In 1500 mL for children 9 to 13 years of age
- Non-GMO¶ and no artificial growth hormones§
- #1 pediatrician recommended brand
- DHA omega-3" to support the brain and eyes^{7,8}
- Excellent source of calcium, vitamin D, and phosphorus for strong bones
- 7 Nutrients to support immune system health: protein, vitamins A & D, zinc, and antioxidants (vitamins C & E and selenium)
- Kosher, Halal, gluten-free, suitable for lactose intolerance

Precautions

- Consult healthcare professional for use with children under 2 years of age.
- Not for children with galactosemia
- Akram DS. et al. J Pak Med Assoc. 2000:50:377-380.
- 2. Alarcon PA, et al. Clin Pediatr. 2003;42:209-217.
- Fisberg M, et al. Int Pediatr. 2002;17:216-222.
- 4. Morales E, et al. J Am Diet Assoc. 1991;91:1233-1238.
- Ramstack M. et al. JPEN. 1991:15:89-92.
- 5. Harristack IVI, et al. JPEIV. 1991;15:69-92.
- Huynh, et al. J Am Hum Nutr Diet. 2015;28:623-635.
- 7. Lauritzen L, et al. Prog Lipid Res. 2001;40:1-94.
- 8. Krinsky NI, et al. Annu Rev Nutr. 2003;23:171-201.
- The WIC (Special Supplemental Nutrition Program for Women, Infants, and Children) name is a service mark of the US Department of Agriculture. USDA does not endorse any goods, services, or enterprises. Requires WIC state medical documentation form. WIC state approval is subject to change, check with your WIC agency on availability.
- † Studied in children at risk for malnutrition, 2 servings per day.
- Contains 9 grams of total fat per serving.
- \S No significant difference has been shown between milk derived from rbST-treated and non-rbST-treated cows.
- 32 mg of DHA per 8-fl-oz serving.
- ¶ Ingredients not genetically engineered.

Availability: Hospital/Institutional

8-fi-oz cans; 24/case	LIST NO
Flavor	
Vanilla	67529
8-fl-oz reclosable bottles; 24/case	
Vanilla	67521

Availability: Retail

8-fl-oz bottles; 6/carton

Flavor

Vanilla	. 58061
Strawberry	. 56368

Ready to Feed Vanilla: Water, Corn Maltodextrin, Sugar, Blend Of Vegetable Oils (Soy, High Oleic Safflower), Milk Protein Concentrate, Short-Chain Fructooligosaccharides, Soy Protein Isolate, Soy Fiber; Less than 0.5% of: Vitamins and Minerals*, Natural and Artificial Flavors, Tuna Oil, Salt, Monoglycerides, Soy Lecithin, Carrageenan, Potassium Hydroxide, Inositol, Taurine, Stevia Leaf Extract, L-Carnitine, and Lutein. *Vitamins And Minerals*: Potassium Citrate, Calcium Phosphate, Potassium Phosphate, Potassium Phosphate, Potassium Phosphate, Calcium Carbonate, Ascorbic Acid, Chionie Chloride, Ferrous Sulfate, dl-Alpha-Tocopheryl Acetate, Zinc Sulfate, Niacinamide, Calcium Pantothenate, Manganese Sulfate, Thiamine Hydrochloride, Pyridoxine Hydrochloride, Riboflavin, Vitamin A Palmitate, Copper Sulfate, Folic Acid, Chromium Chloride, Potassium lodide, Sodium Molybdate, Sodium Selenate, Biotin, Phylloquinone, Vitamin D., Vitamin B., Menaquinone-7. Contains milk and soy ingredients.

	8 fl oz	1L	1.5L	%DV*		8 fl oz	1L	1.5L	%DV
Energy, Cal	240	1000	1500		Riboflavin (Vit B _a), mg	0.33	1.4	2.1	25
Protein, g	7	30	44	14	Vitamin B _e , mg	0.34	1.4	2.1	20
% Total Cal	12	12	12		Vitamin B ₁₂ , mcg	0.48	2	3	20
Source		otein Concentrate, S			Niacin, mg NE	3.2	14	20	20
L-Carnitine, mg	4	17	25		Choline, mg	80	338	506	15
Taurine, mg	18	76	114		Biotin, mcg	8	34	50	25
Fat, g	9	38	57	12	Pantothenic Acid, mg	1.3	5.5	8.2	25
Saturated Fat, q	1.5	6.5	9.5	8	Inositol, mg	20	84	126	
Trans Fat, q	0	0	0		Minerals				
Polyunsaturated Fat, q	3.5	15	22		Calcium, mg	330	1390	2080	25
Monounsaturated Fat, q	3.5	15	22		Calcium, mEg	16.5	69.5	104	
% Total Cal	34	34	34		Phosphorus, mg	250	1050	1580	20
Source	5	Soy Oil, High Oleic S	afflower Oil		Magnesium, mg	40	169	253	10
Oil Ratio		60:40			Iron, mg	2.7	11	17	15
Cholesterol, mg	5	20	30	2	Zinc, mg	1.7	7.2	10.8	15
Carbohydrate, g	33	139	209	12	Manganese, mg	0.46	1.9	2.9	20
% Total Cal	54	54	54		Copper, mg	0.14	0.59	0.89	15
Source Corn Maltodextrin, Sugar				lodine, mcg	23	97	146	15	
Ratio	64:36				Selenium, mcg	8	34	51	15
Total Sugars,† g	12	51	76		Chromium, mcg	9	38	57	25
Added Sugars,† g	12	51	76	24	Molybdenum, mcg	9	38	57	20
Fiber, g	3	13	19	11	Sodium, mg	90	380	570	4
Prebiotic	Sh	ort-Chain Fructoolig	osaccharides		Sodium, mEq	3.9	16.5	24.7	
Vitamins		_			Potassium, mg	470	1980	2970	10
Vitamin A, mcg RAE	140	590	886	15	Potassium, mEq	12.1	51	76	
Vitamin D, mcg	6	25	38	30	Chloride, mg	230	970	1450	10
Vitamin E, mg	3	13	19	20	Chloride, mEq	6.5	27	41	
Vitamin K, mcg	18	76	114	15	Other Characteristics				
Vitamin C, mg	23	97	146	25	PRSL, mOsm	297	297	297	
Folate, mcg DFE	60	253	380	15	Water, g	200	845	1266	
Folic Acid, mcg	36	152	227		Osmolality, mOsm/kg H ₂ O	495	495	495	
Thiamin (Vit B.), mg	0.3	1.3	1.9	25	, , ,				

^{*} Percent Daily Values (%DV) are based on a 2,000 Calorie diet.

[†] Sugar content my vary depending on item number

PediaSure Harvest™

Complete Nutrition Blend Made with Organic Fruits & Vegetables



Description/Indications

PediaSure Harvest is a complete nutrition blend, including real-food organic ingredients from mango, spinach, pumpkin, banana, and carrot juice concentrate.

- May be used as sole source of nutrition or as a supplement
- Formulated for tube feeding including bolus syringe, gravity syringe, and pump-assisted tube feeding; also suitable for oral consumption
- Use under medical supervision

Features

- Provides 6 servings* of combined fruits and vegetables per liter
- Provides 4 servings[†] of organic whole grains per liter
- Provides a good source of plant-based protein from soy and organic rice
- Meets or exceeds 100% of the Dietary Reference Intakes (DRIs) for protein and 25 essential vitamins and minerals for children 1 to 8 years of age in 1000 mL, and for children 9 to 13 years of age in 1500 mL
- Made with no milk-, gluten-, or lactose-containing ingredients
- · No artificial colors or flavors

- Kosher, Halal, Non-GMO[‡]
- From the makers of PediaSure®, the #1 pediatrician recommended brand

Precautions

- Not intended for infants under 1 year of age unless specified by a physician
- KEEP CAP OUT OF REACH OF SMALL CHILDREN—cap may present a choking hazard
- NOT FOR IV USE

Availability: Hospital/Institutional

Size Container List No.

Ready To Drink (1.0 Cal)

8-fl-oz (237.0-mL) Reclosable Carton; 24/case

Flavor List No.

Unflavored 67962

^{* 1} serving = 1/2 cup eq.

^{† 1} serving = 1-oz eq.

[‡] Ingredients not genetically engineered.

Unflavored Ready to Feed: Water, Organic Mango Puree, Organic Spinach Puree, Organic Milled Whole Grain Brown Rice, Organic Pumpkin Puree, Organic Banana Puree, Organic Rice Maltodextrin, Soy Protein Isolate, Soy Oil, High Oleic Safflower Oil, Organic Rice Protein Concentrate, Organic Carrot Juice Concentrate; Less than 0.5% of: Soy Lecithin, L-Lysine, Schizochytrium sp. Oil, Taurine, and L-Carnitine. Vitamins And Minerals: Calcium Phosphate, Potassium Chloride, Potassium Hydroxide, Ascorbic Acid, Choline Chloride, Ferrous Sulfate, dl-Alphatocopheryl Acetate, Zinc Sulfate, Niacinamide, Calcium Pantothenate, Thiamine Hydrochloride, Biotin, Pyridoxine Hydrochloride, Riboflavin, Vitamin B₁₂, Vitamin D₃, Folic Acid, Potassium Iodide, Contains soy ingredients.

	8 fl oz	1000 mL	1500 mL		8 fl oz	1000 mL	1500 mL
Energy, Cal	240	1000	1500	Vitamin B ₁₉ , mg	0.3	1.3	1.9
Protein, g	9	38	57	Vitamin B,, mcg	0.3	1.3	1.9
% Total Cal	15	15	15	Niacin, mg NE	3.2	13.5	20.3
Source	Soy Protein Isolate,	Organic Rice Protein	Concentrate	Choline, mg	80	338	506
L-Carnitine, mg	4	16.9	25.3	Biotin, mcg	5	21.1	31.6
Taurine, mg	15	63	94	Pantothenic Acid, mg	1.3	5.5	8.2
Fat, g	9	38	57	Inositol, mg	20	84	127
Saturated Fat, g	1.5	6.3	9.5	Minerals			
Trans Fat, g	0	0	0	Calcium, mg	330	1392	2089
Polyunsaturated Fat, g	4	16.5	25	Calcium, mEq	16.5	69.6	104
Monounsaturated Fat, g	3.5	15	22	Phosphorus, mg	250	1055	1582
% Total Cal	34	34	34	Magnesium, mg	40	169	253
Source		ligh Oleic Safflower C		Iron, mg	2.7	11.4	17.1
Cholesterol, mg	0	0	0	Zinc, mg	1.8	7.6	11.4
Carbohydrate, g	31	131	196	Manganese, mg	0.5	2.1	3.1
% Total Cal	. 51	51	51	Copper, mg	0.14	0.6	0.9
Source	Organic Whole G	rain Brown Rice, Orga Maltodextrin	anic Rice	lodine, mcg	23	97	146
Total Sugars, g	9	38	57	Selenium, mcg	7	29.5	44.3
Added Sugars, g	2	8	13	Chromium, mcg	7	29.5	44.3
Fiber, g	2	8	13	Molybdenum, mcg	30	127	190
Vitamins				Sodium, mg	135	570	854
Vitamin A, mcg RAE	200	843	1265	Sodium, mEq	5.9	24.8	37.1
Vitamin D, mcg	6	25	38	Potassium, mg	620	2615	3925
Vitamin E, mg	3	12.7	19	Potassium, mEq	14.1	59.5	89.3
Vitamin K, mcg	30	126	189	Chloride, mg	250	1055	1582
Vitamin C, mg	24	101	152	Chloride, mEq	7.1	29.8	44.7
Folate, mcg DFE	70	295	443	Other Characteristics			
Folic Acid, mcg	42	177	265	PRSL, mOsm	85	361	541
Thiamin (Vit B,), mg	0.3	1.3	1.9	Water, g	196	824	1236
Riboflavin (Vit B _a), mg	0.3	1.3	1.9	Osmolality, mOsm/kg H _o O	500	500	500

PediaSure® Reduced Calorie

Complete, Balanced Nutrition® for children with lower calorie needs



Description/Indications

PediaSure Reduced Calorie' has all the nutrition of base PediaSure® Grow & Gain with fewer calories and less fat.

- For oral use for children requiring a lower-calorie supplement
- For tube feeding for children requiring a lower-calorie complete feeding
- Use under medical supervision

Features

- Antioxidants (vitamins C & E and selenium) to support the immune system
- Supplies prebiotic fiber[†] for digestive health
- Kid-approved taste encourages compliance in children
- Kosher, Halal, gluten-free, suitable for lactose intolerance
- From the makers of PediaSure, the #1 pediatrician recommended brand
- Provides 100% or more of the DRIs for protein and 25 vitamins and minerals in a complete feeding for children 1 to 8 years of age in 1000 mL (633 Cal); for 9 to 13 years of age in 1500 mL (949 Cal)

Precautions

- Not intended for infants under 1 year of age unless specified by a physician
- · Not for IV use
- · Not for children with galactosemia

† Contains 5 grams total fat per serving.

NutraFlora and scFOS are not registered trademarks of Abbott Laboratories.

Availability: Hospital/Institutional

Size Container List No.

Ready To Drink (Reduced Calorie)

8-fl-oz bottle; 24/case

^{* 37%} less calories (150 CAL vs 240 CAL) and 40% less fat (5g vs 9g) for PediaSure Reduced Calorie vs PediaSure Grow & Gain per 8-fl-oz serving.

Vanilla Ready to Feed: Water, Sugar, Milk Protein Concentrate, Soy 0il, Short-chain Fructooligosaccharides, Soy Protein Isolate. Less than 0.5% of: Cellulose Gel, Whey Protein Concentrate, Potassium Citrate, Natural & Artificial Flavor, Potassium Chloride, Magnesium Phosphate, Calcium Phosphate, Salt, Potassium Phosphate, Calcium Carbonate, Cellulose Gum, Choline Chloride, Ascorbic Acid, Carrageenan, Soy Lecithin, Monoglycerides, Potassium Hydroxide, m-Inositol, Taurine, Ferrous Sulfate, di-Alpha-Tocopheryl Acetate, L-Carnitine, Zinc Sulfate, Calcium Pantothenate, Niacinamide, Manganese Sulfate, Thiamine Chloride Hydrochloride, Pyridoxine Hydrochloride, Riboflavin, Cupric Sulfate, Vitamin A Palmitate, Folic Acid, Chromium Chloride, Biotin, Potassium Iodide, Sodium Selenate, Sodium Molybdate, Phylloquinone, Vitamin D., Cyanocobalamin. Contains milk and soy ingredients.

	8 fl oz	1000 mL ⁻	1500 mL ⁻		8 fl oz	1000 mL ⁻	1500 mL ⁻
Energy, Cal	150	633	949	Vitamin B ₁₉ , mcg	0.48	2.0	3.0
Protein, g	7	30	44	Niacin, mg NE	3.2	14	20
% Total Cal	19	19	19	Choline, mg	80	338	506
Source Milk Protein	Concentrate, Soy Pro	tein Isolate, Whey Pro	tein Concentrate	Biotin, mcg	8.0	34	50
L-Carnitine, mg	4.0	17	25	Pantothenic acid, mg	1.3	5.5	8.2
Taurine, mg	18	76	114	Inositol, mg	20	84	126
Fat, g	5	21	32	Minerals			
Saturated Fat, g	1	4	6	Sodium, mg	90	380	570
Trans Fat, g	0	0	0	Sodium, mEq	3.9	16.5	24.7
Polyunsaturated Fat, g	3	13	20	Potassium, mg	470	1980	2970
Monounsaturated Fat, g	1	4	6	Potassium, mEq	12.1	51	76
% Total Cal	30	30	30	Chloride, mg	230	970	1450
Source		Soy Oil		Chloride, mEq	6.5	27	41
Cholesterol, mg	<5	20	30	Calcium, mg	330	1390	2080
Carbohydrates, g	21	89	133	Calcium, mEq	16.5	70	104
% Total Cal	51	51	51	Phosphorus, mg	250	1050	1580
Source	5	Sugar, scFOS		Phosphorus, mEq	16.1	68	102
Dietary Fiber, q	3	13	19	Magnesium, mg	40	169	253
Total Sugars, q	17	72	108	Magnesium, mEg	3.3	14	21
Added Sugars, g	17	72	108	lodine, mcg	23	97	145
/itamins				Manganese, mg	0.46	1.9	2.9
Vitamin A, mcg RAE	140	590	886	Copper, mg	0.14	0.59	0.89
Vitamin D, mcg	6.0	25	38	Zinc, mg	1.7	7.2	10.8
Vitamin E (Total), mg	3.0	13	19	Iron, mg	2.7	11	17
Vitamin K, mcg	18	76	114	Selenium, mcg	8.0	34	51
Vitamin C. mg	23	97	146	Chromium, mcg	9.0	38	57
Folate (Total), mcg DFE	60	253	380	Molybdenum, mcg	9.0	38	57
Folic acid, mcq	36	152	227	Other Characteristics			
Thiamin (Vit B.), mg	0.30	1.3	1.9	Water, g	212	895	1342
Riboflavin (Vit B _a), mg	0.33	1.4	2.1	Osmolality, mOsm/kg H _o O	420	420	420
Vitamin B ₆ mg	0.34	1.4	2.1	22	.20	0	.20

^{*} Values extrapolated from 8 fl oz. †Product contains DHA (32 mg DHA/8 fl oz). †scFOS = short-chain fructooligosaccharides.

PediaSure SideKicks®

Balanced Nutrition to Help Fill Gaps



Description/Indications

PediaSure SideKicks is a complete, balanced nutrition shake for kids that is high in protein and can help fill the top 4 nutrient gaps (Vitamin D, Calcium, Fiber, Potassium).

• Formulated for oral feeding; may also be tube fed

Features

- 10 g protein per 8-fl-oz serving
- No artificial growth hormones*
- Suitable for lactose intolerance, gluten-free, Kosher, and Halal
- 7 Nutrients to support immune system health: protein, vitamins A & D, zinc, and antioxidants (vitamins C & E and selenium)

Precautions

- Consult healthcare professional for use with children under 2 years of age.
- · Not for children with galactosemia

Availability: Retail

Ready To Drink

8-fl-oz bottle; 6/carton

Flavor	List No.
Vanilla	66912
Chocolate	66639
Strawberry	66916

^{*} No significant difference has been shown between milk derived from rbST-treated and non-rbST-treated cows.

Vanilla Ready to Feed: Water, Sugar, Milk Protein Concentrate, Corn Maltodextrin, Blend Of Vegetable Oils (Canola, Corn, Soy), Short-Chain Fructooligosaccharides, Soy Protein Isolate; Less than 0.5% of: Vitamins and Minerals!, Nonfat Milk, Cellulose Gel, Natural and Artificial Flavors, Soy Lecithin, Cellulose Gum, Salt, Carrageenan, Inositol, Taurine, Stevia Leaf Extract, and L-carnitine. *Vitamins And Minerals: Potassium Citrate, Potassium Chloride, Magnesium Phosphate, Calcium Phosphate, Calcium Carbonate, Potassium Phosphate, Ascorbic Acid, Choline Chloride, Potassium Hydroxide, Ferrous Sulfate, dl-Alpha-Tocopheryl Acetate, Zinc Sulfate, Niacinamide, Calcium Pantothenate, Manganese Sulfate, Thiamine Hydrochloride, Pyridoxine Hydrochloride, Vitamin A Palmitate, Riboflavin, Copper Sulfate, Folic Acid, Chromium Chloride, Potassium Iodide, Sodium Molybdate, Sodium Selenate, Biotin, Phylloquinone, Vitamin D_{st} and Vitamin B_{st}. Contains milk and soy ingredients.

	Amount	(%DV)		Amount	(%DV)*
Energy, Cal	180	(/051/	Riboflavin (Vit B ₂), mg	0.33	25
Protein, g	10	20	Vitamin B _e , mg	0.34	20
% Total Cal	22	20	Vitamin B ₁₀ , mcg	0.48	20
Source	Milk Protein Concetrate, Soy Protein Isolate, Non	fot Milk	Niacin, mg NE	3.2	20
L-Carnitine, mg	Wilk Protein Concernate, Soy Protein Isolate, Non	ial Will	Choline, mg	80	15
Taurine, mg	18		Biotin, mcg	8	25
Fat, g	6	8	Pantothenic Acid, mg	1.3	25
Saturated Fat, q	1	5	Inositol, mg	20	20
Trans Fat, q	0	3	Minerals	20	
Polyunsaturated Fat, q	2.5		Calcium, mg	330	25
Monounsaturated Fat, g	2.5			16.5	25
% Total Cal	2.5 30		Calcium, mEq	250	20
,	• • • • • • • • • • • • • • • • • • • •		Phosphorus, mg	40	10
Source Oil Ratio	Canola Oil, Corn Oil, Soy Oil 60:20:20		Magnesium, mg	2.7	15
	10	0	Iron, mg		15
Cholesterol, mg		3	Zinc, mg	1.7	
Carbohydrate, g % Total Cal	23 48	8	Manganese, mg	0.46 0.14	20 15
,			Copper, mg		
Source	Sugar, Corn Maltodextrin		lodine, mcg	23	15 15
Ratio	87:13		Selenium, mcg	8	
Total Sugars, g	12	00	Chromium, mcg	9	25
Added Sugars, g	11	22	Molybdenum, mcg	9	20
Fiber, g	3	11	Sodium, mg	90	4
Prebiotic	Short-Chain Fructooligosaccharides		Sodium, mEq	3.9	
Vitamins			Potassium, mg	470	10
Vitamin A, mcg RAE	140	15	Potassium, mEq	12.1	
Vitamin D, mcg	6	30	Chloride, mg	230	10
Vitamin E, mg	3	20	Chloride, mEq	6.5	
Vitamin K, mcg	18	15	Other Characteristics		
Vitamin C, mg	23	25	PRSL, mOsm/L	358	
Folate, mcg DFE	60	15	Water, g	197	
Folic Acid, mcg	36		Osmolality, mOsmL/kg H ₂ O	415	
Thiamin (Vit B ₁), mg	0.3	25			

^{*} Percent Daily Values (%DV) are based on a 2,000 Calorie diet.

PediaSure® Enteral Formula 1.0 Cal

Complete, Balanced Nutrition®



Description/Indications

PediaSure Enteral Formula 1.0 Cal is a source of complete, balanced nutrition especially designed for tube feeding children 1 to 13 years of age.

- May be used as the sole source of nutrition or as a supplement
- · Formulated for tube feeding

Features

- Milk-based
- 1.0 Cal per mL, 240 Cal per 8 fl oz, from a balanced distribution of protein, fat, and carbohydrate
- Meets or exceeds 100% of the DRIs for protein and 25 essential vitamins and minerals in 1000 mL for children ages 1 to 8 and in 1500 mL for children ages 9 to 13
- DHA omega-3* to help support brain and eye development^{1,2}
- Good source of antioxidants[†] to support the immune system
- Gluten-free, suitable for lactose intolerance, low-residue
- · Kosher, Halal

- * 32 mg DHA per 8-fl-oz serving.
- † Vitamins C & E and selenium.
- 1 Lauritzen L, et al. Prog Lipid Res. 2001;40:1-94.
- 2 Krinsky NI, et al. Annu Rev Nutr. 2003;23:171-201.

Precautions

- Not intended for infants under 1 year of age unless specified by a physician
- · Not for children with galactosemia

Availability: Hospital/Institutional

8-fl-oz can; 24/case

Flavor	List No.
Vanilla	67401

Vanilla Ready to Feed: Water, Corn Maltodextrin, Blend Of Vegetable Oils (High Oleic Safflower, Soy, Medium Chain Triglycerides), Milk Protein Concentrate, Sugar, Soy Protein Isolate; Less than 0.5% of: Vitamins And Minerals, * Natural and Artificial Flavor, Cellulose Gel, Tuna Oil, * Soy Lecithin, Monoglycerides, Carrageenan, Salt, Cellulose Gum, Potassium Hydroxide, Inositol, Taurine, L-Carnitine, and Lutein. * Vitamins and Minerals: Potassium Phosphate, Potassium Citrate, Calcium Carbonate, Potassium Chloride, Magnesium Phosphate, Calcium Phosphate, Ascorbic Acid, Choline Chloride, Ferrous Sulfate, Al-Alpha-Tocopheryl Acetate, Zinc Sulfate, Niacinamide, Calcium Pantothenate, Manganese Sulfate, Thiamine Hydrochloride, Pyridoxine Hydrochloride, Riboflavin, Vitamin A Palmitate, Copper Sulfate, Folic Acid, Chromium Chloride, Potassium lodide, Sodium Molybdate, Sodium Selenate, Biotin, Phylloquinone, Vitamin D₃, Vitamin B₁₂: † Source of docosahexaenoic acid (DHA). Contains milk and soy ingredients.

	8 fl oz (237 mL)	1000 mL‡	1500 mL‡		8 fl oz (237 mL)	1000 mL‡	1500 mL‡
Energy, Cal	240	1000	1500	Riboflavin (Vit B _a), mg	0.33	1.4	2.1
Protein, g	7	30	44	Vitamin B _s , mg	0.34	1.4	2.1
% Total Cal	12	12	12	Vitamin B ₁₂ , mcg	0.48	2	3
Source	Milk Protein Con	centrate, Soy Proteir	Isolate	Niacin, mg NE	3.2	14	20
L-Carnitine, mg	4	17	25	Choline, mg	80	338	506
Taurine, mg	18	76	114	Biotin, mcg	8	34	50
Fat, g	9	38	57	Pantothenic Acid, mg	1.3	5.5	8.2
Saturated Fat, g	2.5	10	15	Inositol, mg	20	84	127
Trans Fat, g	0	0	0	Minerals			
Polyunsaturated Fat, g	2.5	11	16	Calcium, mg	330	1390	2080
Monounsaturated Fat, g	3.5	15	22	Calcium, mEq	16.5	70	104
% Total Cal	34	34	34	Phosphorus, mg	250	1050	1580
Source		ver Oil, Soy Oil, Medii Triglycerides	um Chain	Magnesium, mg	40	169	253
Oil Ratio		43:42:15		Iron, mg	2.7	11	17
Cholesterol, mg	5	20	30	Zinc, mg	1.7	7.2	10.8
Carbohydrate, g	33	139	209	Manganese, mg	0.46	1.9	2.9
% Total Cal	54	54	54	Copper, mg	0.14	0.59	0.89
Source	Corn N	Maltodextrin, Sugar		lodine, mcg	23	97	145
Ratio		86:14		Selenium, mcg	8	34	51
Total Sugars, g	7	29	44	Chromium, mcg	9	38	57
Added Sugars, g	7	29	44	Molybdenum, mcg	9	38	57
Fiber, q	<1	<1	1	Sodium, mg	90	380	570
Vitamins				Sodium, mEq	3.9	16.5	24.7
Vitamin A, mcg RAE	140	590	886	Potassium, mg	470	1980	2970
Vitamin D, mcg	6	25	38	Potassium, mEq	12.1	51	76
Vitamin E, mg	3	13	19	Chloride, mg	230	970	1450
Vitamin K, mcg	18	76	114	Chloride, mEg	6.5	27	41
Vitamin C, mg	23	97	146	Other Characteristics			
Folate, mcg DFE	60	253	380	PRSL, mOsm	66	278	417
Folic Acid, mcg	36	152	227	Water, q	200	845	1266
Thiamin (Vit B.), mg	0.3	1.3	1.9	Osmolality, mOsm/kg H _a O	340	340	340

139

[‡]Values extrapolated from 8 fl oz.

PediaSure® Enteral Formula 1.0 Cal with Fiber

Complete, Balanced Nutrition®



PediaSure Enteral Formula 1.0 Cal with Fiber is a source of complete, balanced nutrition especially designed for tube feeding children 1 to 13 years of age.

- May be used as the sole source of nutrition or as a supplement
- · Specially formulated for tube feeding
- Use under medical supervision

Features

- Milk-based
- 1.0 Cal per mL, 240 Cal per 8 fl oz, from a balanced distribution of protein, fat, and carbohydrate
- Meets or exceeds 100% of the DRIs for protein and 25 essential vitamins and minerals in 1000 mL for children ages 1 to 8 and in 1500 mL for children ages 9 to 13
- DHA omega-3* to help support brain and eye development^{4,5}
- Blend of soluble and insoluble fibers and short-chain fructooligosaccharides at a level that is well tolerated by children
- Prebiotics designed to support digestive tract health¹⁻³
- Antioxidants to help support the immune system

- · Gluten-free, suitable for lactose intolerance
- · Kosher, Halal

Precautions

- Not intended for infants under 1 year of age unless specified by a physician
- · Not for children with galactosemia
- 1. Bornet FR, et al. Nutr Rev 2002;60:326-334.
- Tokunaga T, et al. Bifidus 1993;6:143-150.
- 3. Hidaka H, et al. Bifidobact Microflora 1986;5:37-50.
- Lauritzen L, et al. Prog Lipid Res. 2001;40:1-94.
- 5. Krinsky NI, et al. Annu Rev Nutr. 2003;23:171-201.
- * 32 mg DHA per 8-fl-oz serving
- † Vitamins C & E and selenium

NutraFlora and scFOS are not registered trademarks of Abbott Laboratories.

Availability: Hospital/Institutional

Ready To Hang
1000-mL prefilled containers; 8/case
Flavor
List No.

Vanilla Ready to Feed: Water, Corn Maltodextrin, Blend of Vegetable Oils (High Oleic Safflower, Soy, Medium Chain Triglycerides), Milk Protein Concentrate, Sugar, Soy Protein Isolate, Short-Chain Fructooligosaccharides; Less than 0.5% of: Vitamins And Minerals*, Oat Fiber, Soy Fiber, Natural And Artificial Flavor, Gurn Arabic, Cellulose Gel, Tuna Oil*, Cellulose Gurn, Soy Lecithin, Monoglycerides, Carrageenan, Salt, Potassium Hydroxide, Inositol, Taurine, L-Carnitine and Lutein. * Vitamins and Minerals: Potassium Phosphate, Calcium Carbonate, Potassium Chloride, Potassium Citrate, Calcium Phosphate, Ascorbic Acid, Choline Chloride, Ferrous Sulfate, Alpha-Tocopheryl Acetate, Zinc Sulfate, Niacinamide, Calcium Pantothenate, Manganese Sulfate, Thiamine Hydrochloride, Pyridoxine Hydrochloride, Riboflavin, Vitamin A Palmitate, Copper Sulfate, Magnesium Phosphate, Folic Acid, Chromium Chloride, Potassium Iodide, Sodium Molybdate, Sodium Selenate, Biotin, Phylloquinone, Vitamin D., Vitamin B., Contains milk and soy ingredients.

NUTRITION INFO	RMATION						
	8 fl oz (237 mL)	1000 mL‡	1500 mL [‡]		8 fl oz (237 mL)	1000 mL‡	1500 mL‡
Energy, Cal	240	1000	1500	Riboflavin (Vit B ₂), mg	0.33	1.4	2.1
Protein, g	7	30	44	Vitamin B ₆ , mg	0.34	1.4	2.1
% Total Cal	12	12	12	Vitamin B ₁₂ , mcg	0.48	2	3
Source	Milk Protein Cor	centrate, Soy Protein	Isolate	Niacin, mg NE	3.2	14	20
L-Carnitine, mg	4	17	25	Choline, mg	80	338	506
Taurine, mg	18	76	114	Biotin, mcg	8	34	50
Fat, g	9	38	57	Pantothenic Acid, mg	1.3	5.5	8.2
Saturated Fat, g	2.5	10	15	Inositol, mg	20	84	126
Trans Fat, g	0	0	0	Minerals			
Polyunsaturated Fat, g	2.5	11	16	Calcium, mg	330	1390	2080
Monounsaturated Fat, g	3.5	15	22	Calcium, mEq	16.5	70	104
% Total Cal	34	34	34	Phosphorus, mg	250	1050	1580
Source		ver Oil, Soy Oil, Mediu Triglycerides	ım Chain	Magnesium, mg	40	169	253
Oil Ratio		43:42:15		Iron, mg	2.7	11	17
Cholesterol, mg	5	20	30	Zinc, mg	1.7	7.2	10.8
Carbohydrate, g	34	143	214	Manganese, mg	0.46	1.9	2.9
% Total Cal	54	54	54	Copper, mg	0.14	0.59	0.89
Source	Corn I	Maltodextrin, Sugar		lodine, mcg	23	97	145
Ratio		86:14		Selenium, mcg	8	34	51
Total Sugars, g	7	28	42	Chromium, mcg	9	38	57
Added Sugars, q	7	28	42	Molybdenum, mcg	9	38	57
Fiber, q	3	13	19	Sodium, mg	90	380	570
Prebiotic	Short-Chair	Fructooligosacchario	les	Sodium, mEq	3.9	16.5	24.7
Vitamins		·		Potassium, mg	470	1980	2970
Vitamin A, mcg RAE	140	590	886	Potassium, mEq	12.1	51	76
Vitamin D, mcg	6	25	38	Chloride, mg	230	970	1450
Vitamin E, mg	3	13	19	Chloride, mEa	6.5	27	41
Vitamin K, mcg	18	76	114	Other Characteristics			
Vitamin C, mg	23	97	146	PRSL. mOsm	66	278	417
Folate, mcg DFE	60	253	380	Water, q	200	845	1266
Folic Acid, mcq	36	152	227	Osmolality, mOsm/kg H _o O	400	400	400
Thiamin (Vit B.), mg	0.3	1.3	1.9	2 2,	100		100

[‡] Values extrapolated from 8 fl oz. [†] Source of docosahexaenoic acid (DHA).

PediaSure® 1.5 Cal

Complete, Balanced Nutrition®



Description/Indications

PediaSure 1.5 Cal is a higher* caloric density product designed to meet the energy requirements of pediatric patients who are at risk for malnutrition, require a higher caloric density, or have fluid restrictions. PediaSure 1.5 Cal provides a source of complete, balanced nutrition for children 1 to 13 years of age.

- May be used as the sole source of nutrition or as a supplement
- · Formulated for oral or tube feeding
- Use under medical supervision

Features

- Meets or exceeds 100% of the DRIs for protein and 25 essential vitamins and minerals in 1000 mL for children ages 1 to 8 and in 1500 mL for children ages 9 to 13
- Excellent source of high-quality protein designed to meet the acceptable macronutrient distribution range of protein for children ages 1 to 13
- DHA omega-3[†] to help support brain and eye health^{1,2}
- Good source of antioxidants[‡] to support the immune system

- Gluten-free, suitable for lactose intolerance, low-residue
- Kosher, Halal

Precautions

- Not intended for infants under 1 year of age unless specified by a physician
- · Not for children with galactosemia

Availability: Hospital/Institutional

8-fl-oz can: 24/case

Flavor	List No.
Vanilla	67378

^{*} Caloric value is 1.5 Cal/mL compared to base PediaSure® at 1.0 Cal/mL.

^{† 32} mg of DHA per 8-fl-oz serving.

[‡] Vitamins C & E and selenium.

^{1.} Lauritzen L, et al. Prog Lipid Res. 2001;40:1-94.

^{2.} Krinsky NI, et al. Annu Rev Nutr. 2003;23:171-201.

Vanilla Ready to Feed: Water, Corn Maltodextrin, Blend Of Vegetable Oils (High Oleic Safflower, Soy, Medium Chain Triglycerides), Milk Protein Concentrate, Soy Protein Isolate; Less than 0.5% of: Vitamins And Minerals*, Natural And Artificial Flavor, Cellulose Gel, Soy Lecithin, Tuna Oil*, Monoglycerides, Cellulose Gum, Potassium Hydroxide, Carrageenan, Inositol, Taurine, Sucralose, Salt, L-Carnitine and Lutein. *Vitamins and Minerals*: Potassium Citrate, Potassium Chloride, Magnesium Phosphate, Calcium Phosphate, Ascorbic Acid, Choline Chloride, Calcium Carbonate, Potassium Phosphate, Ferrous Sulfate, di-Alpha-Tocopheryl Acetate, Zinc Sulfate, Niacinamide, Calcium Pantothenate, Manganese Sulfate, Thiamine Hydrochloride, Pyridoxine Hydrochloride, Riboflavin, Vitamin A Palmitate, Copper Sulfate, Folic Acid, Chromium Chloride, Potassium Iodide, Sodium Molybdate, Sodium Selenate, Biotin, Phylloquinone, Vitamin B., Contains milk and soy ingredients.

	8 fl oz (237 mL)	1000 mL‡	1500 mL‡		8 fl oz (237 mL)	1000 mL‡	1500 mL‡
Energy, Cal	350	1500	2250	Vitamin B _e , mg	0.34	1.4	2.1
Protein, g	14	59	89	Vitamin B,, mcg	0.48	2	3
% Total Cal	16	16	16	Niacin, mg NE	3.2	14	20
Source	Milk Protein Cor	centrate, Soy Protein	Isolate	Choline, mg	80	338	506
L-Carnitine, mg	4	17	25	Biotin, mcg	8	34	50
Taurine, mg	18	76	114	Pantothenic Acid, mg	1.3	5.5	8.2
Fat, g	16	67	100	Inositol, mg	20	84	126
Saturated Fat, g	3.5	14	22	Minerals			
Trans Fat, g	0	0	0	Calcium, mg	330	1390	2080
Polyunsaturated Fat, g	5	20	30	Calcium, mEq	16.5	70	104
Monounsaturated Fat, g	7	30	44	Phosphorus, mg	250	1050	1580
% Total Cal	41	41	41	Magnesium, mg	40	169	253
Source		ver Oil, Soy Oil, Mediu Triglycerides	m Chain	Iron, mg	2.7	11	17
Oil Ratio		50:40:10		Zinc, mg	1.7	7.2	10.8
Cholesterol, mg	10	40	63	Manganese, mg	0.46	1.9	2.9
Carbohydrate, g	38	160	241	Copper, mg	0.14	0.59	0.89
% Total Cal	43	43	43	lodine, mcg	23	97	145
Source	Co	rn Maltodextrin		Selenium, mcg	8	34	51
Total Sugars, g	3	11	17	Chromium, mcg	9	38	57
Added Sugars, g	3	10	15	Molybdenum, mcg	9	38	57
Fiber, g	<1	1	2	Sodium, mg	90	380	570
Vitamins				Sodium, mEq	3.9	16.5	24.7
Vitamin A, mcg RAE	140	590	886	Potassium, mg	470	1980	2970
Vitamin D, mcg	6	25	38	Potassium, mEq	12.1	51	76
Vitamin E, mg	3	13	19	Chloride, mg	230	970	1450
Vitamin K, mcg	18	76	114	Chloride, mEq	6.5	27	41
Vitamin C, mg	23	97	146	Other Characteristics			
Folate, mcg DFE	60	253	380	PRSL, mOsm	110	462	694
Folic Acid, mcg	36	152	227	Water, g	185	780	1170
Thiamin (Vit B ₁), mg	0.3	1.3	1.9	Osmolality, mOsm/kg H ₂ O	400	400	400
Riboflavin (Vit B _a), mg	0.33	1.4	2.1				

[‡] Source of docosahexaenoic acid (DHA). ¶ Values extrapolated from 8 fl oz.

PediaSure® 1.5 Cal with Fiber

Complete, Balanced Nutrition®



Description/Indications

PediaSure 1.5 Cal with Fiber is a higher* caloric density product designed to meet the energy requirements of pediatric patients who are at risk for malnutrition, require a higher caloric density, or have fluid restrictions. PediaSure 1.5 Cal with Fiber provides a source of complete, balanced nutrition for children 1 to 13 years of age.

- May be used as the sole source of nutrition or as a supplement
- Formulated for oral or tube feeding
- Use under medical supervision

Features

- Meets or exceeds 100% of the DRIs for protein and 25 essential vitamins and minerals in 1000 mL for children ages 1 to 8 and in 1500 mL for children ages 9 to 13
- Excellent source of high-quality protein designed to meet the acceptable macronutrient distribution range of protein for children ages 1 to 13
- Prebiotics designed to support digestive tract health¹⁻³
- Antioxidants[†] to help support the immune system
- DHA omega-3[‡] to support brain and eye health^{4,5}

- Gluten-free, suitable for lactose intolerance
- Kosher, Halal

Precautions

- Not intended for infants under 1 year of age unless specified by a physician
- · Not for children with galactosemia
- Bornet FR. et al. Nutr Rev 2002:60:326-334.
- 2. Tokunaga T, et al. Bifidus 1993;6:143-150.
- 3. Hidaka H. et al. Bifidobact Microflora 1986:5:37-50.
- Lauritzen L, et al. Prog Lipid Res. 2001;40:1-94.
- 5. Krinsky NI, et al. Annu Rev Nutr. 2003;23:171-201.
- * Caloric value is 1.5 Cal/mL compared to base PediaSure® at 1.0 Cal/mL.
- † Vitamins C & E and selenium.
- [‡] 32 mg of DHA per 8-fl-oz serving.

NutraFlora and scFOS are not registered trademarks of Abbott Laboratories.

Availability: Hospital/Institutional

8-fl-oz can; 24/case

Flavor	List No.
Vanilla	67374

Ready To Hang

1000-mL prefilled containers, non-spikeable cap; 8/case
Flavor List No.

Vanilla Ready to Feed: Water, Corn Maltodextrin, Blend Of Vegetable Oils (High Oleic Safflower, Soy, Medium Chain Triglycerides), Milk Protein Concentrate, Soy Protein Isolate, Short-Chain Fructooligosaccharides; Less than 0.5% of: Vitamins And Minerals', Oat Fiber, Natural And Artificial Flavor, Soy Fiber, Cellulose Gel, Soy Lecithin, Tuna Oil*, Monoglycerides, Cellulose Gum, Potassium Hydroxide, Carrageenan, Inositol, Taurine, Sucralose, Salt, L-Carnitine and Lutein. *Vitamins And Minerals: Potassium Chrate, Potassium Chloride, Magnesium Phosphate, Calcium Phosphate, Ascorbic Acid, Choline Chloride, Calcium Carbonate, Potassium Phosphate, Ferrous Sulfate, dl-Alpha-Tocopheryl Acetate, Zinc Sulfate, Niacinamide, Calcium Pantotherate, Manganese Sulfate, Thiamine Hydrochloride, Pyridoxine Hydrochloride, Riboflavin, Vitamin A Palmitate, Copper Sulfate, Folic Acid, Chromium Chloride, Potassium Iodide, Sodium Molybdate, Sodium Selenate, Biotin, Phylloquinone, Vitamin B., Contains milk and soy ingredients.

NUTRITION INFO	RMATION						
	8 fl oz (237 mL)	1000 mL‡	1500 mL‡		8 fl oz (237 mL)	1000 mL‡	1500 mL‡
Energy, Cal	350	1500	2250	Riboflavin (Vit B ₂), mg	0.33	1.4	2.1
Protein, g	14	59	89	Vitamin B ₆ , mg	0.34	1.4	2.1
% Total Cal	16	16	16	Vitamin B ₁₂ , mcg	0.48	2	3
Source	Milk Protein Cond	entrate, Soy Prot		Niacin, mg NE	3.2	14	20
L-Carnitine, mg	4	17	25	Choline, mg	80	338	506
Taurine, mg	18	76	114	Biotin, mcg	8	34	50
Fat, g	16	67	100	Pantothenic Acid, mg	1.3	5.5	8.2
Saturated Fat, g	3.5	14	22	Inositol, mg	20	84	127
Trans Fat, g	0	0	0	Minerals			
Polyunsaturated Fat, g	5	20	30	Calcium, mg	330	1390	2080
Monounsaturated Fat, g	7	30	44	Calcium, mEq	16.5	70	104
% Total Cal	41	41	41	Phosphorus, mg	250	1050	1580
Source		ver Oil, Soy Oil, Medii Triglycerides	um Chain	Magnesium, mg	40	169	253
Oil Ratio		50:40:10		Iron, mg	2.7	11	17
Cholesterol, mg	10	40	63	Zinc, mg	1.7	7.2	10.8
Carbohydrate, g	39	166	247	Manganese, mg	0.46	1.9	2.9
% Total Cal	43	43	43	Copper, mg	0.14	0.59	0.89
Source	Con	n Maltodextrin		lodine, mcg	23	97	145
Total Sugars, g	3	11	17	Selenium, mcg	8	34	51
Added Sugars, g	3	10	15	Chromium, mcg	9	38	57
Fiber, g	3	13	19	Molybdenum, mcg	9	38	57
Prebiotic	Short Chain I	Fructooligosacch	arides	Sodium, mg	90	380	570
Vitamins				Sodium, mEq	3.9	16.5	24.7
Vitamin A, mcg RAE	140	590	886	Potassium, mg	470	1980	2970
Vitamin D, mcg	6	25	38	Potassium, mEq	12.1	51	76
Vitamin E, mg	3	13	19	Chloride, mg	230	970	1450
Vitamin K, mcg	18	76	114	Chloride, mEq	6.5	27	41
Vitamin C, mg	23	97	146	Other Characteristics			
Folate, mcg DFE	60	253	380	PRSL, mOsm	110	465	698
Folic Acid, mcg	36	152	227	Water, g	185	780	1170
Thiamin (Vit B ₁), mg	0.3	1.3	1.9	Osmolality, mOsm/kg H ₂ O	425	425	425

[‡] Source of docosahexaenoic acid (DHA). [¶] Values extrapolated from 8 fl oz.

PediaSure® Peptide 1.0 Cal

Peptide-Based Nutrition for Children



Description/Indications

PediaSure Peptide 1.0 Cal is a nutritionally complete, peptide-based formula for the nutritional needs of children ages 1 to 13 years with malabsorption, maldigestion, and other GI conditions.

- For oral or tube feeding
- For supplemental or sole-source nutrition
- Use under medical supervision

Features

- Complete, Balanced Nutrition®
- Nutrient blend designed to help support tolerance and absorption:
 - Hydrolyzed whey-dominant protein
 - NutraFlora scFOS prebiotic[†]
 - 60% MCT & structured lipids
- Nutrition to help support growth and development
- Meets or exceeds 100% of the DRIs for protein and 25 essential vitamins and minerals for children 1 to 8 years of age in 1000 mL (1000 Cal), and for children 9 to 13 years of age in 1500 mL (1500 Cal)
- Antioxidants (vitamins C & E and selenium)

- DHA omega-3* to help support brain and eye health1,2
- * 32 mg DHA per 8-fl-oz serving.
- † NutraFlora and scFOS are not registered trademarks of Abbott Laboratories.
- 1. Lauritzen L, et al. Prog Lipid Res. 2001;40:1-94.
- 2. Krinsky NI, et al. Annu Rev Nutr. 2003;23:171-201.

Precautions

- Not intended for infants under 1 year of age unless specified by a physician
- · Not for IV use
- Not for children with galactosemia

Availability: Hospital/Institutional

/ tranability incopital, motitational	
8-fl-oz bottle; 24/case	
Flavor	List No.
Unflavored	67413
Vanilla	67407
Strawberry	67411
Chocolate	67073
Doody To Hone	

1000-mL prefilled containers; 8/case	
Flavor	List No
Unflavored	6741

Unflavored Ready to Feed: Water, Corn Maltodextrin, Whey Protein Hydrolysate, Structured Lipid (Interesterified Canola Oil & Medium Chain Triglycerides), Medium-Chain Triglycerides, Hydrolyzed Sodium Caseinate. Less than 0.5% of the following: Vitamins & Minerals (Calcium Phosphate, Potassium Citrate, Potassium Chloride, Magnesium Phosphate, Ascorbic Acid, Choline Chloride, Sodium Citrate, Ferrous Sulfate, Tinc Sulfate, di-Alpha-Tocopheryl Acetate, Niacinamide, Calcium Pantothenate, Manganese Sulfate, Thiamine Hydrochloride, Pyridoxine Hydrochloride, Riboflavin, Copper Sulfate, Vitamin A Palmitate, Folic Acid, Chromium Chloride, Biotin, Potassium Iodide, Sodium Selenate, Sodium Molybdate, Phylloquinone, Vitamin D₃, Vitamin B₁₂), Short-chain Fructooligosaccharides, Canola Oil, Cellulose Gel, Soy Lecithin, Tuna Oil², Cellulose Gum, Carrageenan, Inositol, Potassium Hydroxide, Taurine, L-Carnitine, and Lutein. Contains milk and soy ingredients.

	8 fl oz (237 mL)	1000 mL		8 fl oz (237 mL)	1000 mL
Energy, Cal	237	1000	Riboflavin (Vit B _a), mg	0.5	2.1
Protein, g	7	30	Vitamin B, mg	0.6	2.5
% Total Cal	12	12	Vitamin B ₁₂ , mcg	1.4	5.9
Source	Whey protein hydrolysate, hydrolyzed sodium ca	seinate	Niacin, mg NE	5.2	21.9
L-Carnitine, mg	4.0	17	Choline, mg	80	340
Taurine, mg	30	76	Biotin, mcg	20	84
Fat, g	9.6	41	Pantothenic Acid, mg	2.4	10
Saturated Fat, g	5	21	Inositol, mg	32	135
Trans Fat, g	0	0	Minerals		
Monounsaturated Fat, g	1.5	6	Calcium, mg	330	1390
Polyunsaturated Fat, g	0.5	2	Calcium, mEq	16.5	69.5
Cholesterol, mg	10	42	Phosphorus, mg	250	1055
% Total Cal	35	35	Magnesium, mg	40	170
Source	Structured lipid (Interesterified canola and medium-chain triglycerides), medium-chain triglycerides, canola oil		Iron, mg	3.3	14
Carbohydrate, g	31.7	134	Zinc, mg	2.8	12
% Total Cal	53	53	Manganese, mg	0.46	1.7
Source	Corn maltodextrin, scFOS ^{II}		Copper, mg	0.14	0.6
Ratio	97:3		lodine, mcg	23	97
Total Sugars, g	1	4.2	Selenium, mcg	8	34
Added Sugars, g	1	4.2	Chromium, mcg	9	38
Fiber, g	0.7	3.0	Molybdenum, mcg	9	38
Prebiotic	Short-Chain Fructooligosaccharide	S	Sodium, mg	170	717
Vitamins			Sodium, mEq	7.4	31.2
Vitamin A, mcg RAE	140	570	Potassium, mg	470	1980
Vitamin D, mcg	5.9	25	Potassium, mEq	12.05	50.8
Vitamin E, mg	2.5	10.4	Chloride, mg	240	1010
Vitamin K, mcg	16	68	Chloride, mEq	6.8	28.5
Vitamin C, mg	24	101	Other Characteristics		
Folate, mcg DFE	120	500	PRSL, mOsm	75	293
Folic Acid, mcg	71	300	Water, g	201	845
Thiamin (Vit B ₄), mg	0.6	2.5	Osmolality, mOsm/kg H ₂ O	290	290

[&]quot;scFOS = short-chain fructooligosaccharides.

[‡] Source of docosahexaenoic acid (DHA).

PediaSure® Peptide 1.5 Cal

Peptide-Based Nutrition for Children



Description/Indications

PediaSure Peptide 1.5 Cal is a nutritionally complete, peptide-based formula for the nutritional needs of children 1 to 13 years with malabsorption, maldigestion, and other Gl conditions. It is designed to meet the nutritional and caloric needs of children who require higher caloric density, have fluid restrictions, and are at risk for malnutrition.

- · For oral or tube feeding
- For supplemental or sole-source nutrition
- Use under medical supervision

Features

- Complete, Balanced Nutrition®
- Nutrient blend designed to help support tolerance and absorption for growth and development:
 - Hydrolyzed whey-dominant protein
 - NutraFlora scFOS prebiotic[†]
 - 60% MCT & structured lipids
- Meets or exceeds 100% of the DRIs for protein and 25 essential vitamins and minerals for children 1 to 8 years of age in 667 mL (1000 Cal), and for children 9 to 13 years of age in 1000 mL (1500 Cal)
- Antioxidants (vitamins C & E and selenium)

DHA omega-3* to help support brain and eye health^{1,2}

* 32 mg DHA per 8-fl-oz serving.

† NutraFlora and scFOS are not registered trademarks of Abbott Laboratories.

Precautions

- Not intended for infants under 1 year of age unless specified by a physician
- · Not for IV use

8-fl-oz hottle: 24/case

· Not for children with galactosemia

Availability: Hospital/Institutional

0-11-02 Dottie, 24/ case	
Flavor	List No.
Vanilla	67417
Ready To Hang	
1000-mL prefilled containers; 8/case	
Flavor	List No.
Vanilla	67419

Lauritzen L, et al. Prog Lipid Res. 2001;40:1-94.
 Krinksy NI, et al. Annu Rev Nutr. 2003;23:171-201.

Vanilla Ready to Feed: Water, Corn Maltodextrin, Structured Lipid (Interesterified Canola Oil And Medium-Chain Triglycerides), Whey Protein Hydrolysate, Hydrolyzed Sodium Caseinate, Medium-Chain Triglycerides, Short-Chain Fructooligosaccharides; Less than 0.5% of: Vitamins and Minerals*, Canola Oil, Artificial Flavor, Cellulose Gel, Soy Lecithin, Tuna Oil*, Inositol, Taurine, Cellulose Gum, Potassium Hydroxide, Acesulfame Potassium, Sucralose, L-Carnitine And Lutein. * Vitamins and Minerals: Calcium Phosphate, Potassium Citrate, Potassium Chloride, Potassium Phosphate, Magnesium Phosphate, Calcium Carbonate, Sodium Citrate, Ascorbic Acid, Choline Chloride, Ferrous Sulfate, Magnesium Chloride, Zinc Sulfate, dl-Alpha-Tocopheryl Acetate, Niacinamide, Calcium Pantothenate, Manganese Sulfate, Thiamine Hydrochloride, Pyridoxine Hydrochloride, Riboflavin, Copper Sulfate, Vitamin A Palmitate, Folic Acid, Biotin, Potassium Iodide, Chromium Chloride, Sodium Selenate, Sodium Molybdate, Phylloquinone, Vitamin D., Vitamin B., Contains milk and soy ingredients.

	8 fl oz (237 mL)	1000 mL		8 fl oz (237 mL)	1000 mL
Energy, Cal	356	1500	Riboflavin (Vit B ₂), mg	0.8	3.3
Protein, g	10.7	45.1	Vitamin B _s , mg	1.0	4.2
% Total Cal	12	12	Vitamin B,, mcg	2.1	8.9
Source	Whey protein hydrolysate, hydrolyzed sodiur	n caseinate	Niacin, mg NE	7.6	31.9
L-Carnitine, mg	6.0	25	Choline, mg	120	510
Taurine, mg	45	190	Biotin, mcg	30	127
Fat, g	14.4	60.8	Pantothenic Acid, mg	3.6	15
Saturated Fat, g	8	35	Inositol, mg	47	200
Trans Fat, g	0	0	Minerals		
Polyunsaturated Fat, g	1	4.2	Calcium, mg	500	2090
Monounsaturated Fat, g	2.5	10.5	Calcium, mEq	25	104.5
Cholesterol, mg	15	63	Phosphorus, mg	380	1580
% Total Cal	35	35	Magnesium, mg	60	255
Source	Structured lipid (Interesterified canola and m triglycerides), medium-chain triglycerides,		Iron, mg	5.0	21
Carbohydrate, g	47.6	201	Zinc, mg	4.3	18
% Total Cal	53	53	Manganese, mg	0.7	2.9
Source	Corn maltodextrin, scFOS ^{II}		Copper, mg	0.2	0.89
Ratio	97:3		lodine, mcg	35	148
Total Sugars, g	2	7	Selenium, mcg	12	51
Added Sugars, g	2	7	Chromium, mcg	14	57
Fiber, g	1	4.6	Molybdenum, mcg	14	57
Prebiotic	Short-Chain Fructooligosaccharide	es	Sodium, mg	255	1075
Vitamins			Sodium, mEq	11.1	46.7
Vitamin A, mcg RAE	200	850	Potassium, mg	710	2975
Vitamin D, mcg	5.9	25	Potassium, mEq	18.2	76.3
Vitamin E, mg	3.7	15.5	Chloride, mg	360	1520
Vitamin K, mcg	24	101	Chloride, mEq	10.1	42.8
Vitamin C, mg	36	152	Other Characteristics		
Folate, mcg DFE	180	750	PRSL, mOsm	108	454
Folic Acid, mcg	107	450	Water, g	182	768
Thiamin (Vit B ₄), mg	1.0	4.2	Osmolality, mOsm/kg H _o O	480	480

[&]quot;scFOS = short-chain fructooligosaccharides. † Source of docosahexaenoic acid (DHA).

Vital® Peptide 1.5 Cal

Peptide-Based Nutrition for Ages 14+



Description/Indications

Vital Peptide 1.5 Cal is a nutritionally complete, peptide-based formula for the nutrition needs of patients ages 14+ years with malabsorption, maldigestion, and other GI conditions.

- · For tube or oral feeding
- For supplemental or sole-source nutrition
- Use under medical supervision

Features

- Nutrient blend designed to help support tolerance and absorption
- Hydrolyzed whey-dominant protein
- NutraFlora scFOS, a prebiotic fiber*
- 55% MCT & structured lipids
- Complete, Balanced Nutrition®
- 1000 IU/L vitamin D
- · Antioxidants vitamins C & E and selenium
- Meets or exceeds 100% of the DRIs for protein and 25 essential vitamins and minerals for patients 14 to 18 years of age in 1500 Cal (1000 mL); for patients 18+ years of age in 1800 Cal (1200 mL)
- Kosher, Halal, gluten-free, suitable for lactose intolerance

Precautions

- Not for IV use
- · Not for people with galactosemia

Availability: Hospital/Institutional

8-fl-oz reclosable carton; 24/case
Flavor List No.
Vanilla 66236

^{*} NutraFlora and scFOS are not registered trademarks of Abbott Laboratories.

Vanilla Ready to Feed: Water, Corn Maltodextrin, Whey Protein Hydrolysate, Structured Lipid (Interesterified Canola Oil And Medium Chain Triglycerides), Sugar, Hydrolyzed Sodium Caseinate; Less than 1% of: Canola Oil, Medium Chain Triglycerides, Short-Chain Fructooligosaccharides, Magnesium Phosphate, Natural And Artificial Flavor, Potassium Citrate, DATEM, Calcium Phosphate, Cellulose Gel, Potassium Chloride, Calcium Carbonate, Choline Chloride, Sodium Phosphate, Ascorbic Acid, Potassium Hydroxide, Sucralose, Cellulose Gum, Carrageenan, Magnesium Chloride, L-Carnitine, Taurine, Zinc Sulfate, Ferrous Sulfate, d-Alpha-Tocopheryl Acetate, Niacinamide, Manganese Sulfate, Calcium Pantothenate, Copper Sulfate, Thiamine Hydrochloride, Pyridoxine Hydrochloride, Vitamin A Palmitate, Riboflavin, Chromium Chloride, Folic Acid, Sodium Molybdate, Sodium Selenate, Potassium Iodide, Phylloquinone, Biotin, Vitamin B,, Contains milk ingredients.

	8 fl oz (237 mL)	1000 mL		8 fl oz (237 mL)	1000 mL
Energy, Cal	355	1500	Biotin, mcg	85	359
Protein, g	16	67.5	Pantothenic Acid, mcg	2.4	10
% Total Cal	18	18	Minerals		
Source	Whey protein hydrolysate, hydrolyzed sodiur	n caseinate	Calcium, mg	310	1308
L-Carnitine, mg	5.3	22	Calcium, mEq	15.5	65.4
Taurine, mg	23	97	Phosphorus, mg	300	1266
Fat, g	13.5	57.1	Magnesium, mg	100	422
% Total Cal	33	33	Iron, mg	4.3	18
Source	Structured lipid (Interesterified canola and med triglycerides), medium-chain triglycerides, ca		Zinc, mg	2.7	11
Carbohydrate, g	44.3	187	Manganese, mg	0.6	2.5
% Total Cal	49	49	Copper, mg	0.25	1.1
Source	Corn Maltodextrin, Sugar, scFOS		lodine, mcg	36	152
Fiber, g	1.4	6	Selenium, mcg	15	60
Vitamins			Chromium, mcg	11	46
Vitamin A, IU	775	3270	Molybdenum, mcg	18	76
Vitamin D, IU	240	1013	Sodium, mg	360	1519
Vitamin E, IU	8.1	34	Sodium, mEq	15.7	66
Vitamin K, mcg	25	105	Potassium, mg	480	202
Vitamin C, mg	40	169	Potassium, mEq	12.3	51.9
Folic Acid, mcg	95	401	Chloride, mg	360	151
Thiamin (Vit B ₁), mg	0.5	2.1	Chloride, mEq	10.2	42.
Riboflavin (Vit B ₂), mg	0.5	2.1	Other Characteristics		
Vitamin B ₆ , mg	0.5	2.1	PRSL, mOsm	136	58
Vitamin B ₁₂ , mcg	0.7	3	Water, g	181	76
Niacin, mg NE	3.8	16	Osmolality, mOsm/kg H ₂ O	610	61
Choline, mg	131	553			

[†] scFOS = short-chain fructooligosaccharides.

Similac® Organic Toddler Drink with A2 Milk

Milk-Based Powder



Description/Indications

A non-GMO* milk-based toddler drink with A2 milk to complement toddlers' (12-36 months) diets that is certified USDA Organic and is the first organic toddler drink made with A2 milk.

Features

- First easy-to-digest toddler drink that is USDA-certified organic and made with A2 milk
- Made with premium ingredients, including organic lactose
- · Gentle nutrition and made with A2 milk protein
- Formulated with a unique blend of DHA, lutein, and vitamin E for brain and eye development
- Certified USDA Organic
- Non-GMO*
- Has no artificial growth hormones[†]
- No palm olein oil
- Kosher
- Halal
- Gluten-free
- Has 28 key nutrients to help support growth and development

Precaution

· Not for children with galactosemia

Preparation and Use

- Add 4 fl oz of water to clean sippy cup.
- Using a measuring utensil for dry ingredients, add 2 tablespoons of powder to water.
- Shake well; use within 1 hour; then discard.

Availability: Retail

Size Container List No.

20.6 oz (584 g); about 45 servings container/4 case....68104

^{*} Ingredients not genetically engineered.

[†] No significant difference has been shown between milk derived from rbST-treated and non-rbST-treated cows.

Powder: Organic Lactose, Organic Nonfat Milk, Organic High Oleic Sunflower Oil, Organic Soy Oil, Organic Coconut Oil; Less than 2% of: C. Cohnii Oil, Beta-Carotene, Lutein, Potassium Citrate, Tricalcium Phosphate, Ascorbic Acid, Ascorbyl Palmitate, Ferrous Sulfate, Salt, Choline Chloride, Taurine, Inositol, Magnesium Chloride, Zinc Sulfate, Mixed Tocopherols, d-Alpha-Tocopheryl Acetate, Niacinamide, Calcium Pantothenate, L-Carnitine, Vitamin A Palmitate, Copper Sulfate, Thiamine Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Manganese Sulfate, Phylloquinone, Biotin, Chromium Chloride, Sodium Molybdate, Sodium Selenate, Vitamin D₃, Vitamin B₁₂, Potassium Iodide, and Potassium Hydroxide. Contains milk ingredients.

	Amount	%DV‡		Amount	%DV‡
Serving Size	2 Tablespoons (14.5g)	_	Vitamin B _e , mg	0.125	25
Energy, Cal	70		Vitamin B ₁₃ , mcg	0.225	25
Protein, g	1	8	Niacin, mg	1.2	20
Fat, g	4	10	Choline, mg	40	20
Saturated Fat, g	1.5	15	Biotin, mcg	2.4	30
Polyunsaturated Fat, g	1		Pantothenic Acid, mg	0.4	20
Monounsaturated Fat, g	1.5		Minerals		
Trans Fat, g	0		Calcium, mg	140	20
Cholesterol, mg	0	0	Phosphorus, mg	92	20
Carbohydrate, g	8	5	Magnesium, mg	6.4	8
Total Sugars, g	8		Iron, mg	2.5	35
Added Sugars, g	6	24	Zinc, mg	0.45	15
Dietary Fiber, g	0	0	Manganese, mg	0.12	10
Vitamins			Copper, mg	0.06	20
Vitamin A, mcg RAE	30	10	lodine, mcg	22.5	25
Vitamin D, mcg	4.5	30	Selenium, mcg	2.0	10
Vitamin E, mg	2.4	40	Chromium, mcg	3.3	30
Vitamin K, mcg	5.0	15	Molybdenum, mcg	3.4	20
Vitamin C, mg	5.25	35	Sodium, mg	25	2
Folate, mcg DFE (9 mcg Folic Acid)	15	10	Potassium, mg	100	4
Thiamin, mg	0.075	15	Chloride, mg	60	4
Riboflavin (Vit B ₂), mg	0.25	50			

[‡]The % Daily Value (DV) tells you how much a nutrient in a serving of food contributes to a daily diet. 1,000 calories a day is used for general nutrition advice. C. Cohnii Oil is a source of DHA.

Pure Bliss° by Similac° Toddler Drink

Milk-Based Powder



Description/Indications

A non-GMO,* milk-based drink for toddlers 12 to 36 months old. Starts with fresh milk from grass-fed cows, and has no artificial growth hormones[†] and no antibiotics.

Features

- A non-GMO* toddler drink option
- No artificial growth hormones[†]
- · Contains no antibiotics
- 28 key nutrients in 2 servings[‡]
- Prebiotics to help promote digestive health
- DHA and Lutein to help support brain and eye development
- · Gluten-free
- Kosher, Halal

Precaution

Not for children with galactosemia

Preparation and Use

- Add 4 fl oz of water to clean sippy cup.
- Using a measuring utensil for dry ingredients, add 2 tablespoons of powder to water.
- Shake well; use within 1 hour; then discard.

Availability: Retail

Size Container List No.

Powder:

24.7 oz (700 g); about 38 servings...can; 4/carton 67319

 $^{^{\}star}$ Ingredients not genetically engineered.

 $^{{}^\}dagger \text{No}$ significant difference has been shown between milk derived from rbST-treated and non-rbST-treated cows.

[‡]1 serving is defined as 18 g reconstituted with 4 fl oz water.

Powder: Nonfat Milk, Lactose, High Oleic Sunflower Oil, Soy Oil, Galactooligosaccharides, Coconut Oil, Less than 2% of: C. Cohnii Oil, M. Alpina Oil, Bifidobacterium Lactis, Beta-Carotene, Lutein, Potassium Citrate, Calcium Carbonate, Ascorbic Acid, Soy Lecithin, Ferrous Sulfate, Choline Bitartrate, Ascorbyl Palmitate, Taurine, Mixed Tocopherols, dl-Alpha-Tocopheryl Acetate, Vitamin A Palmitate, Copper Sulfate, Thiamine Hydrochloride, Polic Acid, Manganese Sulfate, Vitamin D₃, Potassium Hydroxide, Magnesium Chloride, Nucleotides (Cytidine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate), Potassium Phosphate, and Potassium Chloride. Contains milk and soy ingredients.

	Amount	%DV [‡]		Amount	%DV [‡]
Serving Size	2 Tablespoons (18 g)		Vitamin B ₆ , mg	0.07	15
Energy, Cal	80		Vitamin B ₁₂ , mcg	0.2	25
Protein, g	3	23	Niacin, mg NE	0.54	8
Fat, g	4	10	Choline, mg	12.4	6
Saturated Fat, g	1.5	15	Biotin, mcg	1.6	20
Trans fat, g	0		Pantothenic Acid, mg	0.15	8
Polyunsaturated Fat, g	1		Minerals		
Monounsaturated Fat, g	2		Calcium, mg	140	20
Cholesterol, mg	less than 5	1	Phosphorus, mg	88	20
Carbohydrate, g	9	6	Magnesium, mg	9	10
Dietary Fiber, g	0	0	Iron, mg	1	15
Total Sugars, g	9		Zinc, mg	0.2	8
Added Sugars, g	4	16	Manganese, mg	0.06	4
Vitamins			Copper, mg	0.01	4
Vitamin A mcg RAE	20	8	lodine, mcg	11.4	15
Vitamin D, mcg	1.8	10	Selenium, mcg	1	6
Vitamin E, mg	1.9	30	Sodium, mg	35	2
Vitamin K, mcg	2	8	Potassium, mg	180	6
Vitamin C, mg	2	15	Chloride, mg	81	6
Thiamin, mg	0.04	8	Other Characteristics		
Folate (15 mcg Folic Acid)	25	15	Osmolality, mOsm/kg H ₂ O	303	
Riboflavin (Vit B ₂), mg	0.2	35			

[‡]The % Daily Value (DV) tells you how much a nutrient in a serving of food contributes to a daily diet. 1,000 calories a day is used for general nutrition advice.

Galactooligosaccharides (GOS) are prebiotics sourced from milk.

C. Cohnii Oil is a source of DHA.

M. Alpina Oil is a source of ARA.

Go & Grow by Similac[®] Non-GMO^{*} with 2'-FL HMO Toddler Drink

Milk-Based Powder



Description/Indications

A milk-based, non-GMO* drink to complement toddlers' (12 to 36 months) diets that has 2'-FL HMO.

Features

- Supports your toddler's immune system and brain & eye development
- Two 4-fl-oz servings[‡] have 25 key nutrients like iron, vitamins C. D. & E
- No artificial growth hormonest
- Gluten-free
- Kosher, Halal

Precaution

· Not for children with galactosemia

Preparation and Use

- Add 4 fl oz of water to clean sippy cup.
- Using a measuring utensil for dry ingredients, add 2 tablespoons of powder to water.
- Shake well; use within 1 hour; then discard.

Availability: Retail

Size	Container	List No.
Powder:		
0.6 oz (17 g); 16 servings (1	0.6-oz packet per ser	ving)
	packet; 16/cartor	n 67149
24 oz (680 g); 45 servings	can; 4/case	67151
36 oz (1.02 kg); 68 servings.	can; 3/case	67146
40 oz (1.13 kg); 75 servings.	can; 6/carton	67153

^{*} Ingredients not genetically engineered.

[‡] 1 serving is defined as 15g or 17.4g (stick pack) reconstituted in 4 fl oz of water

 $^{^\}dagger$ No significant difference has been shown between milk derived from rbST-treated and non-rbST-treated cows.

Powder: Nonfat Milk, Lactose, High Oleic Safflower Oil, Soy Oil, Coconut Oil; Less than 2% of: Short-Chain Fructooligosaccharides, Calcium Phosphate, Potassium Hydroxide, 2'-Fucosyllactose, Ascorbic Acid, Soy Lecithin, Potassium Phosphate, Choline Chloride, C. Cohnii Oil, Ferrous Sulfate, Taurine, Ascorbyl Palmitate, Inositol, Mixed Tocopherols, d-Alpha-Tocopheryl Acetate, Zinc Sulfate, Niacinamide, Potassium Citrate, Manganese Sulfate, Calcium Pantothenate, Vitamin A Palmitate, Thiamine Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Copper Sulfate, Lutein, Beta-Carotene, Folic Acid, Vitamin D., Phylloquinone, Sodium Selenate, Biotin, and Vitamin B., Contains milk and soy ingredients.

	Amount	%DV [‡]		Amount	%DV [‡]
Serving Size	2 Tablespoons (18 g)		Vitamin B ₁₂ , mcg	0.08	8
Energy, Cal	70		Niacin, mg NE	0.9	15
Protein, g	2	15	Choline, mg	20	8
Fat, g	4	10	Biotin, mcg	1.0	10
Saturated Fat, g	1.5	15	Pantothenic Acid, mg	0.1	6
Polyunsaturated Fat, g	1		Minerals		
Monounsaturated Fat, g	1.5		Calcium, mg	150	20
Trans Fat, g	0		Phosphorus, mg	80	15
Cholesterol, mg	0	0	Magnesium, mg	5	8
Carbohydrate, g	7	6	Iron, mg	1.5	20
Total Sugars, g	7		Zinc, mg	0.1	2
Added Sugars, g	4	16	Manganese, mg	0.05	4
Dietary Fiber, g	0	0	Copper, mg	0.02	6
Vitamins			lodine, mcg	10	10
Vitamin A mcg RAE	30	10	Selenium, mcg	2.0	10
Vitamin D, mcg	3.2	20	Chromium, mcg	1.0	10
Vitamin E, mg	1.2	20	Molybdenum, mcg	2.0	10
Vitamin K, mcg	3.0	10	Sodium, mg	25	2
Vitamin C, mg	3.0	20	Potassium, mg	130	4
Thiamin, mg	0.04	8	Chloride, mg	60	4
Riboflavin (Vit B ₂), mg	0.11	20	Other Characteristics		
Vitamin B, mg	0.04	8	Osmolality, mOsm/kg H _o O	295	

[‡]The % Daily Value (DV) tells you how much a nutrient in a serving of food contributes to a daily diet. 1,000 calories a day is used for general nutrition advice. C. Cohnii Oil is a source of DHA.

^{2&#}x27;-fucosyllactose (2'-FL): a prebiotic like that found in breast milk.

Go & Grow by Similac® Toddler Drink

Milk-Based Powder



Description/Indications

A milk-based toddler drink for toddlers 12 to 36 months of age to help balance their diets.

Features

- OptiGRO® our unique blend of brain and eye nutrients, like those found in food, such as:
 - DHA from fish
 - Lutein from spinach
 - Vitamin E from broccoli
- Two 4-fl-oz servings† of Go & Grow by Similac have 25 key nutrients and at least 40% of the recommended Daily Value of calcium, iron, and vitamins C, D, and E
- · Complements toddler's diet
- Gluten-free
- · Kosher, Halal

Precaution

Not for children with galactosemia

Preparation and Use

- Add 4 fl oz of water to clean sippy cup.
- Using a measuring utensil for dry ingredients, add 2 tablespoons of powder to water.
- Shake well; use within 1 hour; then discard.

Availability: Retail

Size	Container	List No.
Powder:		
24 oz (680 g); 45 servings	can; 6/case	64782
30.8 oz (873 g); 58 servings	can; 6/case	66386
36 oz (1.13 kg); 68 servings	can; 3/case	66838

^{*} Daily Values for children 1 to 3 years old.

^{† 1} serving is defined as 15g reconstituted in 4 fl oz of water

Powder: Nonfat Milk, Lactose, High Oleic Safflower Oil, Soy Oil, Coconut Oil; Less than 2% of: Short-Chain Fructooligosaccharides, Calcium Phosphate, Potassium Hydroxide, Ascorbic Acid, Potassium Phosphate, Choline Chloride, Soy Lecithin, C. Cohnii Oil, Ferrous Sulfate, Taurine, Ascorbyl Palmitate, Inositiol, Mixed Tocopherols, d-Alpha-Tocopheryl Acetate, Zinc Sulfate, Niacinamide, Potassium Citrate, Manganese Sulfate, Calcium Pantothenate, Vitamin A Palmitate, Thiaminen Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Copper Sulfate, Lutein, Beta-Carotene, Folic Acid, Vitamin D., Phylloquinone, Sodium Selenate, Biotin, and Vitamin B., Contains milk and soy ingredients.

	Amount	%DV*		Amount	%DV*
Serving Size	2 Tablespoons (15 g)		Vitamin B,, mcg	0.08	8
Energy, Cal	70		Niacin, mg	0.4	15
Protein, g	2	15	Choline, mg	20	8
Fat, g	4	10	Biotin, mcg	1.0	10
Saturated Fat, g	1.5	15	Pantothenic Acid, mg	0.1	6
Polyunsaturated Fat, g	1		Minerals		
Monounsaturated Fat, g	1.5		Calcium, mg	150	20
Trans fat, g	0		Phosphorus, mg	80	15
Cholesterol, mg	0	0	Magnesium, mg	5	8
Carbohydrates, g	7	5	Iron, mg	1.5	20
Total Sugars, g	7		Zinc, mg	0.1	2
Added Sugars, g	4	16	Manganese, mg	0.05	4
Dietary Fiber, g	0	0	Copper, mg	0.02	6
Vitamins			lodine, mg	10	10
Vitamin A, mcg RAE	30	10	Selenium, mcg	2.0	10
Vitamin D, mcg	3.2	20	Chromium, mcg	1.0	10
Vitamin E, mg	1.2	20	Molybdenum, mcg	2.0	10
Vitamin K, mcg	3.0	10	Sodium, mg	25	2
Vitamin C, mg	3.0	20	Potassium, mg	130	4
Folate, mcg DFE (3.85 mcg Folic Acid)	5	4	Chloride, mg	60	4
Thiamin, mg	0.04	8	Other Characteristics		
Riboflavin (Vit B ₂), mg	0.11	20	Osmolality, mOsm/kg H ₂ O	295	
Vitamin B _s , mg	0.04	8			

^{*}The % Daily Value (DV) tells you how much a nutrient in a serving of food contributes to a daily diet. 1,000 calories a day is used for general nutrition advice. C. Cohnii Oil is a source of DHA.

Go & Grow by Similac[®] Sensitive Non-GMO^{*} with 2'-FL HMO Toddler Drink

Milk-Based Powder



Description/Indications

A milk-based, non-GMO* drink to complement toddlers' (12 to 36 months) diets that has 2'-FL HMO and is suitable for lactose sensitivity.

Features

- An easy-to-digest, milk-based toddler drink suitable for children with lactose sensitivity
- Supports your toddler's immune system and brain & eye development
- Two 4-fl-oz servings† of Go & Grow by Similac have 23 key nutrients like iron, calcium, and vitamins C, D, & E.
- · Gluten-free
- Kosher, Halal

Precaution

· Not for children with galactosemia

Preparation and Use

- Add 4 fl oz of water to clean sippy cup.
- Using a measuring utensil for dry ingredients, add 2 tablespoons of powder to water.
- Shake well; use within 1 hour; then discard.

Availability: Retail

Size	Container	List No.
Powder:		
23.3 oz (661 g); 44 servings	scan; 4/case	67581

^{*} Ingredients not genetically engineered.

[†] 1 serving is defined as 15g reconstituted in 4 fl oz of water.

Powder: Corn Syrup, Milk Protein Concentrate, High Oleic Safflower Oil, Sugar, Soy Oil, Coconut Oil, Calcium Phosphate; Less than 2% of: Short-Chain Fructooligosaccharides, Potassium Citrate, Potassium Chloride, Sodium Citrate, Magnesium Phosphate, 2'-Fucosyllactose, Choline Chloride, Calcium Carbonate, Magnesium Chloride, C. Cohnii Oil, Ascorbyl Palmitate, Ascorbic Acid, L-Cystine Dihydrochloride, Ferrous Sulfate, Taurine, Inositol, Mixed Tocopherols, L-Carnitine, d-Alpha-Tocopheryl Acetate, Zinc Sulfate, Niacinamide, Manganese Sulfate, Calcium Pantothenate, Vitamin A Palmitate, Thiamine Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Copper Sulfate, Potassium Iodide, Lutein, Beta-Carotene, Folic Acid, Vitamin D₃, Phylloquinone, Sodium Selenate, Biotin, and Vitamin B.,... Contains milk ingredients.

	Amount	%DV*		Amount	%D V *
Serving Size	2 Tablespoons (15g)		Vitamin B ₁₂ mcg	0.08	8
Energy, Cal	70		Niacin, mg	0.8	15
Protein, g	2	15	Choline, mg	17	8
Fat, g	4	10	Biotin, mcg	1.0	10
Saturated Fat, g	1.5	15	Pantothenic Acid, mg	0.1	6
Trans Fat, g	0		Minerals		
Polyunsaturated Fat, g	1		Calcium, mg	150	20
Monounsaturated Fat, g	1.5		Phosphorus, mg	80	15
Cholesterol, mg	0	0	Magnesium, mg	5	8
Carbohydrate, g	8	5	Iron, mg	1.5	20
Dietary Fiber, g	0	0	Zinc, mg	0.1	2
Total Sugars, g	3		Manganese, mg	0.05	4
Added Sugars, g	3	12	Copper, mg	0.02	6
Vitamins			lodine, mcg	10	10
Vitamin A, mcg RAE	30	10	Selenium, mcg	1.0	4
Vitamin D, mcg	3.2	20	Chromium, mcg	1.0	10
Vitamin E, mg	1.2	20	Molybdenum, mcg	1.0	4
Vitamin K, mcg	3.0	10	Sodium, mg	25	2
Vitamin C, mg	3.0	20	Potassium, mg	130	4
Thiamin, mg	0.04	8	Chloride, mg	60	4
Folate, mcg DFE (4.02 mcg Folic Acid)	5	4	Other Characteristics		
Riboflavin (Vit B ₂), mg	0.11	20	Osmolality, mOsm/kg H ₂ O	200	
Vitamin B _e mg	0.04	8			

^{*} The % Daily Value (DV) tells you how much a nutrient in a serving of food contributes to a daily diet. 1,000 calories a day is used for general nutrition advice.

C. Cohnii Oil is a source of DHA.

^{2&#}x27;-fucosyllactose (2'-FL): a prebiotic like that found in breast milk.





Pedialyte® Classic

Electrolyte Solution for All Ages



Description/Indications

Pedialyte Classic replenishes vital fluids lost due to diarrhea to help prevent dehydration* in infants, children, and adults. A scientifically designed balance of electrolytes and glucose for fast rehydration.

Features

- #1 Pediatrician & Pharmacist recommended brand for hydration
- Medical-grade hydration
- Designed to promote fluid absorption more effectively than common household beverages
- Provides glucose to promote sodium and water absorption
- Quickly replaces fluids and zinc[†]
- · Unflavored liquid available for infants
- Kid-approved taste to encourage compliance in children 1 year of age and older
- · Low osmolality (see table)
- · Kosher, Halal (excluding Mixed Fruit), gluten-free

Precautions

- Consult your doctor for children under 1 year of age.
 Powder packs not for use under 1 year of age.
- Use under medical supervision.
- Mild to moderate dehydration.
 † Liquid only.

Availability: Hospital/Institutional

Ready To Use

2-fl-oz plastic bottle; 48/case	List No.
Unflavored	59892

Availability: Retail

1-Liter (33.6-11-02) plastic bottle; o/case	
Unflavored	00336
Grape	00240
Mixed Fruit	
Strawberry	53983
Coconut Burst (4/case)	
,	

0.6-oz Powder Packs; 6 packs/carton

otrawberry Lerrioriade	04112
Orange	64177
Cherry	
Grapé	64598

6/172

0.3-oz Powder Packs; 8 packs/carton

Liters

Unflavored: Water, Dextrose, Less than 2% of: Potassium Citrate, Salt, Sodium Citrate, Citric Acid, and Zinc Gluconate.

Grape: Water, Dextrose, Less than 0.5% of: Citric Acid, Potassium Citrate, Salt, Artificial Flavor, Sodium Citrate, Sucralose, Acesulfame Potassium. Zinc Gluconate. Red 40, and Blue 1.

Mixed Fruit: Water, Dextrose. Less than 0.5% of: Citric Acid, Natural & Artificial Flavor, Potassium Citrate, Salt, Sodium Citrate, Sucralose, Acesulfame Potassium, Zinc Gluconate, and Yellow 6.

Strawberry: Water, Dextrose. Less than 0.5% of: Citric Acid, Potassium Citrate, Salt, Sodium Citrate, Natural Flavor, Sucralose, Acesulfame Potassium. Zinc Gluconate. Red 40. and Blue 1.

Coconut Burst: Water, Dextrose. Less than 0.5% of: Citric Acid, Potassium Citrate, Salt, Sodium Citrate, Natural Flavor, Sucralose, Acesulfame Potassium, and 7 inc Gluconate.

Powder Packs:

Strawberry Lemonade (17 g): Anhydrous Dextrose, Citric Acid, Potassium Citrate, Salt, Natural Flavors, Sodium Citrate. Less than 2% of: Calcium Silicate, Acesulfame Potassium, Sucralose, Red 40, Blue 1.

Orange (17 g): Anhydrous Dextrose, Citric Acid, Potassium Citrate, Salt, Sodium Citrate, Natural and Artificial Flavor. Less than 2% of: Calcium Silicate, Acesulfame Potassium, Sucralose, and Yellow 6.

Cherry (17 g): Anhydrous Dextrose, Citric Acid, Potassium Citrate, Salt, Sodium Citrate. Less than 2% of: Natural and Artificial Flavor, Calcium Silicate, Acesulfame Potassium, Sucralose, and Red 40.

Grape (17 g): Anhydrous Dextrose, Citric Acid, Potassium Citrate, Salt, Sodium Citrate. Less than 2% of: Natural and Artificial Flavor, Beet Powder Color, Calcium Silicate, Acesulfame Potassium, Sucralose, and Blue 1.

Fruit Punch (8.5 g): Anhydrous Dextrose, Citric Acid, Malic Acid, Potassium Citrate, Salt, Sodium Citrate. Less than 2% of: Natural and Artificial Flavor, Calcium Silicate, Sucralose, Acesulfame Potassium, and Red 40.

Grape (8.5 g): Anhydrous Dextrose, Citric Acid, Malic Acid, Potassium Citrate, Salt, Sodium Citrate. Less than 2% of: Natural and Artificial Flavor, Calcium Silicate, Sucralose, Acesulfame Potassium, Red 40, and Blue 1.

Apple (8.5 g): Anhydrous Dextrose, Citric Acid, Malic Acid, Potassium Citrate, Salt, Sodium Citrate. Less than 2% of: Artificial Flavor, Calcium Silicate, Caramel Color, Sucralose, and Acesulfame Potassium.

Strawberry (8.5 g): Anhydrous Dextrose, Citric Acid, Malic Acid, Potassium Citrate, Salt, Sodium Citrate, Natural Flavor. Less than 2% of: Calcium Silicate. Sucralose. Acesulfame Potassium. Red 40. and Blue 1.

NUTRITION INFORMATION	N*		
	1 L Grape Solution	8 fl oz Grape Powder (Reconstituted)	16 fl oz Grape Powder (Reconstituted)
Calories*	120	25	50
Protein, g	0	0	0
Carbohydrate*, g	32	6	12
Dietary Fiber, g	0	0	0
Sugars*, g	25	6	12
Added Sugars*, g	25	6	12
Fat, g	0	0	0
Sodium, mg	1030	240	490
Sodium, mEq	45	10	21
Potassium, mg	780	180	370
Potassium, mEq	20	5	9
Chloride, mg	1240	290	590
Chloride, mEq	35	8	17
Zinc, mg	7.8	0	0
Osmolality*, mOsm/kg H ₂ O	270	270	270

^{*} There may be small nutritional differences among flavors.

Dosage

Please refer to page 167 for the dosage administration guide for infants and young children.

Pedialyte AdvancedCare®

Electrolyte Solution for All Age



Description/Indications

Pedialyte AdvancedCare replenishes vital fluids lost due to diarrhea to help prevent dehydration* in infants, children, and adults. Scientifically designed balance of electrolytes and glucose for fast rehydration. Contains PreActiv® Prebiotics to help promote digestive health. Includes zinc and selenium for immune support.

Features

- #1 Pediatrician & Pharmacist Recommended Brand
- Designed to promote fluid absorption more effectively than common household beverages
- Provides glucose to promote sodium and water absorption
- Quickly replaces fluids and zinc
- PreActiv® Prebiotics help promote digestive health
- · Low osmolality (see table)
- Ready-to-use reclosable plastic liter bottles allow easy measuring and pouring
- Kid-approved taste to encourage compliance in children 1 year of age and older
- Kosher, Halal (excluding Cherry Punch), gluten-free

Precautions

- Consult your doctor for children younger than 1 year of age
- Use under medical supervision
- · No mixing or diluting is necessary or recommended
- · Not for children with galactosemia

Availability: Retail

1-Liter (33.8-fl-oz) bottle; 8/ct	List No.
Blue Raspberry	63059
Cherry Punch	63057
1-Liter (33.8-fl-oz) bottle; 4/ct	
Strawberry Lemonade	64301
Tropical Fruit	64307

Ingredients

Blue Raspberry: Water, Dextrose, Galactooligosaccharides. Less than 0.5% of: Citric Acid, Potassium Citrate, Salt, Natural and Artificial Flavors, Sodium Citrate, Sucralose, Acesulfame Potassium, Zinc Gluconate, Blue 1, Red 40, and Sodium Selenate. Contains Milk Ingredients.

Cherry Punch: Water, Dextrose, Galactooligosaccharides. Less than 0.5% of: Citric Acid, Natural and Artificial Flavors, Potassium Citrate, Salt, Sodium Citrate, Sucralose, Acesulfame Potassium, Zinc Gluconate, Red 40, and Sodium Selenate. Contains Milk Ingredients.

Strawberry Lemonade: Water, Dextrose, Galactooligosaccharides. Less than 0.5% of: Citric Acid, Potassium Citrate, Salt, Natural and Artificial Flavors, Sodium Citrate, Sucralose, Acesulfame Potassium, Zinc Gluconate, Red 40, Blue 1, and Sodium Selenate. Contains Milk Ingredients.

Tropical Fruit: Water, Dextrose, Galactooligosaccharides. Less than 0.5% of: Citric Acid, Potassium Citrate, Salt, Natural and Artificial Flavors, Sodium Citrate, Sucralose, Acesulfame Potassium, Zinc Gluconate, Yellow 6, and Sodium Selenate. Contains Milk Ingredients.

^{*} Mild to moderate dehydration.

Strawberry Lemonade	12 fl oz	1L
Calories	35	100
Protein, g	0	0
Carbohydrate, g	10	28
Dietary Fiber, g	<1	2
Sugars, g	7	19
Added Sugars, g	7	18
Fat, g	0	0
Selenium, mcg	11	31
Sodium, mg	370	1030
Sodium, mEq	16.0	45
Potassium, mg	280	780
Potassium, mEq	7.1	20
Chloride, mg	440	1240
Chloride, mEq	12.4	35
Zinc, mg	2.8	7.8
Osmolality, mOsm/kg H _o O	240 (CP 225)	240 (CP 225)

† There may be small nutritional differences among flavors.

Dosage

Refer to Administration Guide to restore fluid and minerals lost in diarrhea and vomiting. Pedialyte should be offered frequently in amounts tolerated. Total daily intake should be adjusted to meet individual needs, based on thirst and response to therapy.

The suggested intakes for maintenance are based on water requirements for ordinary energy expenditure.*

Extrapolated from Barness LA, Curran JS: Nutrition, in Nelson WE (sr ed), Behrman RE, Kliegman RIM, Arvin AM (eds): Nelson Textbook of Pediatrics, ed 15. Philadelphia: WB Saunders Co. 1996. pp 141-143.

ADMINISTRATION GUIDE FOR INFANTS AND YOUNG CHILDREN												
	Age	2	3	6	9	1	11/2	2	21/2	3	31/2	4
		Weeks		Months					Years			
Approximate	lb	9	14	18	21	23	26	28	30	32	34	36
Weight ¹	kg	4.0	6.4	8.2	9.5	10.5	11.8	12.7	13.6	14.4	15.3	16.3
Pedialyte for	fl oz/day	16	30	36	39	42	47	48	51	53	54	55
Maintenance ²		to	to	to	to	to	to	to	to	to	to	to
		20	34	42	45	47	52	53	56	57	57	59

Administration Guide does not apply to infants younger than 1 week of age. For children older than 4 years of age, maintenance intakes may exceed 2 liters daily. If there is vomiting or fever, or if diarrhea continues beyond 24 hours, consult the child's physician.

- Weight based on the 50th percentile of weight for age for boys from the National Center for Health Statistics (NCHS) Centers for Disease Control and Prevention (CDC) growth charts. Kuczmarski RJ, Ogden CL, Grummer-Strawn LM, et al: CDC Growth Charts: United States. Data from Vital and Health Statistics of the Centers for Disease Control and Prevention/National Center for Health Statistics. Advance Data, no. 314, December 4, 2000.
- 2. Fluid intake is total fluid requirement from oral electrolyte solution, formula, or other fluids, but does not take into account ongoing stool losses. Fluid loss in the stool should be replaced by consumption of an extra amount of Pedialyte equal to stool losses, in addition to the fluid maintenance requirement in this Administration Guide. Pedialyte Freezer Pops are to be used with Pedialyte Oral Electrolyte Solution or other appropriate fluids to help prevent dehydration.

Pedialyte® Electrolyte Water



Description/Indications

Pedialyte Electrolyte Water drink is designed for hydration purposes only, not mild to moderate dehydration. It is formulated with 3 key electrolytes and zinc for immune support.

Features

- Zero sugar*
- 3 key electrolytes for hydration, including sodium, potassium, and chloride
- · Zinc for immune support
- · No artificial flavors or colors
- Non-GMO[†]
- · Available in Berry Frost and Fruit Punch
- · Contains no fruit juice
- · Kosher, Halal, gluten-free

Precautions

- Not for children under 5 years of age for mild to moderate dehydration due to vomiting and diarrhea
- No mixing or diluting is necessary or recommended

Availability: Retail

1-Liter (33.8-fl-oz) bottle; 4/ct	List No.
Berry Frost	67966
Fruit Punch	67969

Ingredients

Berry Frost: Water; Less than 2% of: Natural Flavor, Citric Acid, Salt, Potassium Citrate, Sodium Citrate. Sucralose. Acesulfame Potassium. and Zinc Gluconate

Fruit Punch: Water; Less than 2% of: Natural Flavor, Citric Acid, Salt, Potassium Citrate, Sodium Citrate, Sucralose, Acesulfame Potassium, and Zinc Gluconate

^{*} Low in calories.

[†] Ingredients not genetically engineered.

NUTRITION INFORMAT	ΓΙΟΝ	
	12 fl oz	1L
Calories	5	10
Protein, g	0	0
Carbohydrate, g	1	3
Total Sugars, g	0	0
Added Sugars, g	0	0
Fat, g	0	0
Sodium, mg	240	690
Sodium, mEq	10.4	30
Potassium, mg	130	390
Potassium, mEq	3.3	10
Chloride, mg	250	700
Chloride, mEq	7.1	20
Zinc, mg	2.8	7.8
Osmolality, mOsm/kg H ₂ O	80	80



Description/Indications

From the #1 doctor recommended brand for hydration, Pedialyte with Immune Support provides hydration with zinc, vitamin C, and other key nutrients.

Features

- 4 key nutrients for immune support:
 - Zinc Builds new immune cells
 - Vitamin C Promotes cell health
 - Vitamin B₁₂ Helps activate immune cells
 - Vitamin E Supports cell membranes
- 33% more electrolytes*
- Available in Mixed Berry and Fruit Punch
- · Contains no fruit juice
- · Kosher, Halal, gluten-free

Precautions

- Not for use for children under 1 year of age
- Use under medical supervision

Availability: Retail

0.49-oz Powder Packs; 6 packs/carton, 6/case	List No.
Fruit Punch	67971
Mixed Berry	67974

Ingredients

Fruit Punch: Anhydrous Dextrose, Short-Chain Fructooligosaccharides, Salt, Citric Acid, Potassium Citrate, Sodium Citrate; Less than 2% of: Natural And Artificial Flavors, Calcium Silicate, Ascorbic Acid, Sucralose, Acesulfame Potassium, dl-Alpha-Tocopheryl Acetate, Zinc Sulfate, Red 40, And Vitamin B₁₂.

Mixed Berry: Anhydrous Dextrose, Salt, Short-Chain Fructooligosaccharides, Citric Acid, Potassium Citrate, Natural Flavor, Sodium Citrate; Less than 2% of: Calcium Silicate, Ascorbic Acid, Beet Powder Color, Sucralose, Acesulfame Potassium, dl-Alpha-Tocopheryl Acetate, Zinc Sulfate, Blue 1, And Vitamin B.,.

 $^{^{\}star}$ 60 mEq sodium electrolytes per liter vs 45 mEq in original Pedialyte® Classic.

	1 Packet (14.1 g)
Calories	40	
Protein, g	0	
Carbohydrate, g	12	
Dietary Fiber, g	1	
Sugars, g	8	
Added Sugars, g	8	
Fat, g	0	
Sodium, mg	650	
Sodium, mEq	60	
Potassium, mg	370	
Potassium, mEq	20	
Chloride, mg	840	
Chloride, mEq	50	
Zinc, mg	3.4	
Vitamin C, mg	90	
Vitamin E, mg	7.5	
Vitamin B12, mcg	2.4	
Osmolality, mOsm/kg H ₂ O	290	

Pedialyte® Sport



Description/Indications

From the #1 doctor recommended brand for hydration, Pedialyte Sport is a new advanced hydration option formulated for athletes with 5 key electrolytes for fast rehydration and muscle support. With a scientifically designed balance of sugar and electrolytes, Pedialyte Sport quickly replenishes fluids and is formulated to replace electrolytes you may lose during exercise.

Features

- Supplies 5 key electrolytes for fast rehydration and muscle support:
 - Sodium to avoid muscle cramps
 - Chloride for fluid balance
 - Potassium for muscle and nerve function
 - Magnesium for muscle health
 - Phosphate for muscle repair
- Increased number of electrolytes to help replace what's lost in sweat
- · Great taste
- · Available in Lemon Lime and Fruit Punch
- · Kosher, Halal, gluten-free

Precautions

- Consult your doctor for children younger than 2 years
- No mixing or diluting is necessary or recommended (liters only)
- · Not for children with galactosemia

Availability: Retail

1-Liter (33.8-fl-oz) plastic bottle; 4/case	List No.
Lemon Lime	67838
Fruit Punch	67840
0.49-oz Powder Packs; 6 packs/carton, 6/case	List No.
Lemon Lime	67845
Fruit Punch	67848

Ingredients

Lemon Lime: Water, Dextrose; Less than 1% of: Galactooligosaccharides, Salt, Potassium Citrate, Citric Acid, Potassium Phosphate, Natural Flavor With Other Natural Flavors, Magnesium Chloride, Sodium Citrate, Sucralose, Acesulfame Potassium, And Yellow 5. Contains milk ingredients.

Fruit Punch: Water, Dextrose; Less than 1% of: Galactooligosaccharides, Salt, Potassium Citrate, Citric Acid, Natural And Artificial Flavors, Potassium Phosphate, Magnesium Chloride, Sodium Citrate, Sucralose, Acesulfame Potassium, and Red 40. Contains milk ingredients.

Lemon Lime (14.1 g powder): Anhydrous Dextrose, Salt, Citric Acid, Short-Chain Fructooligosaccharides, Potassium Citrate, Potassium Phosphate, Magnesium Citrate, Natural And Artificial Flavors; Less than 2% of: Calcium Silicate, Acesulfame Potassium, Sucralose, and Yellow 5

Fruit Punch (14.1 g powder): Anhydrous, Salt, Citric Acid, Short-Chain Fructooligosaccharides, Potassium Citrate, Potassium Phosphate, Magnesium Citrate; Less than 2% of: Natural And Artificial Flavors, Calcium Silicate, Sucralose, Acesulfame Potassium, and Red 40.

NUTRITION INFORMA	TION	
	1 L Lemon Lime Solution	16 fl oz Lemon Lime Powder (Reconstituted)
Calories	85	40
Protein, g	0	0
Carbohydrate, g	25	10
Dietary Fiber, g	2	1
Sugars, g	14	7
Added Sugars, g	14	7
Fat, g	0	0
Sodium, mg	1380	650
Sodium, mEq	60	28
Potassium, mg	1320	600
Potassium, mEq	33	15
Phosphorus, mg	400	190
Chloride, mg	1940	920
Chloride, mEq	55	26
Magnesium, mg	120	55
Osmolality, mOsm/kg H ₂ O	270	270

Pedialyte® Organic



Description/Indications

Pedialyte Organic liters are an advanced hydration solution specially formulated with an optimal balance of sugar and electrolytes to help replenish vital fluids and minerals, which, when lost, can lead to dehydration. Pedialyte Organic liters can help prevent mild to moderate dehydration caused by vomiting, diarrhea, exercise, travel, and extreme heat.

Features

- USDA-certified organic
- No artificial colors, flavors, or sweeteners
- · Quickly replaces fluids and electrolytes
- Has zinc for immune support
- Designed to prevent dehydration more effectively than household beverages, like sports drinks, juice, and soda
- · Great for kids and adults!

Precautions

- Consult your doctor for children younger than 1 year
- Use under medical supervision
- No mixing or diluting is necessary or recommended

Availability: Retail

1-Liter (33.8-fl-oz) plastic bottle; 4/case	List No.
Crisp Lemon Berry	67301
Grape	67303

Ingredients

Crisp Lemon Berry: Water, Organic Apple Juice Concentrate, Less than 2% of: Organic Dextrose, Organic Citric Acid, Organic Flavor, Potassium Citrate, Salt, Sodium Citrate, Organic Stevia Leaf Extract, And Zinc Gluconate.

Grape: Water, Organic Grape Juice Concentrate, **Less than 2% of:** Organic Dextrose, Organic Citric Acid, Organic Flavor, Potassium Citrate, Salt, Sodium Citrate, Organic Stevia Leaf Extract, And Zinc Gluconate.

^{*} Mild to moderate dehydration.

NUTRITION INFORM	ATION	
	12 fl oz	1L
Calories	40	100
Protein, g	0	0
Carbohydrate, g	9	25
Sugars, g	9	25
Added Sugars, g	9	25
Fat, g	0	0
Sodium, mg	370	1030
Sodium, mEq	16.0	45
Potassium, mg	280	780
Potassium, mEq	7.1	20
Chloride, mg	440	1240
Chloride, mEq	12.4	35
Zinc, mg	2.8	7.8

Dosage

Refer to Administration Guide to restore fluid and minerals lost in diarrhea and vomiting. Pedialyte should be offered frequently in amounts tolerated. Total daily intake should be adjusted to meet individual needs, based on thirst and response to therapy. The suggested intakes for maintenance are based on water requirements for ordinary energy expenditure.*

Extrapolated from Barness LA, Curran JS: Nutrition, in Nelson WE (sr ed), Behrman RE, Kliegman RM, Arvin AM (eds): Nelson Textbook of Pediatrics, ed 15. Philadelphia: WB Saunders Co, 1996, pp 141-143.

ADMINISTRATION GUIDE FOR INFANTS AND YOUNG CHILDREN												
	Age	2	3	6	9	1	11/2	2	21/2	3	31/2	4
Weeks Months									Years			
Approximate	lb	9	14	18	21	23	26	28	30	32	34	36
Weight ¹	kg	4.0	6.4	8.2	9.5	10.5	11.8	12.7	13.6	14.4	15.3	16.3
Pedialyte for	fl oz/day	16	30	36	39	42	47	48	51	53	54	55
Maintenance ²		to	to	to	to	to	to	to	to	to	to	to
		20	34	42	45	47	52	53	56	57	57	59

Administration Guide does not apply to infants younger than 1 week of age. For children older than 4 years of age, maintenance intakes may exceed 2 liters daily. If there is vomiting or fever, or if diarrhea continues beyond 24 hours, consult the child's physician.

- Weight based on the 50th percentile of weight for age for boys from the National Center for Health Statistics (NCHS) Centers for Disease Control and Prevention (CDC) growth charts. Kuczmarski RJ, Ogden CL, Grummer-Strawn LM, et al: CDC Growth Charts: United States. Data from Vital and Health Statistics of the Centers for Disease Control and Prevention/National Center for Health Statistics. Advance Data, no. 314, December 4, 2000.
- 2. Fluid intake is total fluid requirement from oral electrolyte solution, formula, or other fluids, but does not take into account ongoing stool losses. Fluid loss in the stool should be replaced by consumption of an extra amount of Pedialyte equal to stool losses, in addition to the fluid maintenance requirement in this Administration Guide. Pedialyte Freezer Pops are to be used with Pedialyte Oral Electrolyte Solution or other appropriate fluids to help prevent dehydration.

Pedialyte® Freezer Pops

Electrolyte Solution

A cool way to help prevent dehydration. Just freeze and eat.



Description/Indications

Pedialyte replenishes vital fluids lost due to diarrhea; to help prevent dehydration* in infants, children, and adults. Scientifically designed balance of electrolytes and glucose for fast rehydration.

Features

- #1 Pediatrician & Pharmacist Recommended Brand
- Medical-grade hydration
- Designed to promote fluid absorption more effectively than common household beverages
- Provides glucose to promote sodium and water absorption
- · Low osmolality (see table)
- · Kosher, Halal

Precautions

- Not for use for children under 1 year of age
- Use under medical supervision

Availability: Retail

Ingredients

Grape: Water, Anhydrous Dextrose. Less than 2% of: Citric Acid, Salt, Cellulose Gum, Potassium Citrate, Potassium Sorbate (Preservative), Sodium Benzoate (Preservative), Natural & Artificial Flavor, Sucralose, Acesulfame Potassium, Red 40, and Blue 1.

Blue Raspberry: Water, Anhydrous Dextrose. Less than 2% of: Citric Acid, Salt, Cellulose Gum, Potassium Citrate, Potassium Sorbate (Preservative), Sodium Benzoate (Preservative), Natural & Artificial Flavor, Sucralose, Acesulfame Potassium, and Blue 1.

Cherry: Water, Anhydrous Dextrose. Less than 2% of: Citric Acid, Salt, Cellulose Gum, Potassium Citrate, Potassium Sorbate (Preservative), Sodium Benzoate (Preservative), Natural & Artificial Flavor, Sucralose, Acesulfame Potassium, and Red 40.

Orange: Water, Anhydrous Dextrose. Less than 2% of: Citric Acid, Salt, Cellulose Gum, Potassium Citrate, Potassium Sorbate (Preservative), Sodium Benzoate (Preservative), Natural & Artificial Flavor, Sucralose, Acesulfame Potassium, Yellow 6, and Red 40.

^{*} Mild to moderate dehydration.

Variety	125 mL (2 pops)	1L	
Calories	15	100	
Protein, g	0	0	
Carbohydrate, g	3	25	
Sugars, g	3	24	
Added Sugars, g	3	24	
Sodium, mg	130	1040	
Sodium, mEq	5.6	45	
Potassium, mg	100	800	
Potassium, mEq	2.5	20	
Chloride, mg	160	1280	
Chloride, mEq	4.4	35	
Osmolality, mOsm/kg H ₂ O	270	270	

Dosage

Refer to Administration Guide to restore fluid and minerals lost in diarrhea and vomiting. Pedialyte should be offered frequently in amounts tolerated. Total daily intake should be adjusted to meet individual needs, based on thirst and response to therapy.

The suggested intakes for maintenance are based on water requirements for ordinary energy expenditure.*

Extrapolated from Barness LA, Curran JS: Nutrition, in Nelson WE (sr ed), Behrman RE, Kliegman RM, Arvin AM (eds): Nelson Textbook of Pediatrics, ed 15. Philadelphia: WB Saunders Co, 1996, pp 141-143

ADMINISTRATION GUIDE FOR YOUNG CHILDREN												
	Age	1	1 11/2 2 21/2 3 31/2 4									
		Years										
Approximate	lb	23 26 28 30 32 34 36										
Weight ¹	kg	10.5	11.8	12.7	13.6	14.4	15.3	16.3				
Pedialyte for	fl oz/day	42	47	48	51	53	54	55				
Maintenance ²		to	to	to	to	to	to	to				
		47	52	53	56	57	57	59				

For children older than 4 years of age, maintenance intakes may exceed 2 liters daily. If there is vomiting or fever, or if diarrhea continues beyond 24 hours, consult the child's physician.

- Weight based on the 50th percentile of weight for age for boys from the National Center for Health Statistics (NCHS) Centers for Disease Control and Prevention (CDC) growth charts. Kuczmarski RJ, Ogden CL, Grummer-Strawn LM, et al: CDC Growth Charts: United States. Data from Vital and Health Statistics of the Centers for Disease Control and Prevention/National Center for Health Statistics. Advance Data, no. 314, December 4, 2000.
- Fluid intake is total fluid requirement from oral electrolyte solution, formula, or other fluids, but does not take into account ongoing stool losses. Fluid loss in the stool should be replaced by consumption of an extra amount of Pedialyte equal to stool losses, in addition to the fluid maintenance requirement in this Administration Guide. Pedialyte Freezer Pops are to be used with Pedialyte Oral Electrolyte Solution or other appropriate fluids to help prevent dehydration.

Pedialyte AdvancedCare® Plus

Electrolyte Solution



Description/Indications

Pedialyte AdvancedCare Plus is formulated with 33% more electrolytes* to replenish more of the electrolytes lost due to vomiting, diarrhea, or extreme perspiration caused by excessive heat or prolonged exercise. It also has PreActiv® Prebiotics to promote digestive health.

Features

- #1 Pediatrician & Pharmacist Recommended Brand
- Designed to promote fluid absorption more effectively than common household beverages
- Provides glucose to promote sodium and water absorption
- Quickly replaces fluids and zinc†
- PreActiv[®] Prebiotics help promote digestive health
- Low osmolality (see table)
- Kid-approved taste to encourage compliance in children 1 year of age and older
- · Kosher, Halal
- * 60 mEq sodium electrolytes per liter vs 45 mEq in original Pedialyte. † Liquid only.

Precautions

- Consult your doctor for children younger than 1 year
- · Use under medical supervision

Availability: Retail

List No.
66641
66645
66855

0.6-oz (17-g) packets; 6/carton	List No.
Berry Frost	66969
Strawberry Freeze	66972
Orange Breeze	67426

Ingredients

Berry Frost: Water, Dextrose, Galactooligosaccharides; Less than 0.5% of: Salt, Citric Acid, Potassium Citrate, Natural and Artificial Flavors, Sodium Citrate, Sucralose, Acesulfame Potassium, Zinc Gluconate, and Blue 1. Contains milk ingredients.

Chilled Cherry Pomegranate: Water, Dextrose, Galactooligosaccharides; Less Than 0.5% Of: Salt, Citric Acid, Potassium Citrate, Sodium Citrate, Natural and Artificial Flavors, Sucralose, Acesulfame Potassium, Zinc Gluconate, Red 40, and Blue 1. Contains milk ingredients.

Iced Grape: Water, Dextrose, Galactooligosaccharides; **Less than0.5% of:** Salt, Citric Acid, Potassium Citrate, Sodium Citrate, Artificial Flavor, Sucralose, Acesulfame Potassium, Zinc Gluconate, Red 40, and Blue 1. **Contains milk ingredients.**

Berry Frost (Powder): Anhydrous Dextrose, Short-Chain Fructooligosaccharides, Citric Acid, Salt, Potassium Citrate, Sodium Citrate, Natural and Artificial Flavor; **Less than 2% of:** Calcium Silicate, Acesulfame Potassium, Sucralose, and Blue 1.

Strawberry Freeze (Powder): Anhydrous Dextrose, Short-Chain Fructooligosaccharides, Citric Acid, Salt, Potassium Citrate, Sodium Citrate, Natural Flavor; Less than 2% of: Calcium Silicate. Acesulfame Potassium, Sucralose. Red 40, and Blue 1.

Orange Breeze (Powder): Anhydrous Dextrose, Short-Chain Fructooligosaccharides, Citric Acid, Salt, Potassium Citrate, Sodium Citrate; Less than 2% of: Natural and Artificial Flavor, Calcium Silicate, Acesulfame Potassium, Sucralose, and Yellow 6.

Dosage

Refer to Administration Guide to restore fluid and minerals lost in diarrhea and vomiting. Pedialyte should be offered frequently in amounts tolerated. Total daily intake should be adjusted to meet individual needs, based on thirst and response to therapy. The suggested intakes for maintenance are based on water requirements for ordinary energy expenditure.*

Extrapolated from Barness LA, Curran JS: Nutrition, in Nelson WE (sr ed), Behrman RE, Kliegman RM, Avrin AM (eds): Nelson Textbook of Pediatrics, ed 15. Philadelphia: WB Saunders Co. 1996. pp 141-143.

ADMINISTRATION GUIDE FOR INFANTS AND YOUNG CHILDREN									
	Age	1	11/2	2	21/2	3	31/2	4	
					Years				
Approximate	lb	23	26	28	30	32	34	36	
Weight ¹	kg	10.5	11.8	12.7	13.6	14.4	15.3	16.3	
Pedialyte for	fl oz/day	42	47	48	51	53	54	55	
Maintenance ²		to	to	to	to	to	to	to	
		47	52	53	56	57	57	59	

Berry Frost	1 L Solution	16 fl oz Powder (Reconstituted)
Calories	100	50
Protein, g	0	0
Carbohydrate, g	28	12
Dietary Fiber, g	2	1
Total Sugars, g	19	12
Added Sugars, g	18	12
Fat, g	0	0
Sodium, mg	1380	650
Sodium, mEq	60	28
Potassium, mg	780	370
Potassium, mEq	20	9
Chloride, mg	1770	840
Chloride, mEq	50	24
Zinc, mg	7.8	0
Osmolality, mOsm/kg H ₂ O	265	265

† Pedialyte AdvancedCare Plus liters may have small differences for the amounts of carbohydrates or sugars within different flavors.

For children older than 4 years of age, maintenance intakes may exceed 2 liters daily. If there is vomiting or fever, or if diarrhea continues beyond 24 hours, consult the child's physician.

- Weight based on the 50th percentile of weight for age for boys from the National Center for Health Statistics (NCHS) Centers for Disease Control and Prevention (CDC) growth charts. Kuczmarski RJ, Ogden CL, Grummer-Strawn LM, et al.: CDC Growth Charts: United States. Data from Vital and Health Statistics of the Centers for Disease Control and Prevention/National Center for Health Statistics. Advance Data, no. 314, December 4, 2000.
- 2. Fluid intake is total fluid requirement from oral electrolyte solution, formula, or other fluids, but does not take into account ongoing stool losses. Fluid loss in the stool should be replaced by consumption of an extra amount of Pedialyte equal to stool losses, in addition to the fluid maintenance requirement in this Administration Guide. Pedialyte Freezer Pops are to be used with Pedialyte Oral Electrolyte Solution or other appropriate fluids to help prevent dehydration.





Calcilo XD⁶

Low-Calcium (2.9 mg/100 Cal)/ Vitamin D-Free Infant Formula With Iron



Description/Indications

Nutrition support of infants with hypercalcemia, as may occur in infants with Williams syndrome, osteopetrosis, and primary neonatal hyperparathyroidism, and when a low-calcium/vitamin D-free formula is needed.

Features

- The only commercially available formula for the management of hypercalcemia in infants
- No vitamin D
- Only a minimal amount of calcium (15 mg/100 g powder; 2.9 mg/100 Cal)
- Nutritionally complete—when prepared as directed, meets the American Academy of Pediatrics Committee on Nutrition recommendations for all nutrients including vitamins and minerals except calcium, vitamin D, and phosphorus
- L-carnitine (7 mg/100 g) and taurine (35 mg/100 g)
- Eliminates the need to prepare a modular formula
- · Lactose-free and gluten-free
- Kosher, Halal

Precautions

 Calcium, vitamin D, and phosphorus requirements should be determined by appropriate laboratory tests

Availability: Hospital/Institutional

Size	Container	List No.
12.4 oz (352 g)	can; 6/case	53328

Preparation

To Make	Water	Scoop(s) of Powder
(Approx)	(fl oz)	(Use scoop in can)
2 fl oz	2	. 1 unpacked, level (8.6 g)
4 fl oz	4	2 unpacked, level
6 fl oz	6	3 unpacked, level

To make a larger amount of formula at standard density (20 Cal/ fl oz) using a standard measuring cup, add 1 unpacked, level cup (NOT the enclosed scoop) of powder (105 g) to 24 fl oz of water. Yields approximately 26 fl oz of formula. To maintain freshness, pour prepared formula into individual feeding bottles, cap and store in refrigerator. Prepared formula should not be left unrefrigerated.

When mixed as directed, the contents of one can (352 g) will make approximately 90 fl oz of formula.

Household	Weight
Measure (US)	(Metric)
1 Tablespoon	7 g
1/4 Cup	26 g
1/3 Cup	35 g
1/2 Cup	53 g
1 Cup	105 g

Powder: Corn Syrup, Coconut Oil, Corn Oil, Whey Protein Concentrate, Sodium Caseinate; Less than 1% of: Potassium Phosphate, Magnesium Chloride, Potassium Chloride, Potassium Citrate, Ascorbic Acid, Inositol, Salt, Choline Chloride, Ferrous Sulfate, Potassium Hydroxide, Taurine, Zinc Sulfate, L-Carnitine, d-Alpha-Tocopheryl Acetate, Niacinamide, Calcium Pantothenate, Vitamin A Palmitate, Copper Sulfate, Thiamine Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Manganese Sulfate, Potassium Iodide, Phylloquinone, Biotin, Sodium Selenate, and Vitamin B.,. Contains milk ingredients.

NUTRITION INF	ORMATION				
	100 g powder F	er 100 Cal (5 fl oz as prepared)		100 g powder	Per 100 Cal (5 fl oz as prepared)
Energy, Cal	513	100	Biotin, mcg	23	4.5
Protein, g	11.4	2.2	Vitamin C, mg	46	9
% Total Calories	9	9	Choline, mg	62	12
Source	Whey protein concentrate, sodium ca	seinate	Inositol, mg	123	24
Fat, g	28.7	5.6	Minerals		
% Total Calories	50	50	Calcium, mg	15	2.9
Source	Coconut and corn oils		Phosphorus, mg	128	25
Linoleic Acid, mg	6660	1300	Magnesium, mg	31	6
Carbohydrate, g	52.3	10.2	Iron, mg	9.2	1.8
% Total Calories	41	41	Zinc, mg	3.8	0.8
Source	Corn syrup		Manganese, mcg	26	5
Vitamins			Copper, mcg	460	90
Vitamin A, IU	1540	300	lodine, mcg	31	6
Vitamin D, IU	0	0	Selenium, mcg	10	2
Vitamin E, IU	10.0	1.9	Sodium, mg	125	24
Vitamin K, mcg	41	8	Potassium, mg	420	82
Thiamin (Vit B ₁), mcg	513	100	Chloride, mg	292	57
Riboflavin (Vit B ₂), mcg	770	150	Other Characteristics		
Vitamin B _s , mcg	310	60	PRSL, mOsm	94	18
Vitamin B ₁₂ , mcg	1.3	0.3	Water, g	2.5	134
Niacin, mcg	5400	1050	Osmolality, mOsm/kg H₂O	-	190
Folic Acid (Folacin), mcg	77	15			
Pantothenic Acid, mcg	2300	450			

Cyclinex®-1

Amino Acid-Modified Infant Formula With Iron



Description/Indications

Nutrition support of infants and toddlers with a urea cycle disorder, gyrate atrophy of the choroid and retina, or HHH syndrome.

Features

- Nonessential amino acid-free to decrease the ingestion of waste nitrogen
- L-carnitine (190 mg/100 g) and taurine (40 mg/100 g) to help supply amounts normally found in human milk and foods of animal origin
- Approximately 43% of energy as fat to help achieve acceptable formula osmolality
- 6.9% of energy as linoleic acid
- Nutrient profile specifically designed for infants and toddlers
- When fed according to the Abbott Support Protocols, provides adequate amounts of all nutrients
- Additional vitamins and minerals to compensate for nutrient losses due to nitrogen-scavenger medications
- · Lactose-free and gluten-free
- Halal
- Non-GMO*
- Includes DHA/ARA

Precautions

- Must be supplemented with protein and fluid in prescribed amounts to completely meet protein and water requirements
- · Do not boil mixture or use terminal sterilization

Availability: Hospital/Institutional

Size	Container	List No.
14.1 oz (400 g)	can; 6/case	67032

Preparation

Follow physician's instructions carefully.

Household Measure (US)	Weight (Metric)
1 Tablespoon	8 g
1/4 Cup	
1/3 Cup	40 g
1/2 Cup	60 g
1 Cup	120 g

^{*} Ingredients not genetically engineered.

Powder: Corn Syrup Solids, High Oleic Safflower Oil, Coconut Oil, Soy Oil, Modified Corn Starch, L-Leucine. Less than 2% of: L-Lysine, Calcium Phosphate, L-Valine, L-Isoleucine, Potassium Phosphate, L-Tyrosine, L-Threonine, L-Phenylalanine, Sodium Citrate, Potassium Chloride, L-Cystine Dihydrochloride, L-Histidine, L-Methionine, M. Alpina Oil, L-Tryptophan, C. Cohnii Oil, Calcium Carbonate, Citric Ackorchic Acsorbic Acid, L-Carnitine, Inositol, Choline Chloride, Taurine, Ferrous Sulfate, Ascorbyl Palmitate, Zinc Sulfate, Alpha-Tocopherols, Calcium Pantothenate, Niacinamide, Manganese Sulfate, Thiamine Hydrochloride, Copper Sulfate, Vitamin A Palmitate, Riboflavin, Pyridoxine Hydrochloride, Potassium Iodide, Folic Acid, Phylloquinone, Chromium Chloride, Sodium Selenate, Sodium Molybdate, Biotin, Vitamin D.,, ultamin B.,, and Salt.

	100 g powder		100 g powder
Energy, Cal	510	Biotin, mcg	26.4
Protein Equivalent, g	7.5	Vitamin C, mg	75
Source	L-amino acids	Choline, mg	106
L-Carnitine, mg	190	Inositol, mg	160
Fat, g	24.6	Minerals	
Source	High-Oleic Safflower Oil, Coconut Oil, Soy Oil	Calcium, mg	650
Linoleic Acid, mg	4000	Phosphorus, mg	455
Linolenic Acid, mg	410	Magnesium, mg	55
DHA, mg	102	Iron, mg	8.75
ARA, mg	124	Zinc, mg	5.8
Carbohydrate, g	57	Manganese, mcg	425
Source	Corn syrup solids	Copper, mcg	450
Vitamins		lodine, mcg	70
Vitamin A, IU	1450	Selenium, mcg	23.8
Vitamin D, IU	510	Chromium, mcg	14
Vitamin E, IU	14	Molybdenum, mcg	15
Vitamin K, mcg	70	Sodium, mg	215
Thiamin (Vit B ₁), mcg	1235	Potassium, mg	760
Riboflavin (Vit B ₂), mcg	725	Chloride, mg	390
Vitamin B ₆ , mcg	625	Other Characteristics	
Vitamin B ₁₂ , mcg	0.92	PRSL, mOsm	98
Niacin, mcg	4840	Osmolality, mOsm/kg H₂O	275 (20 Cal/fl oz)
Folic Acid, mcg	97		
Pantothenic Acid, mcg	4580		

C. cohnii oil is a source of DHA M. alpina oil is a source of ARA

Cyclinex®-2

Amino Acid-Modified Medical Food



Description/Indications

Nutrition support of children and adults with a urea cycle disorder, gyrate atrophy of the choroid and retina, or HHH syndrome.

Features

- Nonessential amino acid-free to decrease the ingestion of waste nitrogen
- L-carnitine (370 mg/100 g) and taurine (140 mg/100 g) to help supply amounts normally found in foods of animal origin
- Approximately 35% of energy as fat to help supply essential fatty acids
- Nutrient profile specifically designed for children and adults
- When fed according to the Abbott Support Protocols, provides adequate amounts of all nutrients
- Additional vitamins and minerals to compensate for nutrient losses due to nitrogen-scavenger medications
- · Lactose-free and gluten-free
- Halal
- Non-GMO*
- Includes DHA

Precautions

- · Do not heat or use in cooking
- Must be supplemented with protein in prescribed amounts to completely meet protein requirements

Availability: Hospital/Institutional

Size	Container	List No.
14.1	oz (400 g) can; 6/case	67034

Preparation

Follow physician's instructions carefully.

Household	Weight
Measure (US)	(Metric)
1 Tablespoon	8 g
1/4 Cup	30 g
1/3 Cup	40 g
1/2 Cup	60 g
1 Cup	120 g

^{*} Ingredients not genetically engineered.

Powder: Com Syrup Solids, High Oleic Safflower Oil, Coconut Oil, Sodium Citrate, Soy Oil, L-Leucine, L-Lysine Acetate, Calcium Phosphate, L-Valine, L-Isoleucine, Magnesium Phosphate, Potassium Chloride; Less than 2% of: L-Tyrosine, L-Phenylalanine, L-Threonine, Silicon Dioxide, Modified Corn Starch, Potassium Citrate, Potassium Phosphate, L-Oystine Dihydrochloride, L-Histidine, L-Methionine, C. Cohnii Oil, L-Tryptophan, Calcium Citrate, Calcium Carbonate, L-Carnitine, Citric Acid, Ascorbic Acid, Taurine, Choline Chloride, Inositol, Ferrous Sulfate, Silos Sulfate, Ascorbyl Palmitate, Niacinamide, Calcium Pantothenate, DI-Alpha-Tocopheryl Acetate, Mixed Tocopherols, Copper Sulfate, Thiamine Hydrochloride, Manganese Sulfate, Ribofavin, Pyridoxine Hydrochloride, Vitamin A Palmitate, Folic Acid, Biotin, Chromium Chloride, Potassium lodide, Sodium Selenate, Phylloquinone, Sodium Molybdate, Vitamin D₃, And Vitamin B₃.

	100 g powder		100 g powder
Energy, Cal	440	Vitamin C, mg	75
Protein equivalent, g	15	Choline, mg	130
Source	L-amino acids	Inositol, mg	100
L-Carnitine, mg	370	Minerals	
Fat, g	17	Calcium, mg	1250
Source	High-Oleic Safflower Oil, Coconut Oil, Soy Oil	Phosphorus, mg	1020
Linoleic Acid, mg	2800	Magnesium, mg	300
Linolenic Acid, mg	275	Iron, mg	17
DHA, mg	192	Zinc, mg	17
Carbohydrate, g	45	Manganese, mg	0.9
Source	Corn syrup solids	Copper, mg	2
Vitamins		lodine, mcg	150
Vitamin A, IU	1650	Selenium, mcg	40
Vitamin D, IU	1100	Chromium, mcg	37
Vitamin E, IU	13	Molybdenum, mcg	40
Vitamin K, mcg	70	Sodium, mg	1175
Thiamin (Vit B₁), mg	4	Potassium, mg	1800
Riboflavin (Vit B ₂), mg	1.8	Chloride, mg	1325
Vitamin B ₆ , mg	1.3	Other Characteristics	
Vitamin B ₁₂ , mcg	7.3	PRSL, mOsm	253
Niacin, mg	21	Osmolality, mOsm/kg H₂O	1015 (30 Cal/fl oz)
Folic Acid, mcg	510		
Pantothenic Acid, mg	10.9		
Biotin, mcg	140		

Glutarex®-1

Amino Acid-Modified Infant Formula With Iron



Description/Indications

Nutrition support of infants and toddlers with glutaric aciduria type I.

Features

- Lysine- and tryptophan-free to allow greater intake of intact protein
- L-carnitine (900 mg/100 g) to help supply amount normally found in human milk and foods of animal origin and to help excrete toxic metabolites
- Taurine (40 mg/100 g) to help supply amount normally found in human milk and foods of animal origin
- Approximately 40% of energy as fat to help achieve acceptable formula osmolality
- . 6.6% of energy as linoleic acid
- Nutrient profile specifically designed for infants and toddlers
- When fed according to the Abbott Support Protocols, provides adequate amounts of all nutrients
- · Lactose-free and gluten-free
- Halal
- Non-GMO*
- Includes DHA/ARA

Precautions

- Must be supplemented with protein and fluid in prescribed amounts to completely meet lysine, tryptophan, and water requirements
- Do not boil mixture or use terminal sterilization

Availability: Hospital/Institutional

Size	Container	List No.
14.1 oz (400 g)	can; 6/case	67036

Preparation

Follow physician's instructions carefully.

Weight
(Metric)
8 g
30 g
40 g
60 g
120 g

^{*} Ingredients not genetically engineered.

Powder: Corn Syrup Solids, High Oleic Safflower Oil, Coconut Oil, Soy Oil, L-Glutamic Acid, Modified Corn Starch. Less than 2% of: L-Leucine, L-Proline, L-Aspartic Acid, L-Arginine, Calcium Phosphate, L-Valine, L-Isoleucine, Potassium Phosphate, Glycine, L-Carnitine, L-Tyrosine, L-Phenylalanine, L-Serine, L-Threonine, Sodium Citrate, Magnesium Chloride, L-Histidine, L-Methionine, M. Alpina Oil, C. Cohnii Oil, Calcium Carbonate, Citric Acid, Ascorbic Acid, L-Cystine Dihydrochloride, Inositol, Choline Chloride, Taurine, Ferrous Sulfate, Ascorbyl Palmitate, Zinc Sulfate, dl-Alpha-Tocopheryl Acetate, Mixed Tocopherols, Calcium Pantothenate, Niacinamide, Manganese Sulfate, Thiamine Hydrochloride, Copper Sulfate, Vitamin A Palmitate, Riboflavin, Pyridoxine Hydrochloride, Potassium Iodide, Folic Acid, Phylloquinone, Chromium Chloride, Sodium Selenate, Sodium Molybdate, Biotin, Vitamin B₁₀, and Salt.

	100 g powder		100 g powder
Energy, Cal	480	Biotin, mcg	23.5
Protein equivalent, g	15	Vitamin C, mg	75
Source	L-amino acids	Choline, mg	95
L-Carnitine, mg	900	Inositol, mg	144
Fat, g	21.7	Minerals	
Source	High-Oleic Safflower Oil, Coconut Oil, Soy Oil	Calcium, mg	575
Linoleic Acid, mg	3650	Phosphorus, mg	400
Linolenic Acid, mg	360	Magnesium, mg	50
DHA, mg	96	Iron, mg	7.6
ARA, mg	115	Zinc, mg	5.2
Carbohydrate, g	53	Manganese, mcg	385
Source	Corn syrup solids	Copper, mcg	408
Vitamins		lodine, mcg	65
Vitamin A, IU	1300	Selenium, mcg	21
Vitamin D, IU	450	Chromium, mcg	12
Vitamin E, IU	12.5	Molybdenum, mcg	13.8
Vitamin K, mcg	60	Sodium, mg	190
Thiamin (Vit B,), mcg	1120	Potassium, mg	675
Riboflavin (Vit B ₂), mcg	650	Chloride, mg	325
Vitamin B ₆ , mcg	560	Other Characteristics	
Vitamin B ₁₂ , mcg	0.85	PRSL, mOsm	133
Niacin, mcg	4450	Osmolality, mOsm/kg H ₂ O	385 (20 Cal/fl oz)
Folic Acid, mcg	88		
Pantothenic Acid, mcg	4000		

C. cohnii oil is a source of DHA M. alpina oil is a source of ARA

Glutarex®-2

Amino Acid-Modified Medical Food



Description/Indications

Nutrition support of children and adults with glutaric aciduria type I.

Features

- Lysine- and tryptophan-free to allow greater intake of intact protein
- L-carnitine (1800 mg/100 g) to help supply amount normally found in foods of animal origin and to help excrete toxic metabolites
- Taurine (125 mg/100 g) to help supply amount normally found in foods of animal origin
- Approximately 30% of energy as fat to help supply essential fatty acids
- Nutrient profile specifically designed for children and adults
- When fed according to the Abbott Support Protocols, provides adequate amounts of all nutrients
- · Lactose-free and gluten-free
- Halal
- Non-GMO*
- Includes DHA

Precautions

- · Do not heat or use in cooking
- Must be supplemented with protein and fluid in prescribed amounts to completely meet lysine, tryptophan, and water requirements

Availability: Hospital/Institutional

ize	Container	List No.
4.1 oz (400 g)	can; 6/case	67038

Preparation

Follow physician's instructions carefully.

Household	Weight
Measure (US)	(Metric)
1 Tablespoon	8 g
1/4 Cup	30 g
1/3 Cup	40 g
1/2 Cup	60 g
1 Cup	120 g

^{*} Ingredients not genetically engineered.

Powder: Corn Syrup Solids, L-Glutamic Acid, High Oleic Safflower Oil, Coconut Oil, Sodium Citrate, Soy Oil, L-Leucine, L-Proline, L-Aspartic Acid, L-Arginine, L-Valine, L-Alanine, Calcium Phosphate, L-Isoleucine, Glycine, L-Carnitine, L-Tyrosine; Less Than 2% of: L-Phenylalanine, Magnesium Phosphate, Potassium Chloride, L-Serine, L-Threonine, Silicon Dioxide, Modified Corn Starch, L-Histidine, Potassium Citrate, Potassium Phosphate, L-Methionine, C. Cohnii Oil, L-Cystine Dihydrochloride, Calcium Citrate, Calcium Carbonate, Citric Acid, Ascorbic Acid, Taurine, Choline Chloride, Inositol, Ferrous Sulfate, Zinc Sulfate, Ascorbyl Palmitate, Niacinamide, Calcium Pantothenate, DI-Alpha-Tocopheryl Acetate, Mixed Tocopherols, Copper Sulfate, Thiamine Hydrochloride, Manganese Sulfate, Riboflavin, Pyridoxine Hydrochloride, Vitamin A Palmitate, Folic Acid, Biotin, Chromium Chloride, Potassium Iodide, Sodium Selenate, Phylloquinone, Sodium Molybdate, Vitamin D., And Vitamin B.,..

	100 g powder		100 g powder
Energy, Cal	410	Vitamin C, mg	60
Protein equivalent, g	30	Choline, mg	100
Source	L-amino acids	Inositol, mg	82
L-Carnitine, mg	1800	Minerals	
Fat, g	13	Calcium, mg	975
Source	High-Oleic Safflower Oil, Coconut Oil, Soy Oil	Phosphorus, mg	760
Linoleic Acid, mg	2200	Magnesium, mg	225
Linolenic Acid, mg	225	Iron, mg	13
DHA, mg	180	Zinc, mg	13
Carbohydrate, g	35	Manganese, mg	0.8
Source	Corn syrup solids	Copper, mg	1.5
Vitamins		lodine, mcg	100
Vitamin A, IU	1500	Selenium, mcg	35
Vitamin D, IU	900	Chromium, mcg	27
Vitamin E, IU	12	Molybdenum, mcg	30
Vitamin K, mcg	60	Sodium, mg	880
Thiamin (Vit B,), mg	3.3	Potassium, mg	1370
Riboflavin (Vit B ₂), mg	1.6	Chloride, mg	940
Vitamin B, mg	1.1	Other Characteristics	
Vitamin B ₁₉ , mcg	5	PRSL, mOsm	296
Niacin, mg	16	Osmolality, mOsm/kg H ₂ O	1360 (30 Cal/fl oz)
Folic Acid, mcg	425		
Pantothenic Acid, mg	8		
Biotin, mcg	100		

Hominex®-1

Amino Acid-Modified Infant Formula With Iron



Description/Indications

Nutrition support of infants and toddlers with vitamin B_e -nonresponsive homocystinuria or hypermethioninemia.

Features

- Methionine-free to allow greater intake of intact protein
- L-cystine, in a soluble form, an essential amino acid that is often deficient in diets of infants and toddlers with homocystinuria
- L-carnitine (20 mg/100 g) and taurine (40 mg/100 g) to help supply amounts normally found in human milk and foods of animal origin
- Approximately 40% of energy as fat to help achieve acceptable formula osmolality
- 6.6% of energy as linoleic acid
- Nutrient profile specifically designed for infants and toddlers
- When fed according to the Abbott Support Protocols, provides adequate amounts of all nutrients
- · Lactose-free and gluten-free
- Halal
- Non-GMO*
- Includes DHA/ARA

Precautions

- Must be supplemented with protein and fluid in prescribed amounts to completely meet methionine and water requirements
- Do not boil mixture or use terminal sterilization

Availability: Hospital/Institutional

Size	Container	List No.
14.1 oz	(400 g) can; 6/case	67040

Preparation

Follow physician's instructions carefully.

Household	Weight
Measure (US)	(Metric)
1 Tablespoon	8 g
1/4 Cup	30 g
1/3 Cup	40 g
1/2 Cup	60 g
1 Cup	120 g

^{*} Ingredients not genetically engineered.

Powder: Corn Syrup Solids, High Oleic Safflower Oil, Coconut Oil, Soy Oil, Modified Corn Starch. Less than 2% of: L-Leucine, L-Lysine, Calcium Phosphate, L-Proline, L-Glutamine, L-Valine, L-Ispecine, L-Arginine, Potassium Citrate, Datassium Phosphate, Glycine, L-Alanine, L-Tyrosine, L-Phenylalanine, L-Asparagine, L-Serine, L-Threonine, Sodium Citrate, L-Cystine Dihydrochloride, Potassium Citrate, Magnesium Chloride, L-Histidine, M. Alpina Oil, C. Cohnii Oil, Calcium Carbonate, Citric Acid, Ascorbic Acid, L-Glutamic Acid, L-Tryptophan, Inositol, Choline Chloride, L-Aspartic Acid, Taurine, Ferrous Sulfate, Ascorbyl Palmitate, L-Carnitine, Zinc Sulfate, dl-Alpha-Tocopheryl Acetate, Mixed Tocopherols, Calcium Pantothenate, Niacinamide, Manganese Sulfate, Thiamine Hydrochloride, Copper Sulfate, Vitamin A Palmitate, Riboflavin, Pyridoxine Hydrochloride, Potassium Iodide, Folic Acid, Phylloquinone, Chromium Chloride, Sodium Selenate, Sodium Molybdate, Biotin, Vitamin D., Vitamin B., and Salt.

	100 g powder		100 g powder
Energy, Cal	480	Biotin, mcg	23.5
Protein equivalent, g	15.0	Vitamin C, mg	75
Source	L-amino acids	Choline, mg	95
L-Carnitine, mg	20	Inositol, mg	144
Fat, g	21.7	Minerals	
Source	High-oleic safflower, coconut, and soy oils	Calcium, mg	575
Linoleic Acid, mg	3650	Phosphorus, mg	400
Linolenic Acid, mg	360	Magnesium, mg	50
DHA, mg	96	Iron, mg	7.6
ARA, mg	115	Zinc, mg	5.2
Carbohydrate, g	53.0	Manganese, mcg	385
Source	Corn syrup solids	Copper, mcg	408
Vitamins		lodine, mcg	65
Vitamin A, IU	1300	Selenium, mcg	21
Vitamin D, IU	450	Chromium, mcg	12
Vitamin E, IU	12.5	Molybdenum, mcg	13.8
Vitamin K, mcg	60	Sodium, mg	190
Thiamin (Vit B,), mcg	1120	Potassium, mg	675
Riboflavin (Vit B _a), mcg	650	Chloride, mg	410
Vitamin B, mcg	560	Other Characteristics	
Vitamin B,, mcg	0.85	PRSL, mOsm	136
Niacin, mcg	4450	Osmolality, mOsm/kg H ₂ O	375 (20 Cal/fl oz)
Folic Acid, mcg	88		
Pantothenic Acid, mcg	4000		

C. cohnii oil is a source of DHA M. alpina oil is a source of ARA

Hominex®-2

Amino Acid-Modified Medical Food



Description/Indications

Nutrition support of children and adults with vitamin B_a -nonresponsive homocystinuria or hypermethioninemia.

Features

- Methionine-free to allow greater intake of intact protein
- L-cystine, in a soluble form, an essential amino acid that is often deficient in diets of children and adults with homocystinuria
- L-carnitine (40 mg/100 g) and taurine (125 mg/100 g) to help supply amounts normally found in foods of animal origin
- Approximately 30% of energy as fat to help supply essential fatty acids
- Nutrient profile specifically designed for children and adults
- When fed according to the Abbott Support Protocols, provides adequate amounts of all nutrients
- · Lactose-free and gluten-free
- Halal
- Non-GMO*
- Includes DHA

Precautions

- · Do not heat or use in cooking
- Must be supplemented with protein and fluid in prescribed amounts to completely meet methionine and water requirements

Availability: Hospital/Institutional

Size		Container		List No.
14.1	oz (400 g)	can; 6/case	9	67042

Preparation

Follow physician's instructions carefully.

mode on one mode of the post do.	
Household	Weight
Measure (US)	(Metric)
1 Tablespoon	8 g
1/4 Cup	30 g
1/3 Cup	40 g
1/2 Cup	60 g
1 Cup	120 g

^{*} Ingredients not genetically engineered.

Powder: Corn Syrup Solids, High Oleic Safflower Oil, Coconut Oil, Sodium Citrate, Soy Oil, L-Leucine, L-Lysine Acetate, L-Proline, L-Glutamine, L-Valine, Calcium Phosphate, L-Isoleucine, L-Arginine, Glycine, L-Alanine, L-Tyrosine; Less than 2% of: Magnesium Phosphate, L-Phenylalanine, Potassium Chloride, Asparagine Monohydrate, L-Serine, L-Thropnine, L-Cystine Dihydrochloride, Silicon Dioxide, Modified Corn Starch, L-Histidine, Potassium Citrate, Potassium Phosphate, C. Cohnii Oil, L-Glutamic Acid, Calcium Citrate, L-Tryptophan, Calcium Carbonate, Citric Acid, L-Aspartic Acid, Ascorbic Acid, Taurine, Cholinde, Inositol, Ferrous Sulfate, Zinc Sulfate, L-Carnitine, Ascorbly Palmitate, Niacinamide, Calcium Pantothenate, DI-Alpha-Tocopheryl Acetate, Mixed Tocopherols, Copper Sulfate, Thiamine Hydrochloride, Manganese Sulfate, Riboflavin, Pyridoxine Hydrochloride, Vitamin A Palmitate, Folic Acid, Biotin, Chromium Chloride, Potassium Iodide, Sodium Selenate, Phylloquinone, Sodium Molybdate, Vitamin D., And Vitamin B.,

	100 g powder		100 g powder
Energy, Cal	410	Vitamin C, mg	60
Protein equivalent, g	30	Choline, mg	100
Source	L-amino acids	Inositol, mg	82
L-Carnitine, mg	40	Minerals	
Fat, g	14	Calcium, mg	975
Source	High-Oleic Safflower Oil, Coconut Oil, Soy Oil	Phosphorus, mg	760
Linoleic Acid, mg	2200	Magnesium, mg	225
Linolenic Acid, mg	225	Iron, mg	13
DHA, mg	180	Zinc, mg	13
Carbohydrate, g	35	Manganese, mg	0.8
Source	Corn syrup solids	Copper, mg	1.5
Vitamins		lodine, mcg	100
Vitamin A, IU	1500	Selenium, mcg	35
Vitamin D, IU	900	Chromium, mcg	27
Vitamin E, IU	12	Molybdenum, mcg	30
Vitamin K, mcg	60	Sodium, mg	880
Thiamin (Vit B,), mg	3.3	Potassium, mg	1370
Riboflavin (Vit B ₂), mg	1.6	Chloride, mg	1160
Vitamin B, mg	1.1	Other Characteristics	
Vitamin B ₁₂ , mcg	5	PRSL, mOsm	302
Niacin, mg	16	Osmolality, mOsm/kg H ₂ O	1350 (30 Cal/fl oz)
Folic Acid, mcg	425		
Pantothenic Acid, mg	8		
Biotin, mcg	100		

I-Valex®-1

Amino Acid-Modified Infant Formula With Iron



Description/Indications

Nutrition support of infants and toddlers with a disorder of leucine catabolism.

Features

- · Leucine-free to allow greater intake of intact protein
- Isoleucine and valine to help prevent deficiency of these amino acids
- L-carnitine (900 mg/100 g) to help supply amount normally found in human milk and foods of animal origin and to help excrete toxic metabolites
- Glycine (1000 mg/100 g) to bind with acyl groups and help enhance their excretion in a nontoxic form
- Approximately 40% of energy as fat to help achieve acceptable formula osmolality
- . 6.6% of energy as linoleic acid
- Nutrient profile specifically designed for infants and toddlers
- When fed according to the Abbott Support Protocols, provides adequate amounts of all nutrients
- · Lactose-free and gluten-free
- Halal
- Non-GMO*
- Includes DHA/ARA

Precautions

- Must be supplemented with protein and fluid in prescribed amounts to completely meet leucine and water requirements
- Do not boil mixture or use terminal sterilization

Availability: Hospital/Institutional

Size	Container	List No.
14.1 oz (400 g)	can; 6/case	67044

Preparation

Follow physician's instructions carefully.

Household	Weight
Measure (US)	(Metric)
1 Tablespoon	8 g
1/4 Cup	30 g
1/3 Cup	40 g
1/2 Cup	60 g
1 Cup	120 g

^{*} Ingredients not genetically engineered.

Powder: Corn Syrup Solids, High Oleic Safflower Oil, Coconut Oil, Soy Oil, L-Alanine, L-Glutamic Acid, Modified Corn Starch. Less than 2% of: L-Aspartic Acid, L-Proline, L-Lysine, Calcium Phosphate, L-Arginine, Potassium Phosphate, Glycine, L-Carnitine, L-Tyrosine, L-Phenylalanine, L-Serine, L-Threonine, Sodium Citrate, Potassium Citrate, L-Valine, Magnesium Chloride, L-Isoleucine, L-Histidine, L-Methionine, M. Alpina Oil, C. Cohnii Oil, Calcium Carbonate, Citric Acid, Ascorbic Acid, L-Cystine Dihydrochloride, L-Tryptophan, Inositol, Choline Chloride, Taurine, Ferrous Sulfate, Ascorbyl Palmitate, Zinc Sulfate, dl-Alpha-Tocopheryl Acetate, Mixed Tocopherols, Calcium Pantothenate, Niacinamide, Manganese Sulfate, Thiamine Hydrochloride, Copper Sulfate, Vitamin A Palmitate, Riboflavin, Pyridoxine Hydrochloride, Potassium Iodide, Folic Acid, Phylloquinone, Chromium Chloride, Sodium Selenate, Sodium Molybdate, Biotin, Vitamin D., Vitamin B., and Salt.

	400		400
	100 g powder		100 g powder
Energy, Cal	480	Biotin, mcg	23.5
Protein equivalent, g	15	Vitamin C, mg	75
Source	L-amino acids	Choline, mg	95
L-Carnitine, mg	900	Inositol, mg	144
Fat, g	21.7	Minerals	
Source	High-Oleic Safflower Oil, Coconut Oil, Soy Oil	Calcium, mg	575
Linoleic Acid, mg	3650	Phosphorus, mg	400
Linolenic Acid, mg	360	Magnesium, mg	50
DHA, mg	96	Iron, mg	7.6
ARA, mg	115	Zinc, mg	5.2
Carbohydrate, g	53	Manganese, mcg	385
Source	Corn syrup solids	Copper, mcg	408
Vitamins		lodine, mcg	65
Vitamin A, IU	1300	Selenium, mcg	21
Vitamin D, IU	450	Chromium, mcg	12
Vitamin E, IU	12.5	Molybdenum, mcg	13.8
Vitamin K, mcg	60	Sodium, mg	190
Thiamin (Vit B ₁), mcg	1120	Potassium, mg	675
Riboflavin (Vit B ₂), mcg	650	Chloride, mg	325
Vitamin B ₆ , mcg	560	Other Characteristics	
Vitamin B ₁₂ , mcg	0.85	PRSL, mOsm	133
Niacin, mcg	4450	Osmolality, mOsm/kg H ₂ O	375 (20 Cal/fl oz)
Folic Acid, mcg	88		
Pantothenic Acid, mcg	4000		

C. cohnii oil is a source of DHA M. alpina oil is a source of ARA

I-Valex®-2

Amino Acid-Modified Medical Food



Description/Indications

Nutrition support of children and adults with a disorder of leucine catabolism.

Features

- · Leucine-free to allow greater intake of intact protein
- Isoleucine and valine to help prevent deficiency of these amino acids
- L-carnitine (1800 mg/100 g) to help supply amount normally found in foods of animal origin and to help excrete toxic metabolites
- Glycine (2000 mg/100 g) to bind with acyl groups and help enhance their excretion in a nontoxic form
- Approximately 30% of energy as fat to help supply essential fatty acids
- Nutrient profile specifically designed for children and adults
- When fed according to the Abbott Support Protocols, provides adequate amounts of all nutrients
- · Lactose-free and gluten-free
- Halal
- Non-GMO*
- Includes DHA

Precautions

- · Do not heat or use in cooking
- Must be supplemented with protein and fluid in prescribed amounts to completely meet leucine and water requirements

Availability: Hospital/Institutional

Size		Container	List No.
14.1	oz (400 g)	can; 6/case.	 67046

Preparation

Follow physician's instructions carefully.

Weight
(Metric)
8 g
30 g
40 g
60 g
120 g

^{*} Ingredients not genetically engineered.

Powder: Corn Syrup Solids, L-Alanine, High Oleic Safflower Oil, Coconut Oil, Sodium Citrate, Soy Oil, L-Glutamic Acid, L-Aspartic Acid, L-Arginine, L-Proline, L-Lysine Acetate, Calcium Phosphate, Glycine, L-Carnitine, L-Tyrosine; Less than 2% of: L-Phenylalanine, Magnesium Phosphate, Potassium Chloride, L-Serine, L-Threonine, Silicon Dioxide, Modified Corn Starch, L-Valine, L-Isoleucine, L-Histidine, Potassium Citrate, Potassium Phosphate, L-Methionine, C. Cohnii Oil, L-Cystine Dihydrochloride, Calcium Citrate, L-Tryptophan, Calcium Carbonate, Citric Acid, Ascorbic Acid, Taurine, Choline Chloride, Inositol, Ferrous Sulfate, Zinc Sulfate, Ascorbyl Palmitate, Niacinamide, Calcium Pantothenee, dl-Alpha-Tocopheryl Acetate, Mixed Tocopherols, Copper Sulfate, Thiamine Hydrochloride, Manganese Sulfate, Riboflavin, Pyridoxine Hydrochloride, Vitamin A Palmitate, Folic Acid, Biotin, Chromium Chloride, Potassium Iodide, Sodium Selenate, Phylloquinone, Sodium Molybdate, Vitamin D₃, and Vitamin B₁,

	100 g powder		100 g powder
Energy, Cal	410	Vitamin C, mg	60
Protein equivalent, g	30	Choline, mg	100
Source	L-amino acids	Inositol, mg	82
L-Carnitine, mg	1800	Minerals	
Fat, g	35	Calcium, mg	975
Source	High-Oleic Safflower Oil, Coconut Oil, Soy Oil	Phosphorus, mg	760
Linoleic Acid, mg	2200	Magnesium, mg	225
Linolenic Acid, mg	225	Iron, mg	13
DHA	192	Zinc, mg	13
Carbohydrate, g	35	Manganese, mg	0.8
Source	Corn syrup solids	Copper, mg	1.5
Vitamins		lodine, mcg	100
Vitamin A, IU	1500	Selenium, mcg	35
Vitamin D, IU	900	Chromium, mcg	27
Vitamin E, IU	12	Molybdenum, mcg	30
Vitamin K, mcg	60	Sodium, mg	880
Thiamin (Vit B,), mg	3.3	Potassium, mg	1370
Riboflavin (Vit B ₂), mg	1.6	Chloride, mg	940
Vitamin B _s , mg	1.1	Other Characteristics	
Vitamin B ₁₂ , mcg	5	PRSL, mOsm	296
Niacin, mg	16	Osmolality, mOsm/kg H ₂ O	1390 (30 Cal/fl oz)
Folic Acid, mcg	425		
Pantothenic Acid, mg	8		
Biotin, mcg	100		

Ketonex®-1

Amino Acid-Modified Infant Formula With Iron



Description/Indications

Nutrition support of infants and toddlers with maple syrup urine disease (MSUD) or beta-ketothiolase deficiency.

Features

- Branched-chain amino acid-free to allow greater intake of intact protein
- L-carnitine (100 mg/100 g) and taurine (40 mg/100 g) to help supply amounts normally found in human milk and foods of animal origin
- Approximately 40% of energy as fat to help achieve acceptable formula osmolality
- 6.6% of energy as linoleic acid
- Nutrient profile specifically designed for infants and toddlers
- When fed according to the Abbott Support Protocols, provides adequate amounts of all nutrients
- · Lactose-free and gluten-free
- Halal
- Non-GMO*
- Includes DHA/ARA

Precautions

- Must be supplemented with protein and fluid in prescribed amounts to completely meet isoleucine, leucine, valine, and water requirements
- Do not boil mixture or use terminal sterilization

^{*} Ingredients not genetically engineered.

Availability: Hosp	Availability: Hospital/Institutional		
Size	Container	List No	
14.1 oz (400 g)	. can; 6/case	67048	

Preparation

Follow physician's instructions carefully.

Weight
(Metric)
8 g
30 g
40 g
60 g
120 g

Powder: Corn Syrup Solids, High Oleic Safflower Oil, Coconut Oil, Soy Oil, L-Alanine, Modified Corn Starch, L-Proline. Less than 2% of: L-Lysine, L-Arginine, Calcium Phosphate, L-Glutamine, Potassium Phosphate, Glycine, L-Tyrosine, L-Phenylalanine, L-Serine, L-Asparagine, L-Threonine, Sodium Citrate, Potassium Citrate, Magnesium Chloride, L-Histidine, M. Alpina Oil, L-Methionine, C. Cohnii Oil, Calcium Carbonate, Citric Acid, L-Glutamic Acid, Ascorbic Acid, L-Cystine Dihydrochloride, Inositol, L-Tryptophan, Choline Chloride, L-Aspartic Acid, L-Carnitine, Taurine, Taurine, Ferrous Sulfate, Ascorbyl Palmitate, Zinc Sulfate, dl-Alpha-Tocopheryl Acetate, Mixed Tocopherols, Calcium Pantothenate, Niacinamide, Manganese Sulfate, Thiamine Hydrochloride, Copper Sulfate, Vitamin A Palmitate, Riboflavin, Pyridoxine Hydrochloride, Potassium Iodide, Folic Acid, Phylloquinone, Chromium Chloride, Sodium Selenate, Sodium Molybdate, Biotin, Vitamin D., Vitamin B.,, and Salt.

	100 g powder		100 g powder
Energy, Cal	480	Biotin, mcg	23.5
Protein equivalent, g	15	Vitamin C, mg	75
Source	L-amino acids	Choline, mg	95
L-Carnitine, mg	100	Inositol, mg	144
Fat, g	21.7	Minerals	
Source	High-Oleic Safflower Oil, Coconut Oil, Soy Oil	Calcium, mg	575
Linoleic Acid, mg	3650	Phosphorus, mg	400
Linolenic Acid, mg	360	Magnesium, mg	50
DHA, mg	96	Iron, mg	7.6
ARA, mg	115	Zinc, mg	5.2
Carbohydrate, g	53	Manganese, mcg	385
Source	Corn syrup solids	Copper, mcg	408
Vitamins		lodine, mcg	65
Vitamin A, IU	1300	Selenium, mcg	21
Vitamin D, IU	450	Chromium, mcg	12
Vitamin E, IU	12.5	Molybdenum, mcg	13.8
Vitamin K, mcg	60	Sodium, mg	190
Thiamin (Vit B ₁), mcg	1120	Potassium, mg	675
Riboflavin (Vit B ₂), mcg	650	Chloride, mg	325
Vitamin B ₆ , mcg	560	Other Characteristics	
Vitamin B ₁₂ , mcg	0.85	PRSL, mOsm	133
Niacin, mcg	4450	Osmolality, mOsm/kg H₂O	365 (20 Cal/fl oz)
Folic Acid, mcg	88		
Pantothenic Acid, mcg	4000		

C. cohnii oil is a source of DHA M. alpina oil is a source of ARA

Ketonex®-2

Amino Acid-Modified Medical Food



Description/Indications

Nutrition support of children and adults with maple syrup urine disease (MSUD) or beta-ketothiolase deficiency.

Features

- Branched-chain amino acid-free to allow greater intake of intact protein
- L-carnitine (200 mg/100 g) and taurine (125 mg/100 g) to help supply amounts normally found in foods of animal origin
- Approximately 30% of energy as fat to help supply essential fatty acids
- Nutrient profile specifically designed for children and adults
- When fed according to the Abbott Support Protocols, provides adequate amounts of all nutrients
- · Lactose-free and gluten-free
- Halal
- Non-GMO*
- Includes DHA

Precautions

- · Do not heat or use in cooking
- Must be supplemented with protein and fluid in prescribed amounts to completely meet isoleucine, leucine, valine, and water requirements

Preparation

Follow physician's instructions carefully.

Weight
(Metric)
8 g
30 g
40 g
60 g
120 g

^{*} Ingredients not genetically engineered.

Powder: Corn Syrup Solids, L-Alanine, High Oleic Safflower Oil, Coconut Oil, Sodium Citrate, Soy Oil, L-Proline, L-Lysine Acetate, L-Arginine, Calcium Phosphate, L-Glutamine, Glycine, Magnesium Phosphate, Potassium Chloride, L-Tyrosine; Less than 2% of: L-Phenylalanine, L-Serine, Asparagine Monohydrate, L-Threonine, Silicon Dioxide, Modified Corn Starch, Potassium Citrate, L-Histidine, Potassium Phosphate, L-Methionine, C. Cohnii Oil, L-Glutamic Acid, Calcium Citrate, L-Cystine Dihydrochloride, Calcium Carbonate, L-Tryptophan, Citric Acid, L-Aspartic Acid, Ascorbic Acid, L-Carnitine, Taurine, Choline Chloride, Inositol, Ferrous Sulfate, Riboflavin, Pyridoxine Hydrochloride, Vitamin A Palmitate, Folic Acid, Biotin, Chromium Chloride, Potassium Iodide, Sodium Selenate, Phylloquinone, Sodium Molybdate, Vitamin D., and Vitamin B.,

	100 g powder		100 g powder
Energy, Cal	410	Vitamin C, mg	60
Protein equivalent, g	30	Choline, mg	100
Source	L-amino acids	Inositol, mg	82
L-Carnitine, mg	200	Minerals	
Fat, g	14	Calcium, mg	975
Source	High-Oleic Safflower Oil, Coconut Oil, Soy Oil	Phosphorus, mg	760
Linoleic Acid, mg	2200	Magnesium, mg	225
Linolenic Acid, mg	225	Iron, mg	13
DHA, mg	180	Zinc, mg	13
Carbohydrate, g	35	Manganese, mg	0.8
Source	Corn syrup solids	Copper, mg	1.5
Vitamins		lodine, mcg	100
Vitamin A, IU	1500	Selenium, mcg	35
Vitamin D, IU	900	Chromium, mcg	27
Vitamin E, IU	12	Molybdenum, mcg	30
Vitamin K, mcg	60	Sodium, mg	880
Thiamin (Vit B,), mg	3.3	Potassium, mg	1370
Riboflavin (Vit B ₂), mg	1.6	Chloride, mg	940
Vitamin B, mg	1.1	Other Characteristics	
Vitamin B ₁₂ , mcg	5	PRSL, mOsm	296
Niacin, mg	16	Osmolality, mOsm/kg H₂O	1315 (30 Cal/fl oz)
Folic Acid, mcg	425		
Pantothenic Acid, mg	8		
Biotin, mcg	100		

Phenex®-1

Amino Acid-Modified Infant Formula With Iron



Description/Indications

Nutrition support of infants and toddlers with phenylketonuria (PKU) or hyperphenylalaninemia.

Features

- Phenylalanine-free to allow greater intake of intact protein
- L-tyrosine, an essential amino acid that is often deficient in infants with PKU
- L-carnitine (20 mg/100 g) and taurine (40 mg/100 g) to help supply amounts normally found in human milk and foods of animal origin
- Approximately 40% of energy as fat to help achieve acceptable formula osmolality
- . 6.6% of energy as linoleic acid
- Nutrient profile specifically designed for infants and toddlers
- When fed according to the Abbott Support Protocols, provides adequate amounts of all nutrients
- · Lactose-free and gluten-free
- Halal
- Non-GMO*
- Includes DHA/ARA

Precautions

- Must be supplemented with protein and fluid in prescribed amounts to completely meet phenylalanine and water requirements
- Do not boil mixture or use terminal sterilization

Availability: Hospital/Institutional

Size	Container	List No.
14.1 oz (400 g)	. can; 6/case	67052

Preparation

Follow physician's instructions carefully.

Household	Weight
Measure (US)	(Metric)
1 Tablespoon	8 g
1/4 Cup	30 g
1/3 Cup	40 g
1/2 Cup	60 g
1 Cup	120 g

^{*} Ingredients not genetically engineered.

Powder: Corn Syrup Solids, High Oleic Safflower Oil, Coconut Oil, Soy Oil, Modified Corn Starch. Less than 2% of: L-Leucine, L-Tyrosine, L-Proline, L-Lysine, Calcium Phosphate, L-Glutamine, L-Valine, L-Isleucine, L-Arginine, Potassium Phosphate, L-Alanine, Glycine, L-Asparagine, L-Serine, L-Threonine, Sodium Citrate, Potassium Citrate, Magnesium Chloride, L-Histidine, M. Alpina Oil, L-Methionine, C. Cohnii Oil, Calcium Carbonate, Citric Acid, Ascorbic Acid, Glutamic Acid, L-Cystine Dihydrochloride, Inositol, L-Tryptophan, Choline Chloride, L-Aspartic Acid, Taurine, Ferrous Sulfate, Ascorbyl Palmitate, L-Carnitine, Zinc Sulfate, dl-Alpha-Tocopheryl Acetate, Mixed Tocopherols, Calcium Pantothenate, Niacinamide, Manganese Sulfate, Thiamine Hydrochloride, Copper Sulfate, Vitamin A Palmitate, Riboflavin, Pyridoxine Hydrochloride, Potassium Iodide, Folic Acid, Phylloquinone, Chromium Chloride, Sodium Selenate, Sodium Molybdate, Biotin, Vitamin D., Vitamin B., and Salt.

	100 g powder		100 g powder
Energy, Cal	480	Biotin, mcg	23.5
Protein equivalent, g	15	Vitamin C, mg	75
Source	L-amino acids	Choline, mg	95
L-Carnitine, mg	20	Inositol, mg	144
Fat, g	21.7	Minerals	
Source	High-Oleic Safflower Oil, Coconut Oil, Soy Oil	Calcium, mg	575
Linoleic Acid, mg	3650	Phosphorus, mg	400
Linolenic Acid, mg	360	Magnesium, mg	50
DHA, mg	96	Iron, mg	7.6
ARA, mg	115	Zinc, mg	5.2
Carbohydrate, g	53	Manganese, mcg	385
Source	Corn syrup solids	Copper, mcg	408
Vitamins		lodine, mcg	65
Vitamin A, IU	1300	Selenium, mcg	21
Vitamin D, IU	450	Chromium, mcg	12
Vitamin E, IU	12.5	Molybdenum, mcg	13.8
Vitamin K, mcg	60	Sodium, mg	190
Thiamin (Vit B,), mcg	1120	Potassium, mg	675
Riboflavin (Vit B ₂), mcg	650	Chloride, mg	325
Vitamin B _e , mcg	560	Other Characteristics	
Vitamin B ₁₂ , mcg	0.85	PRSL, mOsm	133
Niacin, mcg	4450	Osmolality, mOsm/kg H₂O	370 (20 Cal/fl oz)
Folic Acid, mcg	88		
Pantothenic Acid, mcg	4000		

C. cohnii oil is a source of DHA M. alphina oil is a source of ARA

Phenex®-2

Amino Acid-Modified Medical Food



Description/Indications

Nutrition support of children and adults with phenylketonuria (PKU) or hyperphenylalaninemia.

Features

- Phenylalanine-free to allow greater intake of intact protein
- L-tyrosine, an essential amino acid that is often deficient in patients with PKU
- L-carnitine (40 mg/100 g) and taurine (125 mg/100 g) to help supply amounts normally found in foods of animal origin
- Approximately 30% of energy as fat to help supply essential fatty acids
- Nutrient profile specifically designed for children and adults
- When fed according to the Abbott Support Protocols, provides adequate amounts of all nutrients
- · Lactose-free and gluten-free
- Halal
- Non-GMO*
- Includes DHA

Precautions

- . Do not heat or use in cooking
- Must be supplemented with protein and fluid in prescribed amounts to completely meet phenylalanine and water requirements

Availability: Hospital/Institutional

Flavor	Size	Container	List No.
Unflavored	14.1 oz (400 g)	.can; 6/case	67054
Vanilla	14.1 oz (400 g)	.can; 6/case	67056

Preparation

Follow physician's instructions carefully.

Household	Weight
Measure (US)	(Metric)
1 Tablespoon	8 g
1/4 Cup	30 g
1/3 Cup	40 g
1/2 Cup	60 g
1 Cup	

^{*} Ingredients not genetically engineered.

Powder (Unflavored): Corn Syrup Solids, High Oleic Safflower Oil, Coconut Oil, Sodium Citrate, Soy Oil, L-Leucine, L-Tyrosine, L-Proline, L-Lysine Acetate, L-Glutamine, L-Valine, Calcium Phosphate, L-Isoleucine, L-Arginine, L-Alanine, Glycine. Less than 2% of: Magnesium Phosphate, Potassium Chloride, L-Asparagine Monohydrate, L-Serine, L-Threonine, Silicon Dioxide, Moldified Corn Starch, L-Histidine, Potassium Chrate, Potassium Chloride, L-Asparagine Monohydrate, L-Oystine Dihydrochloride, L-Tryptophan, Calcium Carbonate, Citric Acid, L-Aspartic Acid, Ascorbic Acid, Taurine, Choline Chloride, Inositol, Ferrous Sulfate, Zinc Sulfate, L-Carnitine, Ascorbyl Palmitate, Niacinamide, Calcium Pantothenate, dl-Alpha-Tocopheryl Acetate, Mixed Tocopherols, Copper Sulfate, Thiamine Hydrochloride, Manganese Sulfate, Riboflavin, Pyridoxine Hydrochloride, Vitamin A Palmitate, Folic Acid, Biotin, Chromium Chloride, Potassium Iodide, Sodium Selenate, Phylloquinone, Sodium Molybdate, Vitamin D., and Vitamin B.,

	100 g powder		100 g powder
Energy, Cal	410	Vitamin C, mg	60
Protein equivalent, g	30	Choline, mg	100
Source	L-amino acids	Inositol, mg	82
L-Carnitine, mg	40	Minerals	
Fat, g	14	Calcium, mg	975
Source	High-Oleic Safflower Oil, Coconut Oil, Soy Oil	Phosphorus, mg	760
Linoleic Acid, mg	2200	Magnesium, mg	225
Linolenic Acid, mg	225	Iron, mg	13
DHA, mg	180	Zinc, mg	13
Carbohydrate, g	35	Manganese, mg	0.8
Source	Corn syrup solids	Copper, mg	1.5
Vitamins		lodine, mcg	100
Vitamin A, IU	1500	Selenium, mcg	35
Vitamin D, IU	900	Chromium, mcg	27
Vitamin E, IU	12	Molybdenum, mcg	30
Vitamin K, mcg	60	Sodium, mg	880
Thiamin (Vit B ₁), mg	3.3	Potassium, mg	1370
Riboflavin (Vit B ₂), mg	1.6	Chloride, mg	940
Vitamin B ₆ , mg	1.1	Other Characteristics	
Vitamin B ₁₂ , mcg	5	PRSL, mOsm	296
Niacin, mg	16	Osmolality, mOsm/kg H₂O	1215 (30 cal/fl oz)
Folic Acid, mcg	425		
Pantothenic Acid, mg	8		
Biotin, mcg	100		

Pro-Phree®

Protein-Free Energy Module With Iron, Vitamins & Minerals



Description/Indications

Nutrition support of infants and toddlers who require extra calories, minerals, and vitamins and/or protein restriction.

Features

- Protein-free to permit protein restriction or the addition of extra energy, minerals, and vitamins
- L-carnitine (25 mg/100 g) and taurine (50 mg/100 g) to help supply amounts normally found in human milk and foods of animal origin
- Approximately 49% of energy as fat to help achieve acceptable formula osmolality
- 7.8% of energy as linoleic acid
- Nutrient profile specifically designed for infants and toddlers. May be used by children and adults
- · Lactose-free and gluten-free
- Halal
- Non-GMO^{*}
- Includes DHA/ARA

Precautions

- Must be used with a source of intact protein or amino acids and fluid to completely meet nutrient and water requirements
- Not a substitute for standard infant formula or intended for use as the sole source of nutrition
- Do not boil mixture or use terminal sterilization

Availability: Hospital/Institutional

Size	Container	List No.
14.1 oz (400 g)	can; 6/case	67030

Preparation

Follow physician's instructions carefully.

Household	Weight
Measure (US)	(Metric)
1 Tablespoon	8 g
1/4 Cup	30 g
1/3 Cup	40 g
1/2 Cup	60 g
1 Cup	120 g

^{*} Ingredients not genetically engineered.

Powder: Corn Syrup Solids, High Oleic Safflower Oil, Coconut Oil, Soy Oil, Modified Corn Starch. Less than 2% of: Calcium Phosphate, Potassium Phosphate, Sodium Citrate, Potassium Citrate, Magnesium Chloride, M. Alpina Oil, C. Cohnil Oil, Calcium Carbonate, Citric Acid, Ascorbic Acid, Inositol, Choline Chloride, Taurine, Ferrous Sulfate, Ascorbyl Palmitate, L-Carnitine, Circ Sulfate, dl-Alpha-Tocopheryl Acetate, Mixed Tocopherols, Calcium Pantothenate, Niacinamide, Manganese Sulfate, Thiamine Hydrochloride, Copper Sulfate, Vitamin A Palmitate, Riboflavin, Pyridoxine Hydrochloride, Potassium Iodide, Folic Acid, Phylloquinone, Chromium Chloride, Sodium Selenate, Sodium Molybdate, Biotin, Vitamin D., Vitamin B.,, and Salt.

	100 g powder		100 g powder
Energy, Cal	510	Biotin, mcg	30
Protein equivalent, g	0	Vitamin C, mg	85
Source		Choline, mg	120
L-Carnitine, mg	25	Inositol, mg	180
Fat, g	28	Minerals	
Source	High-Oleic Safflower Oil, Coconut Oil, Soy Oil	Calcium, mg	750
Linoleic Acid, mg	4600	Phosphorus, mg	525
Linolenic Acid, mg	470	Magnesium, mg	70
DHA, mg	120	Iron, mg	10
ARA, mg	145	Zinc, mg	6.6
Carbohydrate, g	65	Manganese, mcg	485
Source	Corn syrup solids	Copper, mcg	510
Vitamins		lodine, mcg	80
Vitamin A, IU	1700	Selenium, mcg	27
Vitamin D, IU	575	Chromium, mcg	16
Vitamin E, IU	16	Molybdenum, mcg	17
Vitamin K, mcg	80	Sodium, mg	250
Thiamin (Vit B ₁), mcg	1400	Potassium, mg	875
Riboflavin (Vit B ₂), mcg	825	Chloride, mg	350
Vitamin B ₆ , mcg	710	Other Characteristics	
Vitamin B ₁₂ , mcg	1.1	PRSL, mOsm	60
Niacin, mcg	5550	Osmolality, mOsm/kg H ₂ O	205 (20 Cal/fl oz)
Folic Acid, mcg	110		
Pantothenic Acid, mcg	5200		

C. cohnii oil is a source of DHA M. alphina oil is a source of ARA

Propimex®-1

Amino Acid-Modified Infant Formula With Iron



Description/Indications

Nutrition support of infants and toddlers with propionic or methylmalonic acidemia.

Features

- Methionine- and valine-free to allow greater intake of intact protein; low in isoleucine and threonine
- L-carnitine (900 mg/100 g) to help supply amount normally found in human milk and foods of animal origin and to help excrete toxic metabolites
- Taurine (40 mg/100 g) to supply amount normally found in human milk and foods of animal origin
- Approximately 40% of energy as fat to help achieve acceptable formula osmolality
- 6.6% of energy as linoleic acid
- Nutrient profile specifically designed for infants and toddlers
- When fed according to the Abbott Support Protocols. provides adequate amounts of all nutrients
- · Lactose-free and gluten-free
- Halal
- Non-GMO*
- Includes DHA/ARA

Precautions

- Must be supplemented with protein and fluid in prescribed amounts to completely meet isoleucine, methionine, threonine, valine, and water requirements
- Do not boil mixture or use terminal sterilization

Availability: Hospital/Institutional

Size	Container	List No.
14.1 oz (400 g)	can; 6/case	67058

Preparation

Follow physician's instructions carefully.

Household	Weight
Measure (US)	(Metric)
1 Tablespoon	8 g
1/4 Cup	30 g
1/3 Cup	40 g
1/2 Cup	60 g
1 Cup	120 g

^{*} Ingredients not genetically engineered.

Powder: Corn Syrup Solids, High Oleic Safflower Oil, Coconut Oil, Soy Oil, L-Alanine, L-Glutamic Acid, Modified Corn Starch. Less than 2% of: L-Proline, L-Aspartic Acid, L-Arginine, L-Lysine, L-Leucine, Calcium Phosphate, Potassium Phosphate, L-Carnitine, L-Tyrosine, L-Phenylalanine, L-Serine, Sodium Citrate, L-Cystine Dihydrochloride, Potassium Citrate, Glycine, Magnesium Chloride, L-Histotine, L-Histotine, L-Threonine, Taurine, Ferrous Sulfate, Ascorbyl Palmitate, Zinc Sulfate, dl-Alpha-Tocopheryl Acetate, Mixed Tocopherols, Calcium Pantothenate, Niacinamide, Manganese Sulfate, Thiamine Hydrochloride, Copper Sulfate, Vitamin A Palmitate, Riboflavin, Pyridoxine Hydrochloride, Potassium Iodide, Folic Acid, Phylloquinone, Chromium Chloride, Sodium Selenate, Sodium Molybdate, Biotin, Vitamin B₁₁, and Salt.

	100 g powder		100 g powder
Energy, Cal	480	Biotin, mcg	23.5
Protein equivalent, g	15	Vitamin C, mg	75
Source	L-amino acids	Choline, mg	95
L-Carnitine, mg	900	Inositol, mg	144
Fat, g	21.7	Minerals	
Source	High-Oleic Safflower Oil, Coconut Oil, Soy Oil	Calcium, mg	575
Linoleic Acid, mg	3650	Phosphorus, mg	400
Linolenic Acid, mg	360	Magnesium, mg	50
DHA, mg	96	Iron, mg	7.6
ARA, mg	115	Zinc, mg	5.2
Carbohydrate, g	53	Manganese, mg	385
Source	Corn syrup solids	Copper, mg	408
Vitamins		lodine, mcg	65
Vitamin A, IU	1300	Selenium, mcg	21
Vitamin D, IU	450	Chromium, mcg	12
Vitamin E, IU	12.5	Molybdenum, mcg	13.8
Vitamin K, mcg	60	Sodium, mg	190
Thiamin (Vit B₁), mg	1120	Potassium, mg	675
Riboflavin (Vit B ₂), mg	650	Chloride, mg	410
Vitamin B ₆ , mg	560	Other Characteristics	
Vitamin B ₁₂ , mcg	0.85	PRSL, mOsm	136
Niacin, mg	4450	Osmolality, mOsm/kg H ₂ O	370 (20 Cal/fl oz)
Folic Acid, mg	88		
Pantothenic Acid, mcg	4000		

C. cohnii oil is a source of DHA M. alpina oil is a source of ARA

Propimex®-2

Amino Acid-Modified Medical Food



Description/Indications

Nutrition support of children and adults with propionic or methylmalonic acidemia.

Features

- Methionine- and valine-free to allow greater intake of intact protein; low in isoleucine and threonine
- L-carnitine (1800 mg/100 g) to help supply amount normally found in foods of animal origin and to help excrete toxic metabolites
- Taurine (125 mg/100 g) to help supply amount normally found in foods of animal origin
- Approximately 30% of energy as fat to help supply essential fatty acids
- Nutrient profile specifically designed for children and adults
- When fed according to the Abbott Support Protocols, provides adequate amounts of all nutrients
- · Lactose-free and gluten-free
- Halal
- Non-GMO*
- Includes DHA

Precautions

- · Do not heat or use in cooking
- Must be supplemented with protein and fluid in prescribed amounts to completely meet isoleucine, methionine, threonine, valine, and water requirements

Availability: Hospital/Institutional

Size	Container	List No.
14.1 oz (400 g)	can; 6/case	67060

Preparation

Follow physician's instructions carefully.

ight tric)
tric)
,
. 8 g
30 g
40 g
60 g
20 g

^{*} Ingredients not genetically engineered.

Powder: Corn Syrup Solids, L-Alanine, L-Glutamic Acid, High Oleic Safflower Oil, Coconut Oil, Sodium Citrate, Soy Oil, L-Proline, L-Aspartic Acid, L-Arginine, L-Lysine Acetate, L-Leucine, Calcium Phosphate, L-Carritine, L-Tyrosine. Less than 2% of: L-Phenylalanine, L-Serine, Magnesium Phosphate, Potassium Chloride, L-Cystine Dihydrochloride, Silicon Dioxide, Glycine, Modified Corn Starch, L-Histidine, Potassium Chrate, Potassium Phosphate, C. Cohnii Oil, Calcium Citrate, L-Tryptophan, Calcium Carbonate, Citrach, Acid, L-Isoleucine, Ascorbic Acid, L-Threonine, Taurine, Choline Chloride, Inositol, Ferrous Sulfate, Zinc Sulfate, Ascorbyl Palmitate, Niacinamide, Calcium Pantothenate, dl-Alpha-Tocopheryl Acetate, Mixed Tocopherols, Copper Sulfate, Thiamine Hydrochloride, Manganese Sulfate, Riboflavin, Pyridoxine Hydrochloride, Vitamin A Palmitate, Folic Acid, Biotin, Chromium Chloride, Potassium lodide, Sodium Selenate, Phylloquinone, Sodium Molybdate, Vitamin D,, and Vitamin B,.

	100 g powder		100 g powder
	Too y powder		100 g powder
Energy, Cal	410	Vitamin C, mg	60
Protein equivalent, g	30	Choline, mg	100
Source	L-amino acids	Inositol, mg	82
L-Carnitine, mg	1800	Minerals	
Fat, g	13	Calcium, mg	975
Source	High-Oleic Safflower Oil, Coconut Oil, Soy Oil	Phosphorus, mg	760
Linoleic Acid, mg	2200	Magnesium, mg	225
Linolenic Acid, mg	225	Iron, mg	13
DHA, mg	180	Zinc, mg	13
Carbohydrate, g	35	Manganese, mg	0.8
Source	Corn syrup solids	Copper, mg	1.5
Vitamins		lodine, mcg	100
Vitamin A, IU	1500	Selenium, mcg	35
Vitamin D, IU	900	Chromium, mcg	27
Vitamin E, IU	12	Molybdenum, mcg	30
Vitamin K, mcg	60	Sodium, mg	880
Thiamin (Vit B ₁), mg	3.3	Potassium, mg	1370
Riboflavin (Vit B ₂), mg	1.6	Chloride, mg	1160
Vitamin B _e , mg	1.1	Other Characteristics	
Vitamin B ₁₂ , mcg	5	PRSL, mOsm	302
Niacin, mg	16	Osmolality, mOsm/kg H ₂ O	1350 (30 Cal/fl oz)
Folic Acid, mcg	425		
Pantothenic Acid, mg	8		
Biotin, mcg	100		

ProViMin®

Protein-Vitamin-Mineral Module With Iron



Description/Indications

For use in the management of patients who require a formula modified in carbohydrate, fat, and/or increased protein: abetalipoproteinemia; cholestasis; chylothorax; fatty acid oxidation defects; glutaric aciduria type II; hyperlipoproteinemia type I (fasting chylomicronemia); hypobetalipoproteinemia; lymphangiectasis, intestinal malabsorption of carbohydrate and/or fat; supplement for any patient who requires increased protein, minerals, and vitamins; X-linked adrenoleukodystrophy.

Features

- · High-quality source of protein (casein)
- Virtually carbohydrate- and fat-free; provides flexibility in prescribing a formula in which the amount and type of carbohydrate and fat can be added per individual infant's need and tolerance
- Meets the Dietary Reference Intakes and the American Academy of Pediatrics Committee on Nutrition (AAPCON) recommendations for vitamin and mineral intakes for infants when fed at a dilution of 3.25 g protein/100 Cal
- Calcium-to-phosphorus ratio of 1.4:1, which is similar to that of standard infant formulas and meets AAPCON recommendations for that ratio

- L-carnitine (40 mg/100 g) and taurine (110 mg/100 g)
- · Lactose-free and gluten-free
- · Kosher, Halal

Precautions

- Does not supply sufficient amounts of energy as carbohydrate and fat, or linoleic acid. These nutrients should be supplied from other sources under the supervision of a physician
- Do not boil mixture or use terminal sterilization

Availability: Hospital/Institutional

Size	Container	List No.
5.3 oz (150 g)	can; 6/case	50260

Preparation

Follow physician's instructions carefully. To prepare 1000 mL of a 20 Cal/fl oz formula, add 30 g ProViMin, 34 g fat, and 69 g carbohydrate to 900 mL (30 fl oz) water.

Household	Weight
Measure (US)	(Metric)
1 Tablespoon	2.9 g
1/4 Cup	11 g
1/2 Cup	20 g
2/3 Cup	30 g
1 Cup	44 g

Powder: Sodium Caseinate, Calcium Phosphate, Potassium Citrate, Potassium Chloride. Less than 2% of: Magnesium Chloride, Ascorbic Acid, Coconut Oil, Choline Chloride, Ferrous Sulfate, Taurine, m-Inositol, L-Methionine, d-Alpha Tocopheryl Acetate, Zinc Sulfate, L-Carritine, Niacinamide, Calcium Pantothenate, Vitamin A Palmitate, Copper Sulfate, Thiamine Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Manganese Sulfate, Potassium Iodide, Phylloquinone, Biotin, Sodium Selenate, Vitamin D₃, Vitamin B₁₂, Salt and Calcium Carbonate. Contains milk ingredients.

NUTRITION INFORMA	TION		
100 g powder			100 g powder
Energy, Cal	313	Minerals	
Protein equivalent, g	73	Calcium, mg	2400
Source	Casein, L-amino acids	Calcium, mEq	120.0
Fat, g	1.4	Phosphorus, mg	1700
Source	Coconut oil	Magnesium, mg	200
Linoleic Acid, mg	0.0	Iron, mg	40
Carbohydrate, g	2.0	Zinc, mg	17
Source	None added	Manganese, mcg	200
Vitamins		Copper, mcg	2100
Vitamin A, IU	6740	lodine, mcg	335
Vitamin D, IU	1000	Selenium, mcg	46
Vitamin E, IU	67	Sodium, mg	1200
Vitamin K, mcg	90	Sodium, mEq	52
Thiamin (Vit B₁), mcg	2240	Sodium, mg	3300
Riboflavin (Vit B ₂), mcg	2020	Potassium, mg	84
Vitamin B ₆ , mcg	1350	Chloride, mg	2300
Vitamin B ₁₂ , mcg	5.6	Chloride, mEq	65
Niacin, mcg	24,000	Other Characteristics	
Folic Acid, mcg	320	PRSL, mOsm	673
Pantothenic Acid, mcg	10,100	Osmolality, mOsm/kg H ₂ O	Will vary with the amount of
Biotin, mcg	100		carbohydrate, fat, and water added
Vitamin C, mg	200		
Choline, mg	335		
Inositol, mg	105		

RCF®

No Added Carbohydrate Soy Infant Formula Base With Iron



Description/Indications

For use in the dietary management of patients unable to tolerate the type or amount of carbohydrate in milk or conventional infant formulas; or seizure disorders requiring a ketogenic diet.

Features

- The only commercial formula available for the dietary management of seizures in infants
- Formulated to allow physician to prescribe type and amount of carbohydrate (that can be tolerated) with the assurance that other nutrient needs will be met
- Soy protein isolate to avoid symptoms of cow's-milkprotein allergy or sensitivity
- 1.8 mg of iron (as ferrous sulfate) per 100 Cal if carbohydrate is added to make a 20 Cal/fl oz feeding
- L-carnitine (3 mg/100 mL) and taurine (12 mg/100 mL)
- · Lactose-free and gluten-free
- · Kosher, Halal

Availability: Hospital/Institutional

Size	Container	List No.
Concentrat	ed Liquid: (24 Cal/fl oz)	
13 fl oz (38	4 mL) can; 12/case	00108

Preparation

Add water and other ingredients as directed by physician before feeding. Use only under the supervision of a physician.

Ingredients

Concentrated Liquid: Water (88%), Soy Protein Isolate (5%), High Oleic Safflower Oil (3%), Soy Oil (2%), Coconut Oil (2%). Less than 1% of: Calcium Phosphate, Potassium Citrate, Potassium Chloride, Mangnesium Chloride, Monoglycerides, Soy Lecithin, Carrageenan, Salt, L-Methionine, Ascorbic Acid, Potassium Hydroxide, Choline Chloride, Taurine, m-Inositol, Ferrous Sulfate, Zinc Sulfate, L-Carnitine, Niacinamide, d-Alpha-Tocopheryl Acetate, Calcium Pantothenate, Cupric Sulfate, Thiamine Chloride Hydrochloride, Vitamin A Palmitate, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Potassium Iodide, Phylloquinone, Biotin, Beta-Carotene, Sodium Selenate, Vitamin D, and Cyanocobalamin. Contains soy ingredients.

	100 mL concentrated liquid	100 Cal		100 mL concentrated liquid	100 Cal
Energy, Cal	81	100	Vitamin C, mg	12	9
Volume, mL	100	148	Choline, mg	15.7	12
Protein, g	4.0	3.0	Inositol, mg	6.5	5
% Total Calories	20	12	Minerals		
Source	Soy protein isolate, L-methionine		Calcium, mg	140	105
Fat, g	7.2	5.3	Calcium, mEq	7.0	5.2
% Total Calories	80	48	Phosphorus, mg	100	75
Source	High oleic safflower, coconut, and soy oils		Magnesium, mg	10.0	7.5
Linoleic Acid, mg	1352	1000	Iron, mg	2.4	1.8
Carbohydrate, g	0.07	10.1	Zinc, mg	1.0	0.75
% Total Calories	0	40	Manganese, mcg	34	25
Source	None added		Copper, mcg	100	75
Vitamins			lodine, mcg	20.3	15
Vitamin A, IU	405	300	Selenium, mcg	2.7	2.0
Vitamin D, IU	81	60	Sodium, mg	59.1	44
Vitamin E, IU	2.0	1.5	Sodium, mEq	2.6	1.9
Vitamin K, mcg	15	11	Potassium, mg	146	108
Thiamin (Vit B,), mcg	80	60	Potassium, mEg	3.7	2.8
Riboflavin (Vit B ₂), mcg	120	90	Chloride, mg	83	62
Vitamin B ₆ , mcg	80	60	Chloride, mEq	2.3	1.8
Vitamin B ₁₂ , mcg	0.6	0.4			
Niacin, mcg	1800	1350	Other Characteristics		
Folic Acid, mcg	20	15	PRSL, mOsm	35	25.8
Pantothenic Acid, mcg	1000	750	Water, g	88	133
Biotin, mcg	6.1	4.5	Osmolality, mOsm/kg H ₂ O	90	168

^{*} As prepared: 54 g of carbohydrate, 12 fl oz of water and mixed with 13 fl oz of RCF.

Tyrex®-1

Amino Acid-Modified Infant Formula With Iron



Description/Indications

Nutrition support of infants and toddlers with tyrosinemia type I, II or III.

Features

- Phenylalanine- and tyrosine-free to allow greater intake of intact protein
- L-carnitine (20 mg/100 g) and taurine (40 mg/100 g) to help supply amounts normally found in human milk and foods of animal origin
- Approximately 40% of energy as fat to help achieve acceptable formula osmolality
- 6.6% of energy as linoleic acid
- Nutrient profile specifically designed for infants and toddlers
- When fed according to the Abbott Support Protocols, provides adequate amounts of all nutrients
- · Lactose-free and gluten-free
- Halal
- Non-GMO*
- Includes DHA/ARA

Precautions

- Must be supplemented with protein and fluid in prescribed amounts to completely meet phenylalanine, tyrosine, and water requirements
- Do not boil mixture or use terminal sterilization

^{*} Ingredients not genetically engineered.

Availab	ility: Hospital/Institutional	
Size	Container	List No.
14.1 oz ((400 g) can; 6/case	67062

Preparation

Follow physician's instructions carefully.

Approximate weights for unpacked, level, dry US standard household measures for powder

modeonicia modeodico ici pomaci	
Household	Weight
Measure (US)	(Metric)
1 Tablespoon	8 g
1/4 Cup	30 g
1/3 Cup	40 g
1/2 Cup	60 g
1 Cup	120 g

Ingredients

Powder: Corn Syrup Solids, High Oleic Safflower Oil, Coconut Oil, Soy Oil, Modified Corn Starch. Less than 2% of: L-Leucine, L-Proline, L-Lysine, Calcium Phosphate, L-Arginine, L-Gutamine, L-Valine, L-Islamine, E-Valine, L-Islamine, E-Valine, L-Islamine, E-Valine, L-Islamine, E-Valine, L-Islamine, E-Serine, L-Threonine, Sodium Citrate, Potassium Citrate, Magnesium Chloride, L-Histidine, M. Alpina Oil, L-Methionine, C. Cohnii Oil, Calcium Carbonate, Citric Acid, L-Gutamic Acid, Ascorbic Acid, L-Cystine Dihydrochloride, Inositol, L-Tryptophan, Choline Chloride, L-Aspartic Acid, Taurine, Ferrous Sulfate, Ascorbyl Palmitate, L-Carnitine, Zinc Sulfate, dl-Alpha-Tocopheryl Acetate, Mixed Tocopherols, Calcium Pantothenate, Niacinamide, Manganese Sulfate, Thiamine Hydrochloride, Copper Sulfate, Vitamin A Palmitate, Riboflavin, Pyridoxine Hydrochloride, Potassium Iodide, Folic Acid, Phylloquinone, Chromium Chloride, Sodium Selenate, Sodium Molybdate, Biotin, Vitamin B, vitamin B, and Salt.

	100 g powder		100 g powder
Energy, Cal	480	Biotin, mcg	23.5
Protein equivalent, g	15	Vitamin C, mg	75
Source	L-amino acids	Choline, mg	95
L-Carnitine, mg	20	Inositol, mg	144
Fat, g	21.7	Minerals	
Source	High-Oleic Safflower Oil, Coconut Oil, Soy Oil	Calcium, mg	575
Linoleic Acid, mg	3650	Phosphorus, mg	400
Linolenic Acid, mg	360	Magnesium, mg	50
DHA, mg	96	Iron, mg	7.6
ARA, mg	115	Zinc, mg	5.2
Carbohydrate, g	53	Manganese, mcg	385
Source	Corn syrup solids	Copper, mcg	408
Vitamins		lodine, mcg	65
Vitamin A, IU	1300	Selenium, mcg	21
Vitamin D, IU	450	Chromium, mcg	12
Vitamin E, IU	12.5	Molybdenum, mcg	13.8
Vitamin K, mcg	60	Sodium, mg	190
Thiamin (Vit B ₁), mcg	1120	Potassium, mg	675
Riboflavin (Vit B ₂), mcg	650	Chloride, mg	325
Vitamin B, mcg	560	Other Characteristics	
Vitamin B ₁₂ , mcg	0.85	PRSL, mOsm	133
Niacin, mcg	4450	Osmolality, mOsm/kg H₂O	360 (20 Cal/fl oz)
Folic Acid, mcg	88		
Pantothenic Acid, mcg	4000		

C. cohnii oil is a source of DHA M. alpina oil is a source of ARA

Tyrex®-2 Amino Acid-Modified Medical Food



Description/Indications

For nutrition support of children and adults with tyrosinemia type I, II, or III.

Features

- Phenylalanine- and tyrosine-free to allow greater intake of intact protein
- L-carnitine (40 mg/100 g) and taurine (125 mg/100 g) to supply amounts normally found in foods of animal origin
- Approximately 30% of energy as fat to help supply essential fatty acids
- Nutrient profile specifically designed for children and adults
- When fed according to the Abbott Support Protocols. provides adequate amounts of all nutrients
- · Lactose-free and gluten-free
- Halal
- Non-GMO^{*}
- Includes DHA

Precautions

- · Do not heat or use in cooking
- Must be supplemented with protein and fluid in prescribed amounts to completely meet phenylalanine, tyrosine, and water requirements

Availability: Hospital/Institutional

Size	Container	List No.
14.1 oz (400 g)	can; 6/case	67064

Preparation

Follow physician's instructions carefully.

Approximate weights for unpacked, level, dry US standard household measures for powder

Household	Weight
Measure (US)	(Metric)
1 Tablespoon	8 g
1/4 Cup	30 g
1/3 Cup	40 g
1/2 Cup	60 g
1 Cup	120 g

^{*} Ingredients not genetically engineered.

Ingredients

Powder: Corn Syrup Solids, High Oleic Safflower Oil, Coconut Oil, Sodium Citrate, Soy Oil, L-Leucine, L-Proline, L-Lysine Acetate, L-Arginine, L-Glutamine, Calcium Phosphate, L-Valine, L-Isoleucine, L-Alanine, Glycine, Magnesium Phosphate, Potassium Chloride. Less than 2% of. Asparagine Monohydrate, L-Serine, L-Threonine, Silicon Dioxide, Modified Corn Starch, Potassium Citrate, L-Histidine, Potassium Phosphate, L-Methionine, C. Cohnii Oil, L-Glutamic Acid, Calcium Citrate, L-Cystine Dihydrochloride, L-Tryptophan, Calcium Carbonate, Citric Acid, L-Aspartic Acid, Ascorbic Acid, Taurine, Choline Chloride, Inositol, Ferrous Sulfate, Zinc Sulfate, L-Carnitine, Ascorbyl Palmitate, Niacinamide, Calcium Pantothenate, dl-Alpha-Tocopheryl Acetate, Mixed Tocopherols, Copper Sulfate, Thiamine Hydrochloride, Manganese Sulfate, Riboflavin, Pyridoxine Hydrochloride, Vitamin A Palmitate, Folic Acid, Biotin, Chromium Chloride, Potassium Iolide, Sodium Selenate, Phylloquinone, Sodium Molybdate, Vitamin D³, and Vitamin B³, and Vitamin B³, and Vitamin B³.

	100 g powder		100 g powder
Energy, Cal	410	Vitamin C, mg	60
Protein equivalent, g	30	Choline, mg	100
Source	L-amino acids	Inositol, mg	82
L-Carnitine, mg	40	Minerals	
Fat, g	14	Calcium, mg	975
Source	High-Oleic Safflower Oil, Coconut Oil, Soy Oil	Phosphorus, mg	760
Linoleic Acid, mg	2200	Magnesium, mg	225
Linolenic Acid, mg	225	Iron, mg	13
DHA, mg	180	Zinc, mg	13
Carbohydrate, g	35	Manganese, mg	0.8
Source	Corn syrup solids	Copper, mg	1.5
Vitamins		lodine, mcg	100
Vitamin A, IU	1500	Selenium, mcg	35
Vitamin D, IU	900	Chromium, mcg	27
Vitamin E, IU	12	Molybdenum, mcg	30
Vitamin K, mcg	60	Sodium, mg	880
Thiamin (Vit B ₁), mg	3.3	Potassium, mg	1370
Riboflavin (Vit B ₂), mg	1.6	Chloride, mg	940
Vitamin B, mg	1.1	Other Characteristics	
Vitamin B ₁₉ , mg	5	PRSL, mOsm	296
Niacin, mg	16	Osmolality, mOsm/kg H ₂ O	1330 (30 Cal/fl oz)
Folic Acid, mcg	425		
Pantothenic Acid, mg	8		
Biotin, mcg	100		

C. cohnii oil is a source of DHA





Additional Information

Newborn Metric Conversion Tables

Temperature Fahrenheit (F) to Celsius (C)

°F	°C	°F	°C	°F	°C	°F	°C
95.0	35.0	98.0	36.7	101.0	38.3	104.0	40.0
95.2	35.1	98.2	36.8	101.2	38.4	104.2	40.1
95.4	35.2	98.4	36.9	101.4	38.6	104.4	40.2
95.6	35.3	98.6	37.0	101.6	38.7	104.6	40.3
95.8	35.4	98.8	37.1	101.8	38.8	104.8	40.4
96.0	35.6	99.0	37.2	102.0	38.9	105.0	40.6
96.2	35.7	99.2	37.3	102.2	39.0	105.2	40.7
96.4	35.8	99.4	37.4	102.4	39.1	105.4	40.8
96.6	35.9	99.6	37.6	102.6	39.2	105.6	40.9
96.8	36.0	99.8	37.7	102.8	39.3	105.8	41.0
97.0	36.1	100.0	37.8	103.0	39.4	106.0	41.1
97.2	36.2	100.2	37.9	103.2	39.6	106.2	41.2
97.4	36.3	100.4	38.0	103.4	39.7	106.4	41.3
97.6	36.4	100.6	38.1	103.6	39.8	106.6	41.4
97.8	36.6	100.8	38.2	103.8	39.9	106.8	41.6

Note: ${}^{\circ}C = ({}^{\circ}F-32) \times 5/9$. Celsius temperature equivalents rounded to one decimal place by adding 0.1 when second decimal place is 5 or greater.

The metric system replaces the term "Centigrade" with "Celsius" (the name of the inventor of the scale).

Length Inches to centimeters

One-inch increments To obtain centimeters equivalent to 22 inches, read "20" on top scale, "2" on side scale; equivalent is 55.9 centimeters.

Inches	0	10	20	30	40	
0	0 0		50.8	76.2	101.6	
1	2.5	27.9	53.3	78.7	104.1	
2	5.1	30.5	55.9	81.3	106.7	
3	7.6	33.0	58.4	83.8	109.2	
4	10.2	35.6	61.0	86.4	111.8	
5	12.7	38.1	63.5	88.9	114.3	
6	15.2	40.6	66.0	91.4	116.8	
7	17.8	43.2	68.6	68.6 94.0		
8	20.3	45.7	71.1	96.5	121.9	
9	22.9	48.3	73.7	99.1	124.5	

One-Quarter (1/4) in increments To obtain centimeters equivalent to 14% inches, read "14" on top scale, "%" on side scale; equivalent is 37.5 centimeters.

10-15 Inches

	10	11	12	13	14	15
0	25.4	27.9	30.5	33.0	35.6	38.1
1/4	26.0	28.6	31.1	33.7	36.2	38.7
1/2	26.7	29.2	31.8	34.3	36.8	39.4
3/4	27.3	29.8	32.4	34.9	37.5	40.0

16-21 Inches

	16	17	18	19	20	21
0	40.6	43.2	45.7	48.3	50.8	53.3
1/4	41.3	43.8	46.4	48.9	51.4	54.0
1/2	41.9	44.5	47.0	49.5	52.1	54.6
3/4	42.5	45.1	47.6	50.2	52.7	55.2

Note: 1 inch = 2.540 centimeters. Centimeter equivalents rounded to one decimal place by adding 0.1 when second decimal place is 5 or greater; for example, 33.48 becomes 33.5.

Weight (Mass) Pounds and ounces to grams

Example: To obtain grams equivalent to 6 pounds, 8 ounces, read "6" on top scale, "8" on side scale; equivalent is 2948 grams.

Note: 1 pound = 453.592 g; 1 ounce = 28.350 g; 1000 g = 1 kilogram. Gram equivalents have been rounded to whole numbers by adding one when the first decimal place is 5 or greater.

	Pounds															
		0	1	2	3	4	5	6	7	8	9	10	-11	12	13	14
	0	0	454	907	1361	1814	2268	2722	3175	3629	4082	4536	4990	5443	5897	6350
	1	28	482	936	1389	1843	2296	2750	3203	3657	4111	4564	5018	5471	5925	6379
	2	57	510	964	1417	1871	2325	2778	3232	3685	4139	4593	5046	5500	5953	6407
	3	85	539	992	1446	1899	2353	2807	3260	3714	4167	4621	5075	5528	5982	6435
	4	113	567	1021	1474	1928	2381	2835	3289	3742	4196	4649	5103	5557	6010	6464
	5	142	595	1049	1503	1956	2410	2863	3317	3770	4224	4678	5131	5585	6038	6492
ွှ	6	170	624	1077	1531	1984	2438	2892	3345	3799	4252	4706	5160	5613	6067	6520
Ounces	7	198	652	1106	1559	2013	2466	2920	3374	3827	4281	4734	5188	5642	6095	6549
Ę۱	8	227	680	1134	1588	2041	2495	2948	3402	3856	4309	4763	5216	5670	6123	6577
ا ۲	9	255	709	1162	1616	2070	2523	2977	3430	3884	4337	4791	5245	5698	6152	6605
	10	283	737	1191	1644	2098	2551	3005	3459	3912	4366	4819	5273	5727	6180	6634
	11	312	765	1219	1673	2126	2580	3033	3487	3941	4394	4848	5301	5755	6209	6662
	12	340	794	1247	1701	2155	2608	3062	3515	3969	4423	4876	5330	5783	6237	6690
	13	369	822	1276	1729	2183	2637	3090	3544	3997	4451	4904	5358	5812	6265	6719
	14	397	850	1304	1758	2211	2665	3118	3572	4026	4479	4933	5386	5840	6294	6747
	15	425	879	1332	1786	2240	2693	3147	3600	4054	4508	4961	5415	5868	6322	6776

Pediatric Metric Conversion Tables

Temperature Fahrenheit (F) to Celsius (C)

°F	°C	°F	°C	°F	°C	°F	°C
95.0	35.0	98.0	36.7	101.0	38.3	104.0	40.0
95.2	35.1	98.2	36.8	101.2	38.4	104.2	40.1
95.4	35.2	98.4	36.9	101.4	38.6	104.4	40.2
95.6	35.3	98.6	37.0	101.6	38.7	104.6	40.3
95.8	35.4	98.8	37.1	101.8	38.8	104.8	40.4
96.0	35.6	99.0	37.2	102.0	38.9	105.0	40.6
96.2	35.7	99.2	37.3	102.2	39.0	105.2	40.7
96.4	35.8	99.4	37.4	102.4	39.1	105.4	40.8
96.6	35.9	99.6	37.6	102.6	39.2	105.6	40.9
96.8	36.0	99.8	37.7	102.8	39.3	105.8	41.0
97.0	36.1	100.0	37.8	103.0	39.4	106.0	41.1
97.2	36.2	100.2	37.9	103.2	39.6	106.2	41.2
97.4	36.3	100.4	38.0	103.4	39.7	106.4	41.3
97.6	36.4	100.6	38.1	103.6	39.8	106.6	41.4
97.8	36.6	100.8	38.2	103.8	39.9	106.8	41.6

Note: ${}^{\circ}C = ({}^{\circ}F-32) \times 5/9$. Celsius temperature equivalents rounded to one decimal place by adding 0.1 when second decimal place is 5 or greater.

The metric system replaces the term "Centigrade" with "Celsius" (the name of the inventor of the scale).

Length/Height Feet and inches to centimeters

Example: To obtain centimeters equivalent to 3 feet, 9 inches, read "3" on the top scale, "9" on the side scale; equivalent is 114.3 centimeters.

Feet

	0	1	2	3	4	5
0	0	30.5	61.0	91.4	121.9	152.4
1	2.5	33.0	63.5	94.0	124.5	154.9
2	5.1	35.6	66.0	96.5	127.0	157.5
3	7.6	38.1	68.6	99.1	129.5	160.0
4	10.2	40.6	71.1	101.6	132.1	162.6
5	12.7	43.2	73.7	104.1	134.6	165.1
6	15.2	45.7	76.2	106.7	137.2	167.6
7	17.8	48.3	78.7	109.2	139.7	170.2
8	20.3	50.8	81.3	111.8	142.2	172.7
9	22.9	53.3	83.8	114.3	144.8	175.3
10	25.4	55.9	86.4	116.8	147.3	177.8
11	27.9	58.4	88.9	119.4	149.9	180.3

Note: 1 inch = 2.540 centimeters. Centimeter equivalents rounded to one decimal place by adding 0.1 when second decimal place is 5 or greater; for example, 102.37 becomes 102.4.

Pounds	0	1	2	3	4	5	6	7	8	9
0	0.00	0.45	0.91	1.36	1.81	2.27	2.72	3.18	3.63	4.08
10	4.54	4.99	5.44	5.90	6.35	6.80	7.26	7.71	8.16	8.62
20	9.07	9.53	9.98	10.43	10.89	11.34	11.79	12.25	12.70	13.15
30	13.61	14.06	14.51	14.97	15.42	15.88	16.33	16.78	17.24	17.69
40	18.14	18.60	19.05	19.50	19.96	20.41	20.87	21.32	21.77	22.23
50	22.68	23.13	23.59	24.04	24.49	24.95	25.40	25.85	26.31	26.76
60	27.22	27.67	28.12	28.58	29.03	29.48	29.94	30.39	30.84	31.30
70	31.75	32.21	32.66	33.11	33.57	34.02	34.47	34.93	35.38	35.83
80	36.29	36.74	37.19	37.65	38.10	38.56	39.01	39.46	39.92	40.37
90	40.82	41.28	41.73	42.18	42.64	43.09	43.54	44.00	44.45	44.91
100	45.36	45.81	46.27	46.72	47.17	47.63	48.08	48.53	48.99	49.44
110	49.90	50.35	50.80	51.26	51.71	52.16	52.62	53.07	53.52	53.98
120	54.43	54.88	55.34	55.79	56.25	56.70	57.15	57.61	58.06	58.51
130	58.97	59.42	59.87	60.33	60.78	61.23	61.69	62.14	62.60	63.05
140	63.50	63.96	64.41	64.86	65.32	65.77	66.22	66.68	67.13	67.59
150	68.04	68.49	68.95	69.40	69.85	70.31	70.76	71.21	71.67	72.12
160	72.57	73.03	73.48	73.94	74.39	74.84	75.30	75.75	76.20	76.66
170	77.11	77.56	78.02	78.47	78.93	79.38	79.83	80.29	80.74	81.19
180	81.65	82.10	82.55	83.01	83.46	83.91	84.37	84.82	85.28	85.73
190	86.18	86.64	87.09	87.54	88.00	88.45	88.90	89.36	89.81	90.26
200	90.72	91.17	91.63	92.08	92.53	92.99	93.44	93.89	94.35	94.80

Weight (Mass) Pounds to kilograms

Example: To obtain kilograms equivalent to 44 pounds, read "40" on side scale, "4" on top scale; equivalent is 19.96 kilograms.

Note: 1 pound = 0.453592 kilogram. Kilogram equivalents have been rounded to 2 decimal places by adding 0.01 when the third decimal place is 5 or greater.

Abbott Infant Products: Approximate Powder Container Yield

Formula	Container size	Grams of powder per container	Grams of powder per scoop	Approximate container yield at standard caloric density	Approximate calories per container ¹	Pediatric approximate scoops per container ²
EleCare® for Infants	14.1 oz	400 g	9.4	95 fl oz	1900 Calories	42
Pure Bliss® by Similac®	24.7 oz	700 g	8.3	189 fl oz	3591 Calories	84
C	20.6 oz	584 g	8.6	151 fl oz	3020 Calories	67
Similac® 360 Total Care®	30.3 oz	859 g	8.6	222 fl oz	4440 Calories	99
360 lotal Care®	30.8 oz	874 g	8.6	226 fl oz	4520 Calories	101
Similac®	20.1 oz	570 g	8.8	145 fl oz	2900 Calories	64
360 Total Care®	29.5 oz	837 g	8.8	213 fl oz	4260 Calories	95
Sensitive	30.2 oz	857 g	8.8	218 fl oz	4360 Calories	97
	12.4 oz	352 g	8.7	90 fl oz	1800 Calories	40
Similac® Advance®	1.28 lb (20.6 oz)	584 g	8.7	149 fl oz	2980 Calories	67
Similac Advance	1.92 lb (30.8 oz)	874 g	8.7	224 fl oz	4480 Calories	100
	2.5 lb (40 oz)	1134g	8.7	290 fl oz	5800 Calories	130
Similac® Advance® Lamehadrin	1.54 lb (24.7 oz)	700 g	8.8	180 fl oz	3600 Calories	79
Similac®	12.1 oz	343 g	8.7	87 fl oz	1740 Calories	39
Alimentum® with 2'-FL HMO	19.8 oz	561 g	8.7	143 fl oz	2860 Calories	64
Similac® NeoSure®	13.1 oz	371 g	9.6	87 fl oz	1914 Calories	38
Similac Neosure	22.8 oz	646 g	9.6	151 fl oz	3322 Calories	67
Similac® Organic	1.28 lb (20.6 oz)	584 g	8.6	152 fl oz	3040 Calories	67
Similac® Organic with A2 Milk	1.28 lb (20.6 oz)	584 g	8.6	152 fl oz	3040 Calories	67
Similac® PM 60/40	14.1 oz	400 g	8.7	102 fl oz	2040 Calories	45

Formula	Container size	Grams of powder per container	Grams of powder per scoop	Approximate container yield at standard caloric density	Approximate calories per container ¹	Pediatric approximate scoops per container ²
	1.28 lb (20.6 oz)	584 g	8.6	151 fl oz	3020 Calories	67
Similac	1.92 lb (30.8 oz)	874 g	8.6	226 fl oz	4520 Calories	101
Pro-Advance®	2.12 lb (34 oz)	964 g	8.6	249 fl oz	4980 Calories	112
	2.25 lb (36 oz)	1021 g	8.6	264 fl oz	5280 Calories	118
0. "	1.25 lb (20.1 oz)	570 g	8.8	145 fl oz	2900 Calories	64
Similac Pro-Sensitive®	1.86 lb (29.8 oz)	845 g	8.8	215 fl oz	4300 Calories	96
1 10-Octionive	2.12 lb (34 oz)	964 g	8.8	245 fl oz	4900 Calories	109
0: "	1.25 lb (20.1 oz)	570 g	8.8	144 fl oz	2880 Calories	64
Similac Pro-Total Comfort®	1.86 lb (29.8 oz)	845 g	8.8	214 fl oz	4280 Calories	96
TTO TOTAL CONTION	2.25 lb (36 oz)	1021 g	8.8	258 fl oz	5160 Calories	116
	12.5 oz	354 g	8.8	90 fl oz	1800 Calories	40
Similac	1.25 lb (20.1 oz)	570g	8.8	145 fl oz	2900 Calories	64
Sensitive®	1.86 lb (29.8 oz)	845 g	8.8	215 fl oz	4300 Calories	96
	2.13 lb (34 oz)	964 g	8.8	245 fl oz	4900 Calories	109
Similac®	12.4 oz	352 g	8.8	90 fl oz	1800 Calories	40
Soy Isomil®	1.92 lb (30.8 oz)	873 g	8.8	223 fl oz	4460 Calories	99
Similac [®] Total Comfort	12.6 oz	357 g	8.8	90 fl oz	1800 Calories	40

^{&#}x27;Approximate calories per container derived from multiplying the fluid ounce (fl oz) container yield by the standard caloric density of that product (20 Cal/fl oz; exceptions: Pure Bliss by Similac at 19 Cal/fl oz and Similac NeoSure at 22 Cal/fl oz).

²Approximate scoops per container derived from dividing the grams of powder per container by the grams of powder per scoop; all numbers are rounded down to the nearest whole scoop (e.g. if 42.7, rounded to 42 scoops).

Macronutrient Values for Pediatric Formula Powders

All values are approximated; rounded from more precise numbers

Nutrients per Scoop

Measurements are based upon unpacked, level scoops. Use only the scoop provided in the container. Scoops from different products are not interchangeable.

Product	Grams of powder	Calories	Carbohydrate	Protein	Fat
EleCare® (for Infants)	9.4 g	44.7 Cal	4.78 g	1.38 g	2.14 g
EleCare® Jr (unflavored & vanilla)	9.5 g	44.6 Cal	4.68 g	1.36 g	2.16 g
EleCare® Jr (banana)	9 g	42.2 Cal	4.44 g	1.29 g	2.04 g
EleCare® Jr (chocolate)	8.9 g	41.7 Cal	4.39 g	1.27 g	2.02 g
Pure Bliss™ by Similac®	8.3 g	42.5 Cal	4.55 g	0.88 g	2.38 g
Similac®Advance®	8.7 g	44.5 Cal	4.77 g	0.92 g	2.50 g
Similac® Advance® Lamehadrin	8.8 g	45.2 Cal	5.06 g	0.96 g	2.35 g
Similac® Alimentum® with 2'-FL HMO	8.7 g	44.3 Cal	4.52 g	1.22 g	2.45 g

Nutrients per Scoop — cont'd					
Product	Grams of powder	Calories	Carbohydrate	Protein	Fat
Similac® NeoSure®	9.6 g	49.3 Cal	4.98 g	1.38 g	2.71 g
Similac® Organic	8.6 g	44.7 Cal	4.65 g	0.93 g	2.52 g
Similac® Organic with A2 Milk	8.6 g	44.7 Cal	4.65 g	0.93 g	2.52 g
Similac® PM 60/40	8.7 g	44.7 Cal	4.56 g	0.98 g	2.50 g
Similac Pro-Advance®	8.6 g	44.5 Cal	4.67 g	0.92 g	2.49 g
Similac Pro-Sensitive®	8.8 g	44.7 Cal	4.88 g	0.94 g	2.42 g
Similac Pro-Total Comfort®	8.8 g	44.5 Cal	4.76 g	1.03 g	2.40 g
Similac® 360 Total Care®	8.6 g	44.5 Cal	4.67 g	0.92 g	2.49 g
Similac® 360 Total Care® Sensitive	8.8 g	44.7 Cal	4.88 g	0.94 g	2.42 g
Similac Sensitive®	8.8 g	44.8 Cal	4.88 g	0.94 g	2.42 g
Similac® Soy Isomil®	8.8 g	44.9 Cal	4.67 g	1.1 g	2.45 g
Similac Total Comfort®	8.8 g	44.5 Cal	4.77 g	1.03 g	2.41 g

Nutrients per Packet:						
Product	Packet Size	Calories	Carbohydrate	Protein	Fat	Displacement
Similac® Human Milk Fortifier Concentrated Liquid	5 mL	7 Cal	0.81 g	0.35 g	0.27 g	5 mL
Similac® Human Milk Fortifier Hydrolyzed Protein Concentrated Liquid	5 mL	7 Cal	0.75 g	0.5 g	0.21 g	5 mL

Nutrients per 100 Grams:					
Product	Grams of powder	Calories	Carbohydrate	Protein	Fat
EleCare (for Infants)	9.4 g	475 Cal	50.85 g	14.73 g	22.81 g
EleCare Jr (unflavored & vanilla)	9.5 g	469 Cal	49.3 g	14.3 g	22.7 g
EleCare Jr (banana)	9 g	469 Cal	49.3 g	14.3 g	22.7 g
EleCare Jr (chocolate)	8.9 g	469 Cal	49.3 g	14.3 g	22.7 g
Pure Bliss by Similac	8.3 g	512 Cal	54.79 g	10.6 g	28.67 g
Similac Advance	8.7 g	512 Cal	54.79 g	10.6 g	28.67 g
Similac Advance Lamehadrin	8.8 g	514 Cal	57.52 g	10.89 g	26.71 g
Similac Alimentum with 2'-FL HMO	8.7 g	509 Cal	51.9 g	13.99 g	28.19 g
Similac NeoSure	9.6 g	513 Cal	51.85 g	14.37 g	28.23 g
Similac Organic	8.6 g	520 Cal	54.05 g	10.76 g	29.26 g
Similac Organic with A2 Milk	8.6 g	520 Cal	54.05 g	10.76 g	29.26 g
Similac PM 60/40	8.7 g	514 Cal	52.38 g	11.3 g	28.76 g
Similac Pro-Advance	8.6 g	517 Cal	54.32 g	10.71 g	28.97 g
Similac Pro-Sensitive	8.8 g	508 Cal	55.41 g	10.68 g	27.45 g
Similac Pro-Total Comfort	8.8 g	506 Cal	54.12 g	11.73 g	27.31 g
Similac 360 Total Care	8.6 g	517 Cal	54.32 g	10.71 g	28.97 g
Similac 360 Total Care Sensitive	8.8 g	508 Cal	55.41 g	10.68 g	27.45 g
Similac Sensitive	8.8 g	509 Cal	55.47 g	10.69 g	27.48 g
Similac Soy Isomil	8.8 g	510 Cal	53.09 g	12.51 g	27.87 g
Similac Total Comfort	8.8 g	506 Cal	54.18 g	11.75 g	27.34 g

Approximate Displacement			
Product	Powder grams per scoop	Per scoop	Per gram of powder
EleCare® (for Infants)	9.4 g	7.0 mL	0.74 mL
EleCare® Jr (unflavored & vanilla)	9.5 g	7.0 mL	0.74 mL
EleCare® Jr (banana)	9 g	6.7 mL	0.74 mL
EleCare® Jr (chocolate)	8.9 g	6.6 mL	0.74 mL
Pure Bliss™ by Similac®	8.3 g	6.3 mL	0.77 mL
Similac® Advance®	8.7 g	6.7 mL	0.77 mL
Similac® Advance® Lamehadrin	8.8 g	6.8 mL	0.77 mL
Similac® Alimentum®	8.7 g	6.6 mL	0.76 mL
Similac® NeoSure®	9.6 g	7.3 mL	0.76 mL
Similac® Organic	8.6 g	6.6 mL	0.77 mL
Similac® Organic with A2 Milk	8.6 g	6.6 mL	0.77 mL
Similac® PM 60/40	8.7 g	6.9 mL	0.79 mL
Similac Pro-Advance®	8.6 g	6.6 mL	0.77 mL
Similac Pro-Sensitive®	8.8 g	6.8 mL	0.78 mL
Similac Pro-Total Comfort®	8.8 g	6.8 mL	0.78 mL
Similac Sensitive®	8.8 g	6.8 mL	0.78 mL
Similac® Soy Isomil®	8.8 g	6.7 mL	0.77 mL
Similac Total Comfort®	8.8 g	6.8 mL	0.77 mL
Similac® 360 Total Care®	8.6 g	6.6 mL	0.77mL
Similac® 360 Total Care® Sensitive	8.8 g	6.7 mL	0.76 mL

Item Number	Product	Flavor	Unit	HCPCS Code	NDC Format Code
Term Infant Form	ulas				
68136	Similac® 360 Total Care® (RTF)		2 fl oz bottle		70074-0681-36
68145	Similac® 360 Total Care® (RTF)		1 qt bottle		70074-0681-46
68063	Similac® 360 Total Care® (Powder)		20.6 oz container		70074-0680-64
68065	Similac® 360 Total Care® (Powder)		30.3 oz container		70074-0680-66
68116	Similac® 360 Total Care® (Powder)		30.8 oz can		70074-0681-17
68111	Similac® 360 Total Care® (Powder)		36 oz container		70074-0681-12
68067	Similac® 360 Total Care® (Powder)		40 oz container		70074-0680-68
67323	Similac Pro-Advance® (RTF)		2 fl oz bottle	B4158	70074-0673-24
64248	Similac Pro-Advance® (RTF)		1 qt bottle	B4158	70074-0642-49
68088	Similac Pro-Advance® (Powder)		20.6 oz container	B4158	70074-0680-89
66439	Similac Pro-Advance® (Powder)		30.8 oz container	B4158	70074-0664-38
66655	Similac Pro-Advance® (Powder)		34 oz container	B4158	70074-0666-56
66781	Similac Pro-Advance® (Powder) 0.61 oz		16 stick packs/case	B4158	70074-0667-83
53363	Similac® Advance® (RTF)		32 fl oz bottle	B4158	70074-0533-64
56973	Similac® Advance® (CL)		13 fl oz can	B4158	70074-0569-74
55957	Similac® Advance® (Powder)		12.4 oz can	B4158	70074-0559-58
68084	Similac® Advance® (Powder)		20.6 oz container	B4158	70074-0680-85
58585	Similac® Advance® (Powder)		30.8 oz container	B4158	70074-0585-86
68092	Similac® Organic (Powder)		20.6 oz container	B4158	70074-0680-93
67391	Similac® Soy Isomil®		2 fl oz bottle	B4159	70074-0673-92
67012	Similac® Soy Isomil® (RTF)		32 fl oz bottle	B4159	70074-0559-68
56975	Similac® Soy Isomil® (CL)		13 fl oz can	B4159	70074-0569-76
55963	Similac® Soy Isomil® (Powder)		12.4 oz can	B4159	70074-0559-64
67699	Similac® Soy Isomil®		30.8 oz container	B4159	70074-0677-00

Item Number	Product	Flavor	Unit	HCPCS Code	NDC Format Code
68155	Similac® 360 Total Care® Sensitive (RTF)		2 fl oz bottle		70074-0681-55
68160	Similac® 360 Total Care® Sensitive (RTF)		1 qt bottle		70074-0681-61
68070	Similac® 360 Total Care® Sensitive		20.1 oz container		70074-0680-71
68072	Similac® 360 Total Care® Sensitive		29.5 oz container		70074-0680073
68120	Similac® 360 Total Care® Sensitive		34.9 oz container		70074-0681-21
68074	Similac® 360 Total Care® Sensitive		40 oz container		70074-0680-75
67325	Similac Pro-Sensitive® (RTF)		2 fl oz bottle	B4158	70074-0673-26
67517	Similac Pro-Sensitive® (RTF)		8 fl oz bottle	B4158	70074-0675-19
64253	Similac Pro-Sensitive® (RTF)		32 fl oz bottle	B4158	70074-0642-54
68090	Similac Pro-Sensitive® (Powder)		20.1 oz container	B4158	70074-0680-91
66441	Similac Pro-Sensitive® (Powder)		29.8 oz container	B4158	70074-0664-40
57535	Similac Sensitive® (CL)		13 fl oz can	B4158	70074-0575-36
57533	Similac Sensitive® (RTF)		1 qt bottle	B4158	70074-0575-34
57539	Similac Sensitive® (Powder)		12.5 oz can	B4158	70074-0575-41
68082	Similac Sensitive® (Powder)		20.1 oz container	B4158	70074-0680-83
62951	Similac Sensitive® (Powder)		29.8 oz container	B4158	70074-0629-52
64440	Similac Sensitive® (Powder)		34.9 oz container	B4158	70074-0662-86
67132	Similac Pro-Total Comfort®		2 fl oz bottle	N/A	70074-0671-33
68107	Similac Pro-Total Comfort®		20.1 oz container	N/A	70074-0681-08
66937	Similac Pro-Total Comfort®		29.8 oz can	N/A	70074-0669-38
62599	Similac Total Comfort® (Powder)		12.6 oz can	B4158	70074-0626-00
59738	Similac® Alimentum® (RTF)		2 fl oz bottle	B4161	70074-0111-30
57512	Similac® Alimentum® (RTF)		1 qt bottle	B4161	70074-0575-13
64715	Similac® Alimentum® (Powder)		12.1 oz can	B4161	70074-0647-12
64719	Similac® Alimentum® (Powder)		19.8 oz can	B4161	70074-0647-18

Item Number	Product	Flavor	Unit	HCPCS Code	NDC Format Code			
55251	EleCare® (for infants)(Powder)	Unflavored	14.1	B4161	70074-0535-11			
00850	Similac® PM 60/40 (Powder)		14.1 oz can	B4154	70074-0608-50			
Premature Infant	Premature Infant Formulas							
67366	Similac® Special Care® 20 (RTF)		2 fl oz bottle	N/A	70074-0673-67			
67368	Similac® Special Care® 24 (RTF) (Institutional)		2 fl oz bottle	B4160	70074-0676-87			
67446	Similac® Special Care® 24 (RTF) (Retail)		2 fl oz bottle	B4160	70074-0114-85			
67370	Similac® Special Care® 24 High Protein (RTF)		2 fl oz bottle	N/A	70074-0673-71			
67372	Similac® Special Care® 30 (RTF)		2 fl oz bottle	B4160	70074-0676-89			
67448	Similac® Special Care® 30 (RTF) (Retail)		2 fl oz bottle	B4160	70074-0114-86			
56177	Similac® NeoSure® (RTF)		2 fl oz bottle	B4160	70074-0561-78			
57455	Similac® NeoSure® (RTF)		1 qt bottle	B4160	70074-0574-56			
57430	Similac® NeoSure® (Powder)		13.1 oz can	B4160	70074-0574-31			
66923	Similac® NeoSure® (Powder)		22.8 oz can	B4160	70074-0669-24			
Human Milk Fort	Human Milk Fortification							
56649	Similac® Human Milk Fortifier Concentrated Liquid (Institutional)		5. mL packet	B4155	70074-0566-50			
62317	Liquid Protein Fortifier		1.8 fl oz bottle	N/A	70074-0623-15			
Children								
55253	EleCare® Jr (Powder)	Unflavored	14.1	B4161	70074-0552-54			
56585	EleCare® Jr (Powder)	Vanilla	14.1	B4161	70074-0565-86			
66273	EleCare® Jr (Powder)	Chocolate	14.1	B4161	70074-0662-71			
66275	EleCare® Jr (Powder)	Banana	14.1	B4161	70074-0662-76			
67522	PediaSure® (Oral Use Instititutional)	Vanilla	8 fl oz can	B4160	70074-0671-85			
67523	PediaSure® (Oral Use Instititutional)	Chocolate	8 fl oz can	B4160	70074-0675-24			
67525	PediaSure® (Oral Use Instititutional)	Strawberry	8 fl oz can	B4160	70074-0675-26			
67527	PediaSure® (Oral Use Instititutional)	Banana	8 fl oz can	B4160	70074-0675-28			
67533	PediaSure® (Oral Use Instititutional)	Vanilla	8 fl oz bottle	B4160	70074-0675-34			

Item Number	Product	Flavor	Unit	HCPCS Code	NDC Format Code
67535	PediaSure® (Oral Use Instititutional)	Chocolate	8 fl oz bottle	B4160	70074-0675-36
67537	PediaSure® (Oral Use Instititutional)	Strawberry	8 fl oz bottle	B4160	70074-0675-38
58049	PediaSure® Retail	Vanilla	8 fl oz bottle	B4160	70074-0580-51
58058	PediaSure® Retail	Chocolate	8 fl oz bottle	B4160	70074-0580-60
58055	PediaSure® Retail	Strawberry	8 fl oz bottle	B4160	70074-0580-57
58052	PediaSure® Retail	Banana	8 fl oz bottle	B4160	70074-0580-54
53818	PediaSure® Retail	Berry	8 fl oz bottle	B4160	70074-0538-20
66918	PediaSure® Retail	S'mores	8 fl oz bottle	B4160	70074-0669-20
67529	PediaSure® Grow & Gain with Fiber (Institutional)	Vanilla	8 fl oz can	B4160	70074-0675-30
67531	PediaSure® Grow & Gain with Fiber (Institutional)	Vanilla	8 fl oz bottle	B4160	70074-0675-32
58061	PediaSure® Grow & Gain with Fiber	Vanilla	8 fl oz bottle	B4160	70074-0580-63
56368	PediaSure® Grow & Gain with Fiber	Strawberry	8 fl oz bottle	B4160	70074-0563-69
67214	PediaSure® Grow & Gain Organic	Classic Vanilla	8 fl oz bottle	N/A	70074-0672-15
67217	PediaSure® Grow & Gain Organic	Milk Chocolate	8 fl oz bottle	N/A	70074-0672-18
67962	PediaSure Harvest™ (Institutional)	Unflavored	8 fl oz carton	B4149	70074-0679-63
67612	Pediasure® Reduced Calorie (Institutional)	Vanilla	8 fl oz bottle	B4158	70074-0676-13
66912	PediaSure SideKicks®	Vanilla	8 fl oz bottle	N/A	70074-0669-13
66916	PediaSure SideKicks®	Chocolate	8 fl oz bottle	N/A	70074-0669-17
66639	PediaSure SideKicks®	Strawberry	8 fl oz bottle	N/A	70074-0666-40
67401	PediaSure® Enteral Formula 1.0 Cal (Enteral Formula Only - Institutional)	Vanilla	8 fl oz can	B4160	70074-0674-02
67403	PediaSure® Enteral Formula 1.0 Cal with Fiber (Enteral Formula Only - Institutional)	Vanilla	8 fl oz can	B4160	70074-0674-04
67405	PediaSure® Enteral Formula 1.0 Cal with Fiber RTH (Institutional)	Vanilla	1 liter bottle	B4160	70074-0674-06
67378	PediaSure® 1.5 Cal (Oral Use Instititutional)	Vanilla	8 fl oz can	B4160	70074-0673-86

Item Number	Product	Flavor	Unit	HCPCS Code	NDC Format Code
67374	PediaSure® 1.5 Cal with Fiber (Oral Use Institutional)	Vanilla	8 fl oz can	B4160	70074-0673-75
67376	PediaSure® 1.5 Cal with Fiber (Institutional)	Vanilla	1 liter bottle	B4160	70074-0673-77
67413	PediaSure® Peptide 1.0 Cal (Institutional)	Unflavored	8 fl oz bottle	B4161	70074-0674-14
67407	PediaSure® Peptide 1.0 Cal (Institutional)	Vanilla	8 fl oz bottle	B4161	70074-0674-10
67411	PediaSure® Peptide 1.0 Cal (Institutional)	Strawberry	8 fl oz bottle	B4161	70074-0674-12
67415	PediaSure® Peptide 1.0 Cal RTH (Institutional)	Unflavored	1 liter bottle	B4161	70074-0674-16
67417	PediaSure® Peptide 1.5 Cal (Institutional)	Vanilla	8 fl oz bottle	B4161	70074-0674-18
67419	PediaSure® Peptide 1.5 Cal RTH (Institutional)	Vanilla	1 liter bottle	B4161	70074-0674-20
66236	Vital® Peptide 1.5 Cal (Institutional)	Vanilla	8 fl oz carton	B4161	70074-0662-37
67151	Go & Grow by Similac® Non-GMO with 2'-FL HMO Toddler Drink		24 oz can	N/A	70074-0671-52
67011	Go & Grow by Similac® Toddler Drink		1.93 lb can	N/A	70074-0663-85
67581	Go & Grow by Similac® Sensitive Non-GMO with 2'-FL HMO Toddler Drink		23.2 oz container	N/A	70074-0647-88
Therapeutic Hyd	ration				
00336	Pedialyte® (Solution)	Unflavored	1 liter bottle	B4103	70074-0803-36
00240	Pedialyte® (Solution)	Grape	1 liter bottle	B4103	70074-0802-40
00365	Pedialyte® (Solution)	Mixed Fruit	1 liter bottle	B4103	70074-0803-65
53983	Pedialyte® (Solution)	Strawberry	1 liter bottle	B4103	70074-0539-84
67461	Pedialyte® (Solution)	Coconut Burst	1 liter bottle	B4103	70074-0674-62
63059	Pedialyte AdvancedCare®	Blue Raspberry	1 liter bottle	B4103	70074-0630-60
63057	Pedialyte AdvancedCare®	Cherry Punch	1 liter bottle	B4103	70074-0630-58
64301	Pedialyte AdvancedCare®	Strawberry Lemonade	1 liter bottle	B4103	70074-0643-02
64307	Pedialyte AdvancedCare®	Tropical Fruit	1 liter bottle	B4103	70074-0643-08
66641	Pedialyte AdvancedCare® Plus	Berry Frost	1 liter bottle	N/A	70074-0666-42

Item Number	Product	Flavor	Unit	HCPCS Code	NDC Format Code
66645	Pedialyte AdvancedCare® Plus	Chilled Cherry Pomegranate	1 liter bottle	N/A	70074-0666-46
66855	Pedialyte AdvancedCare® Plus	Iced Grape	1 liter bottle	N/A	70074-0668-56
66643	Pedialyte AdvancedCare® Plus	Kiwi Berry Mist	1 liter bottle	N/A	70074-0666-44
67434	Pedialyte AdvancedCare® Plus	Orange Breeze	1 liter bottle	N/A	70074-0674-35
Metabolic Produ	cts				
53328	Calcilo XD®		13.2 oz can	B4162	70074-0533-29
67032	Cyclinex®-1		14.1 oz can	B4162	70074-0670-33
67034	Cyclinex®-2		14.1 oz can	B4157 or B4162	70074-0670-35
67036	Glutarex®-1		14.1 oz can	B4162	70074-0670-37
67038	Glutarex®-2		14.1 oz can	B4157 or B4162	70074-0670-39
67040	Hominex®-1		14.1 oz can	B4162	70074-0670-41
67042	Hominex®-2		14.1 oz can	B4157 or B4162	70074-0670-43
67044	I-Valex®-1		14.1 oz can	B4162	70074-0670-45
67046	I-Valex®-2		14.1 oz can	B4157 or B4162	70074-0670-47
67048	Ketonex®-1		14.1 oz can	B4162	70074-0670-49
67050	Ketonex®-2		14.1 oz can	B4157 or B4162	70074-0670-51
67052	Phenex®-1		14.1 oz can	B4162	70074-0670-53
67054	Phenex®-2	Unflavored	14.1 oz can	B4157 or B4162	70074-0670-55
67056	Phenex®-2	Vanilla	14.1 oz can	B4157 or B4162	70074-0670-57
67030	Pro-Phree®		14.1 oz can	B4155	70074-0670-31
67058	Propimex®-1		14.1 oz can	B4162	70074-0670-59

Item Number	Product	Flavor	Unit	HCPCS Code	NDC Format Code
67060	Propimex®-2		14.1 oz can	B4157 or B4162	70074-0670-61
50260	ProViMin®		5.3 oz can	B4155	70074-0502-61
00108	RCF® (Liquid)		13 fl oz can	B4155	70074-0401-08
67062	Tyrex®-1		14.1 oz can	B4162	70074-0670-63
67064	Tyrex®-2		14.1 oz can	B4157 or B4162	70074-0670-65

Abbott does not represent these codes to be actual National Drug Codes (NDCs). NDC-format codes are product codes adjusted according to standard industry practice to meet the format requirements of pharmacy and health insurance computer systems.

NOTE: The HCPCS codes provided in this book are taken from government publications and are provided from information correct at the time of publication. They are provided for your information only. Healthcare providers are ultimately responsible for making appropriate product selections per individual patient and verifying that codes used for third-party billing are accurate for the items provided.

